



Field Trip Guide Book - B22

Florence - Italy
August 20-28, 2004

Volume n° 2 - from B16 to B33

**32nd INTERNATIONAL
GEOLOGICAL CONGRESS**

**PALEOZOIC OROGENIES IN THE
FRENCH MASSIF CENTRAL
A CROSS SECTION FROM
BÉZIERS TO LYON**



Leader: M. Faure

Associate Leaders: J.M. Lardeaux, P. Matte

Pre-Congress

B22

The scientific content of this guide is under the total responsibility of the Authors

Published by:

**APAT – Italian Agency for the Environmental Protection and Technical Services - Via Vitaliano
Brancati, 48 - 00144 Roma - Italy**



Series Editors:

Luca Guerrieri, Irene Rischia and Leonello Serva (APAT, Roma)

English Desk-copy Editors:

**Paul Mazza (Università di Firenze), Jessica Ann Thonn (Università di Firenze), Nathalie Marlène
Adams (Università di Firenze), Miriam Friedman (Università di Firenze), Kate Eadie (Freelance
independent professional)**

Field Trip Committee:

**Leonello Serva (APAT, Roma), Alessandro Michetti (Università dell'Insubria, Como), Giulio Pavia
(Università di Torino), Raffaele Pignone (Servizio Geologico Regione Emilia-Romagna, Bologna) and
Riccardo Polino (CNR, Torino)**

Acknowledgments:

**The 32nd IGC Organizing Committee is grateful to Roberto Pompili and Elisa Brustia (APAT, Roma)
for their collaboration in editing.**

Graphic project:

Full snc - Firenze

Layout and press:

Lito Terrazzi srl - Firenze

Volume n° 2 - from B16 to B33



**32nd INTERNATIONAL
GEOLOGICAL CONGRESS**

**PALEOZOIC OROGENIES IN THE
FRENCH MASSIF CENTRAL
A CROSS SECTION FROM
BÈZIERS TO LYON**

AUTHORS:

M. Faure (University of Orléans - France)

P. Ledru (BRGM, Orléans - France)

J.M. Lardeaux (University of Nice - France)

P. Matte (CNRS, University of Montpellier - France)

**Florence - Italy
August 20-28, 2004**

Pre-Congress

B22

Front Cover:

View of gneiss-migmatite of the Montagne Noire axial zone (visited during D2) looking to the north. The village of Olargues is located in the micaschist envelope of the dome. The picture is taken from the northernmost part of the recumbent folds of the Montagne Noire southern side (visited during D1).

Leader: M. Faure

Associate Leaders: J.M. Lardeaux, P. Ledru, P. Matte


1. Introduction


The formation of the continental substratum of Medio-Europa occurred in Paleozoic times. The names of “Hercynian” or “Variscan” are used to deal with the geodynamic processes that took place from Cambrian to Carboniferous. It is now widely accepted that this Paleozoic Belt that crops out from Iberia to Bohemia (Fig. 1) results from a complex interplay of rifting, convergence and collision between three large continents, namely Laurentia, Baltica and Gondwana and several microcontinental stripes such as Avalonia or Armorica (Matte, 2001). Continental drifting and welding resulted in the opening and closure of several oceans such as Iapetus, Rheic and Medio-European. There is however a wide range of opinions concerning the location and width of these oceanic domains and the number, kinematics and timing of collisional processes (e. g. Autran and Cogné, 1980; Franke, 1989, 2000; Ledru et al., 1989; Matte, 1991; 2001; Faure et al., 1997).

The French Massif Central is one of the largest pieces of the Variscan Belt. The whole Massif Central provides a reference cross section throughout the north Gondwana margin deformed and metamorphosed during the Paleozoic. During the last two decades, and recently through the GéoFrance 3D program, developments made in the areas of geochronology, structural geology, metamorphic and magmatic petrology, allow us to draw a comprehensive structural map of the Massif Central and to discuss a possible scenario accounting for the Paleozoic tectono-thermal evolution.

This field trip presents representative lithological, structural, magmatic, metamorphic and geochronological data of the French Massif Central from unmetamorphosed kilometer-scale recumbent folds to UHP metamorphic rocks. Most of the controversial aspects of collisional orogens such as continental subduction and exhumation

Figure 1 - Location of the French Massif Central in the frame of the Paleozoic belt of Medio-Europa (Matte, 1991).





of ultrametamorphic rocks, nappe kinematics, inverted metamorphism, syn- to post-orogenic extensional tectonics, crustal melting and tectonic setting of pluton emplacement will be addressed.

Field References

Topographic maps IGN 1/100 000: n°65 Béziers-Montpellier; n°58 Rodez-Mende; n°59 Privas-Alès; n°50 St-Etienne-Le Puy; n°51 Lyon-Grenoble. Geologic maps BRGM 1/50 000: n°1014 St-Chinian; n°988 Bédarieux; n°862 Mende; n°863 Le Bleymard; n°839 Langogne; n°840 Burzet; n°792 Yssingeaux; n°745 St-Etienne; n°721 St-Symphorien-sur-Coise; n°697 Tarare.

2. Regional geological setting

2.1. A structural map of the French Massif Central.

It is now widely accepted that the structure of the French Massif Central is a stack of nappes (Ledru et al., 1989, 1994 Fig. 3). From top to bottom and also from south to north, six main tectonic units are distinguished.

i) The Southern Palaeozoic Fold and Thrust Belt involves a set of continental margin/platform series recording a more or less continuous sedimentation spanning from Early Cambrian to Early Carboniferous. The series is deformed within kilometer-scale recumbent folds well observed in the Montagne Noire area (Arthaud, 1970).

ii) The Para-autochthonous Unit that overthrusts the previous unit consists of a thick metapelite-metagrauwacke series (also called “Cévennes micaschists”) with some quartzite beds and volcanic rocks. Although stratigraphic ages are lacking, a Neoproterozoic to Ordovician age is

Figure 2 - Cross-sections from the Massif Armoricaïn to Ardenne (B) and through Massif Central (C). C' is an alternative section through the Lèvezou klippe (Matte, 1991).



Figure 3 - Structural map of the Massif Central
(adapted from Ledru et al., 1989).

generally accepted.

iii) The Lower Gneiss Unit (LGU) is lithologically quite similar to the Para-autochthonous Unit. Early Cambrian and Early Ordovician alkaline granitoids, now transformed in augen orthogneiss, are also widespread. Both the Para-autochthonous Unit and Lower Gneiss Unit are interpreted as Proterozoic-Early Paleozoic remnants of the northern Gondwana margin that experienced crustal thinning and rifting in Ordovician times.

iv) The Upper Gneiss Unit (UGU) is made up of a bi-modal association called "leptynite-amphibolite" sequence which is a peculiar assemblage of mafic and felsic rocks. This unit experienced a higher metamorphic pressure under eclogite and HP granulite facies (ca. 20Kb). Ultra high-pressure

metamorphism is reached locally near Lyon, coesite-eclogite facies rocks crop-out (Lardeaux et al., 2001). The protoliths of the UGU also include metasediments and granitoids. The upper part of the UGU consists of migmatites formed by the partial melting of pelitic and quartzo-feldspathic rocks within which amphibolite block are preserved as restites. Radiometric dates show that the magmatism occurred in Early Ordovician times (ca. 480 Ma) and the high-pressure metamorphism in Late Silurian (ca. 420-410 Ma, Pin and Lancelot, 1982; Ducrot et al., 1983). Due to the occurrence of rare metagabbros and serpentized ultramafics, the UGU is considered by some authors as a remnant of an oceanic domain, the Medio-European Ocean, that opened in Early Paleozoic times during the rifting that led to the separation of Armorica from North Gondwana (e. g. Dubuisson et al., 1989; Matte, 1991). However, it is worth noting that the Upper Gneiss Unit is not a true ophiolitic sequence since oceanic sedimentary rocks such as radiolarites or siliceous

shales are lacking and ultramafics or serpentinites are rare. A likely interpretation would be to consider that the UGU is a transitional crust between true continental and oceanic ones.

v) The Thiviers-Payzac Unit that crops out in the south Limousin, is the highest tectonic unit of the allochthonous stack in the French Massif Central. It is formed by Cambrian metagraywackes, rhyolites and quartzites intruded by Ordovician granite. Conversely to the underlying UGU, the Thiviers-Payzac Unit never experienced the high-pressure metamorphism. As revealed by seismic reflection line (Bitri et al., 1999), these relatively low grade rocks tectonically overly the UGU.

vi) In the NE Massif Central, near Lyon, the Brevenne Unit consists of mafic magmatic rocks (pyroclastites, pillow basalts, diabases, gabbros), serpentized ultramafics, acidic volcanic rocks,


and siliceous sediments (radiolarites, siltites). The acidic rocks are dated of 366 ± 5 Ma (U/Pb method on zircon, Pin and Paquette, 1998). Petrology and geochemistry show that the Brevenne Unit and its extension in the Beaujolais area is a Middle Devonian oceanic sequence formed within an oceanic or a back-arc basin opened within the UGU (Sider and Ohnenstetter, 1986; Pin, 1990; Pin and Paquette, 1998). The Brevenne Unit records an early thrusting to the NW over the UGU followed by a NE-SW dextral strike-slip (Feybesse et al., 1988; Leloix et al., 1999). The precise age of the thrusting is unknown but since the metamorphic rocks are concealed below the Early Visean calcareous sandstone of the famous unconformity of Le Goujet (east of Lyon) an Early Carboniferous age is likely (see below).

2.2. The tectono-metamorphic evolution.

Structural information related to the high-pressure

metamorphism and the prograde metamorphic evolution is poorly documented since these rocks are known only as relics. It is therefore quite difficult to draw a general view of this event. Moreover, three main synmetamorphic ductile events are recognized. The earliest deformation found in the UGU, D1, is characterized by a NE-SW trending lineation with a top-to-the-SW shearing developed coevally with an intermediate pressure/intermediate temperature metamorphism and anatexis dated around 385-380 Ma (e. g. Floc'h, 1983; Quenardel and Rolin, 1984; Costa, 1992; Boutin and Montigny, 1993; Duthou et al., 1994; Roig and Faure, 2000; Figs. 4, 5). Since the D1 event is found in the migmatites that form the upper part of the UGU, it occurred during or at the end of the exhumation of the high-pressure metamorphic rocks. The radiometric dates comply with the Devonian stratigraphic age of the unmetamorphosed rocks (e. g. Villedé d'Ardin, Génis, Somme, Belfort

Figure 4 - Structural and geochronologic features related to the Late Silurian-Middle Devonian D1 event.






Figure 5 - P-T paths of the Silurian-Devonian and Carboniferous events for the different units.

areas, Fig. 4). Although a direct unconformity is never observed, field relationships suggest that D1 is older than Middle Devonian.

The second event, D2, is characterized by a NW-SE trending lineation coeval with a barrovian type metamorphism (Figs. 5, 6). $^{40}\text{Ar}/^{39}\text{Ar}$ dates on biotite, muscovite and amphibole range around 360-350 Ma. Most of the shear criteria developed along the NW-SE lineation indicate a top-to-the-NW shearing. In the Rouergue area, the Naucelle thrust is related to this event (Duguet and Faure, in press). The last increment of the ductile deformation in the metamorphic series is associated with the emplacement of peraluminous cordierite bearing granitoids such as the Guéret pluton that is the largest massif of this type. These granitoids exhibit magmatic to sub-solidus fabrics that comply with the synkinematic character of these plutons (e. g. Roig et al., 1998). A similar tectonic-metamorphic-magmatic pattern is also recognized in the south part of the Massif Armoricain. The closure of the Brevenne oceanic basins is chronologically and kinematically in agreement with the D2 event (Leloix et al., 1999). The geodynamic significance of the NW-SE lineation parallel to the belt is not clearly understood yet. Several hypotheses have been proposed (e. g. Burg et al., 1987; Bouchez and Jover, 1986; Mattauer et al., 1988) but none of them appears fully convincing. As discussed in section 2.4,

this Early Carboniferous deformation is coeval with the closure of the Rheic Ocean and collision between Gondwana and Laurussia.

The third event, D3, is restricted to the southern part of the Massif Central. In the Para-autochthonous Unit of Cévennes-Albigeois, upper greenschist to amphibolite facies rocks are deformed by top-to-the-south ductile shearing along a submeridian lineation (Fig. 7). Available $^{40}\text{Ar}/^{39}\text{Ar}$ dates on the metamorphic minerals yield Viséan ages around 340 Ma (Monié et al., 2000; Faure et al., 2001). This thrusting propagates southward in the Fold and Thrust Belt where kilometer-scale recumbent folds develop from Viséan to Namurian. Although in the South Massif Central south-directed compressional regime lasts from Viséan to Namurian (345 to 325Ma), conversely, in the northern part of the massif, the Late Viséan (ca. 340 Ma) is a turning point in the tectonic evolution. From Morvan to Limousin, the Late Viséan time corresponds to the onset of syn-orogenic extension characterized by a huge crustal melting. Structural studies indicate that the syn-orogenic extension is controlled by a NW-SE maximum stretching direction (Fig. 7). The NW-SE spreading of the inner part of the Massif Central is also partly accommodated by ductile wrench faults well developed in Limousin (e. g. La Courtine or S. Limousin faults, Fig. 7) and in the Massif Armoricain. In the scale of the whole



Figure 6 - Structural, magmatic and geochronologic features related to the Late Devonian-Early Carboniferous D2 event (AMBP: Magnetic Anomaly of Paris Basin).

belt, the Late Visean to Namurian compression is also responsible for the development of north-directed thrusts in Ardenne and SE England.

The last ductile deformation events (ca. 320 Ma and younger ones) took place during the collapse of the belt. Since they are closely associated to magmatism, they will be considered in the next section.

2.3. A magmatic outline.

Like all the Variscan massifs, the Massif Central is also characterized by a voluminous magmatism mainly derived from crustal melting. Several generations of migmatites and granitoids are recognized (e. g. Duthou et al., 1984).

2.3.1. The pre-orogenic magmatism is not presented in detail here. The Early Ordovician bimodal magmatism, responsible for the formation of the leptynite-amphibolite complex in the UGU, and the Cambrian or Ordovician magmatic rocks are ductilely deformed, metamorphosed and included in the stack of nappes.

2.3.2. The Middle to Late Devonian calc-alkaline volcanic and volcanoclastic rocks that crop out in the Morvan area (called the Somme series) belong to a magmatic arc (Fig. 8; Pin et al., 1982; Delfour, 1989). In the south part of the Massif Armoricain, Eifelian-Givetian basaltic pillow lavas form the Meilleraie series. These rocks are interpreted as the aerial part of a magmatic arc. Moreover, mafic calc-alkaline rocks well known for a long time in the Limousin (Didier and Lameyre, 1971), are interpreted as the deep part of the same Devonian arc. Its geodynamic significance will be discussed in section 2.4.

2.3.3. The Tournaisian late-collisional magmatism is represented by the Guéret-type granites peraluminous plutons. Their magmatic fabric suggests that those plutons emplacement was controlled by the

same strain field than the D2 deformation (Fig. 6) .

2.3.4. The Visean magmatism is well developed in the north and west part of the massif Central (Fig. 7). It consists in aerial products with lava flows, ignimbrites, pyroclastic deposits, called “Tufs anthacifères series”, rhyolitic to dacitic dykes and hypovolcanic microgranites. Geochemistry indicates that crustal melting was triggered by heat input from the mantle. Moreover, a mantle contribution as the source of magma is also likely (Pin and Duthou, 1990). The structural control of dyke intrusion complies with a NW-SE stretching related to the early stage of orogenic collapse. In the northern Cévennes, the Para-autochthonous Unit is underlain by migmatitic ortho- and paragneiss called “the Masméjean Unit” or pre-Velay migmatites (Faure et al., 2001). The anatexis is dated between 333 and 324 Ma by the Chemical U/Th/Pb method on monazite (Be Mezème, 2002). The migmatites and cordierite granites of the



Figure 7 - Structural and geochronologic features related to the Visean-Namurian D3 tectonics and Late Visean magmatism (AMBP: Magnetic Anomaly of Paris Basin).

Montagne Noire Axial Zone (cf D2 field itinerary in section 3) yield similar ages. In the present state of knowledge, this Late Visean event is still poorly studied. It is likely that other pre-Velay migmatites are not yet recognized also within the Velay dome.

2.3.5. The Namurian-Westphalian plutonism corresponds to the main period of magma production in the French Massif Central. It is well acknowledged (Didier and Lameyre, 1971) that this magmatism is represented by two types of granitoids, namely porphyritic monzogranites, such as the Margeride or Pont-de-Montvert-Borne plutons, and biotite-muscovite leucogranites such as the Brame or Millevalche




Figure 8 - Map showing the distribution of the Devonian plutons and volcanic rocks related to the magmatic arc and ophiolites (Brèvenne, Ligne Klippes) interpreted as back arc basins (adapted from Faure et al., 1997).

Figure 9 - Distribution of the main granitic plutons coeval with the stretching lineation and kinematics related to the Namurian-Westphalian extensional tectonics. During this event, the pre-Viséan GuÉret pluton behaves as a rigid body.

massifs (Fig. 9). Although both granite types crop out throughout the Massif Central, the former type is best represented in the central and southern parts of the massif and the later type is more abundant in the north and west parts. The two types were derived from different magmas, but field relationships and geochronology show that these two magmatic types emplaced coevally. Petro-structural and AMS studies of the Namuri-Westphalian plutons show that these bodies are characterized by a conspicuous NW-SE trending mineral, stretching and magnetic lineation. The same trend is also observed in biotite and andalusite contact minerals in the pluton host rocks (Fig. 9). In the north Limousin, the Brame pluton is bounded to the west by the Nantiat ductile normal fault that also exhibits a NW-SE trending hot slickenline. A similar kinematics is also found along the Argentat ductile normal fault. This structural pattern is interpreted as the consequence of the syn-orogenic extensional tectonics of the Massif Central (Faure, 1995).

2.3.6. The Stephanian magmatism is represented by cordierite granite and migmatites of the Velay dome, and also by acidic tuff, ash layers and more rarely alkaline basalts interlayers with terrigenous formations in the coal basins. The Velay dome is bounded to the north by a detachment fault, the Pilat ductile normal fault (Malavieille et al., 1990). Gneiss and micaschists belonging to the LGU that crop out north of the Pilat fault form the substratum of the Late Carboniferous St-Etienne basin.

2.4. A possible geodynamic scenario.

The above-presented data allow us to discuss a geodynamic evolution model. Presently, two scenarios for the evolution of the French Massif Central are proposed. The first one emphasizes a continuous convergence between Gondwana and Laurussia from Silurian to Early Carboniferous (e. g. Matte, 1991, Lardeaux et al., 2001). The second one points out a polycyclic evolution (Pin, 1990; Faure et al., 1997). According to this model, an Early Paleozoic cycle, (Cambrian to Early Devonian), is related to the opening and closure of the Medio-European Ocean and correlatively drifting and rewelding of Armorica



to Gondwana. A second orogenic cycle ranging from Middle Devonian to Carboniferous accounts for which the closure of the Rheic Ocean and the collision of Gondwana and Laurussia. Whatever the preferred model, the following stages are acknowledged.

2.4.1. The breaking of the north Gondwana margin. From Cambrian to Early Silurian, the Massif Central belongs to the northern passive margin of Gondwana which extends from South America to China. From the study of Montagne Noire, the Cambrian-Ordovician corresponds to a terrigenous environment, followed in Devonian by a carbonate platform. The lack of Late Ordovician and Silurian deposits is interpreted as the result of erosion on tilted blocks (Robardet et al., 1994). Evidence of an Ordovician rifting is also inferred from magmatism. In the Para-autochthonous Unit, alkaline mafic volcanics (sometimes with pillow lava), diabase dykes, gabbro intrude the grauwacke-pelite series (Pin and Marini, 1993). In LGU, the alkaline Ordovician granitoids also comply with continental rifting. It is worth noting that "pseudo-calc-alkaline" geochemistry of these granitoids is due to crustal contamination (Duthou et al., 1984; Pin and Duthou, 1990). In the UGU, crustal thinning due to continental rifting is coeval with the emplacement of the leptynite-amphibolite complexes. As a matter of




Figure 10 - Devonian geodynamic reconstruction (map and section) showing the closure of the Rheic Ocean by southward subduction below Gondwana and related microcontinents (from Faure et al., 1997).

fact, the Cambrian-Ordovician period is characterized by the formation of continental stripes, such as the Armorica microcontinent drifted from the north Gondwana margin. The question of the maximum width of the intervening Medio-European Ocean has not been settled yet (see discussion in Robardet, 2003). A rough estimate suggests that this oceanic area was of limited extension (i. e. between 500 and 1000 km).

2.4.2. The closure of the Medio-European Ocean.

On the basis of available dates on the high-pressure metamorphism, the closure of the Medio-European

Ocean started in Silurian. All authors accept a northward subduction of the Gondwana margin, however, structural constraints (i. e. kinematics coeval with the development of high-pressure assemblages) or geodynamic evidence (i. e. relics of a magmatic arc) are lacking. By Middle Devonian time, the Armorica microcontinent is rewelded to Gondwana. In NE Massif Central, (North of Lyon), undeformed and unmetamorphosed Givetian sedimentary rocks unconformably cover the migmatites and high pressure rocks (Delfour, 1989; Godard, 1990). Subduction of oceanic and continental rocks is followed by their exhumation in Early to Middle Devonian, around 390-385 Ma. The lack of large volumes of Devonian clastic rocks suggests that exhumation was tectonically assisted. Exhumation results in the extensive retrogression of the high-pressure rocks of the UGU and migmatisation of the pelitic parts.


2.4.3. Mid-Devonian magmatic arc-back arc system.

Frasnian-Fammenian calc-alkaline volcanism in the NE Massif Central and Vosges argue for subduction. In addition, the 380-370 Ma calc-alkaline diorite, tonalite, granodiorite plutons that crop out in NW Massif Central are interpreted as the deep part of this magmatic arc. However, in their present position, these plutons are rootless and tectonically included into the Hercynian nappes. Southward subduction of the Rheic Ocean is viewed as the cause of the calc-alkaline magmatism. At the same time, distension also occurred in the upper plate, giving rise to limited oceanic zones such as the Brévenne in the Massif Central or other areas in the Massif Armoricain and Vosges. Therefore, an arc-back arc pattern appears as the most likely geodynamic setting for Devonian times (Fig. 10). However a discussion of the Léon and microcontinents is beyond the scope of this presentation.

2.4.4. The closure of the Rheic Ocean and the Tournaisian collision.

Since the Late Fammenian, a complete closure of the Rheic Ocean led to a collision between the North European continent made by the assembly of Laurentia, Baltica and Avalonia during the Caledonian orogeny and Gondwana, including Armorica microcontinent rewelded to it. Intracontinental shortening follows the Lizard ophiolite obduction

Figure 11 - Interpretative lithosphere scale cross section through the French Hercynian Belt in Late Visean. Mantle lithosphere delamination may account for the contrasted tectonic regimes (extensional and compressional), magmatism and heat flow in the central part of the belt (modified from Faure et al., 2002).



(which probably extends along the Magnetic Anomaly of Paris Basin). North-directed thrusts develop from the South of England to the Ardennes. In the northern Massif Central, the closure of the Brévenne oceanic area is characterized by top-to-the-NW shearing under upper greenschist-lower amphibolite facies in the mafic rocks. Top-to-the-NW ductile shearing coeval with middle temperature / middle pressure metamorphism, and dated around 360 Ma is also widespread in western and northern Massif Central. In the southern part of the Massif Central, southward shearing and recumbent folding develops with a progressively younging southward: ca. 345-340 Ma in the Para-autochthonous Unit to 330-325 Ma in the Fold and Thrust Belt.

2.4.5. Late Visean-Namurian syn-convergence extension.

As soon as Late Visean (ca 335 Ma), the northern Massif Central experienced crustal melting responsible for the "tufs anthracifères" acidic volcanism and related plutonism. The structural analysis of the Late Visean plutons and dykes emplacement is controlled by a NW-SE maximum stretching direction and argue for an incipient stage of syn-orogenic collapse in the inner part of the belt (Fig. 7). However, the southern and northern external zones of the Hercynian Belt, such as Montagne Noire and Ardenne respectively, are still under compression as shown by the development of kilometer-scale recumbent folds and thrusts.

In the Central and southern Massif Central, the thermal

overprint is responsible for migmatite and cordierite granite formation. The ca 330 Ma migmatites that crop out in the Montagne Noire Axial Zone and south of the late Carboniferous Velay massif belong to this event. However, in the present state of knowledge, the tectonic setting (namely extensional or compressional tectonic regime) is not settled yet. Decoupling of lithospheric mantle from crust, i. e. lithospheric delamination, is likely to play a significant role to account for the magmatism (Fig. 11).

From Namurian to Westphalian (ca 325-310 Ma), orogen parallel extension is well recorded by emplacement fabrics of leucogranites and granodiorites in the Massif Central (Fig. 9). In the Massif Armoricain, leucogranitic are also widespread. There, they are syn-kinematic plutons coeval with dextral wrenching (e. g. Berthé et al., 1978). However, it is worth noting that both in the Massif Armoricain and Massif Central the wrench or extension controlled synkinematic plutons exhibit the same NW-SE maximum stretching direction. This tectonic stage corresponds also to the main metallogenic epoch for mesothermal gold deposits.

2.4.6. Stephanian post-orogenic NNE-SSW extension

The last stage of the Hercynian orogeny in the French Massif Central corresponds to the collapse of the whole belt. Extensional regime is well recorded by the tectonic setting of intra-mountain Stephanian coal basins. Two structural types of basins are recognized : 1) half-graben bounded by pure normal faults or normal faults with a



Figure 12 - Massif Central map showing the Carboniferous extensional structures: coal basins, Velay granite-migmatite dome.





Figure 13 - Structural map of the Montagne Noire modified from Gèze (1949) and Arthaud (1970).

strike-slip component or 2) pull-apart controlled by wrench faults (Fig. 12). Among these intra-mountain basins, the St-Etienne coal basin is one of the most famous since it corresponds to the para-stratotype of the Stephanian stage. Nevertheless the structural control, either as a left-lateral pull apart or a half-graben is not settled yet (Mattauer and Matte, 1998). In the scale of the Massif Central, the deformation pattern of Stephanian extension is characterized by NE-SW stretching, NW-SE and vertical shortening. The amount of extension increases from west to east. NE-SW extension and correlatively coal basins are widespread in eastern Massif Central but are rare in western Massif Central and almost lacking in the Massif Armoricain. Several N-S to NNE-SSW trending wrench faults such as the Sillon Houiller and Argentat fault are interpreted as transfer faults that accommodate different amounts of extension. Magmatism and mid-crustal deformation associated to Late Carboniferous extension are less developed than during syn-convergence extension. The most spectacular structure is the 100 km diameter migmatitic-granitic Velay dome (Ledru et al., 2002). Heat input from the mantle is responsible for high temperature granulitization of the lower crust (Pin and Vielzeuf, 1983).

Figure 14 - Schematic stratigraphic column of the Paleozoic series found in the Montagne Noire southern side recumbent folds (adapted from Arthaud, 1970).



3. Field itinerary

DAY 1

Recumbent folding in the Montagne

Noire Southern Side

Montpellier--> Béziers --> Cessenon D 136 to St-Nazaire de Ladarez

A. Geologic setting

Following Gèze (1949) and Arthaud (1970), the Montagne Noire area is classically divided from South to North, into a Southern Side, an Axial Zone and a Northern Side (Fig. 13). This last area is less studied than the previous two. The geology of the Axial Zone will be presented during D2. The Southern Side is worldwide famous for Paleozoic stratigraphy (Lower Cambrian,




Figure 15 - Sketch of the main units observed in the eastern part of the Montagne Noire.

Five tectonic units are recognized in the southern side of the Montagne Noire, namely from top to bottom (Fig. 15):

- i) The Pardailhan Nappe (or recumbent fold)
 - ii) The Mont Peyroux Nappe
 - iii) The Monts de Faugères Unit
 - iv) The Cabrières Unit
 - v) The Para-autochthonous domain
- The Pardailhan Nappe consists of

Lower Ordovician, Devonian, Carboniferous) and the development of kilometer-scale recumbent folds (or nappes). The stratigraphic column is schematically summarized in Fig. 14.

folded and overturned Cambrian to Devonian rocks. The Mont Peyroux Nappe includes Ordovician to Viséan rocks. The Monts de Faugères Unit consists of several overturned folds of Devonian to Viséan rocks. The Cabrières Unit is an olistostrome, with

large-scale olistoliths of Carboniferous and Devonian limestones, Silurian volcanites and Ordovician turbidites are resedimented within a wild-flysch matrix corresponding to the foreland basin of the belt (Engel et al., 1980).

The Pardailhan Nappe exhibits a conspicuous axial planar cleavage, whereas in the Mont Peyroux Nappe, the transition between ductile deformation with axial planar cleavage folds and synsedimentary structures can be observed. On the basis




Figure 16 - D1 route map.




Figure 17 - Synthetic cross section of the Montagne Noire with location of the D1 and D2 stops.

of stratigraphy, those nappes were emplaced in Visean-Namurian times (around 330-325Ma).

The aim of this first day is to present the polyphase deformation of the Mont Peyroux recumbent fold through a South to North cross section along the Orb river (Fig. 16). There, the stratigraphic succession ranges from Early Ordovician (Tremadoc-Arenig) to Middle Carboniferous (Visean). Along this route, deformation and metamorphism increase from South to North, however most of the observed structures develop after recumbent folding during an upright deformation linked with the formation of the Axial Zone dome (cf. D2).

B. Stop description

The stops will show most of the lithological and structural aspects of the Mont Peyroux nappe and Para-autochthonous domain underneath (Figs. 16, 17).

Stop D1.1:

Visean flysch with limestone blocks.

The landscape shows Eocene limestone

unconformably overlying at low angle Paleozoic rocks. The S. part of the section exposes Visean flysch with continuous sandstone beds dipping east (20E 60). The northern part that is folded and sheared exposes disrupted beds with sandstone lenses and limestone blocks. Devonian limestone overlying the Visean turbidite is seen on the other side of the valley and nearly 200m to the north of this stop.

Stop D1.2:

Coumiac quarry (protected area)

-Frasnian/Famennian boundary.

This old quarry was mined for red nodular limestone exported all over the world (e. g. the White House in Washington, or the Maison de la France in Rio de Janeiro). The vertical beds (N 30. 90) are Late Devonian (365 Ma) Goniatile limestone called "Griotte marble" (griotte is a type of cherry). This section has been chosen as the Global Stratotype Section for the Frasnian/Famennian boundary. This series corresponds to the "Famenian Biological Crisis" responsible for one of the most severe mass extinction in the Earth history (Kapper et al., 1993).

Stop D1.3:

Early Ordovician turbidite. NE of Ligné.

In the landscape, looking to the SE, the vertical cliffs are Early Devonian limestones continuous with those seen in the previous stop (D1-2) in the Coumiac quarry.

Sandstone-mudstone alternations lie subhorizontally, however graded bedding and load cast show that the sequence is upside down. Folds are apparently overturned to the north but correspond in reality to the inverted limb of the Mont Peyroux recumbent fold. This Early Ordovician turbidite is interpreted as deposited along the northern passive margin of Gondwana.

Stop D1.4:

Early Ordovician turbidite.

A few hundred meters from the previous stop.

The subvertical to north dipping upside down of the beds exhibits numerous load casts, ripple marks and bioturbation evidence (worm burrows). Locally *Lingula* shells can be abundant. Sandstones contain abundant floated muscovite and heavy minerals. In this outcrop, like the previous one, cleavage is lacking.

Stop D1.5:

Panorama on Roquebrun synform and the Orb River. Col de la Vernède.

Below the road, the vineyards and Orb river are located in the Ordovician turbidites which form the core of the Roquebrun synform. Looking northward, above the Roquebrun village, the white cliffs are Devonian limestones. In the background, the hills with bushes are Visean flysch and in the distance, the last hills are made of Devonian limestone belonging to the Monts de Faugères Unit. To the West (left), the highest mountain is the Pic de Naudech made of inverted Cambrian rocks overlying inverted Ordovician turbidites belonging to the Pardailhan Nappe. Lastly, the farthest mountain to the NW is the Mt Caroux composed of orthogneiss belonging to the Axial Zone.

Along the other side of the road, the Ordovician turbidite is complexly folded. Superimposed folds are observed in the next stop.

Stop D1.6:

Superimposed folds in Ordovician turbidite. 200 m down to Roquebrun.

The Ordovician turbidite experienced two folding

phases. Recumbent isoclinal folds (F1) are deformed by upright open folds (F2) with axes plunging 50° NW. F1 are related to the Mont Peyroux recumbent fold and F2 belong to the kilometer-scale upright folding responsible for the Roquebrun synform, Vieussan synform, and Axial Zone antiform (Fig. 17).

Stop D1.7:

Ordovician-Devonian contact. North of Roquebrun.

This stop shows the inverted stratigraphic contact between Ordovician detritals and Devonian carbonates in the northern limb of the Roquebrun synform. From south to north: Ordovician turbidite with top-to-the-S base (with load casts) dipping southward is underlain by Devonian calcareous sandstone, followed by limestone and dolomite. At the northern end of the outcrop, undeformed crinoid stems can be observed in the Devonian carbonates.

Stop D1.8:

Visean flysch. Chapelle St-Poncian, S. of Ceps.

Looking to the NW, the white rocks above the village of Ceps are inverted Devonian limestone, and to the W and SW, the vineyards are located in the Ordovician turbidite. The highest white cliff in the background (La Tour du Pin summit) is the northern extension of Devonian formations. Below the cliff and up to Ceps, the lowest parts of the mountains are made of Visean flysch, belonging to several tectonic units.

The outcrop exposes Visean mudstone-sandstone with limestone intercalations dipping south-westwards (S0-1: 130 SW 50). Contrasting with the Visean rocks observed at the first stop (point D1-1), here the Visean pelites are slightly metamorphosed (sericite) and exhibit a N70E trending crenulation lineation. Chevron folds and south-directed brittle shear zones with quartz veins deform S0-1. Along the road, Devonian rocks are not observed, a late fault separates the Visean and Devonian rocks.

Stop D1.9:

Monts de Faugères Unit. Large curve of Orb River below Chapelle St-Geminian.

Tournaisian (?) - Visean limestone and sericite metapelite present a westward (170W40) dipping foliation and a well marked mineral, stretching and crenulation lineation trending N 70E. Pressure-resolution is the dominant deformation mechanism. S0-1 is also cut at high angle by west dipping tension

gashes filled by fibrous calcite.

Stop D1.10:

Para-autochthonous Domain. Le Pin and Le Lau anticlines.

Turning right to the road of Le Pin, we can observe the underlying Para-autochthonous Domain. North of Le Pin, this outcrop exposes the deepest part of the Orb section. From North to South, the upside down sequence consists of the Upper Devonian red nodular limestone (griotte marble) with goniatites (S0: 60NW60) with an inverted limb subhorizontal cleavage; Tournaisian radiolarian black cherts (lydiennes) and nodular limestones (calcaires de Faugères) and Visean flysch. North of this outcrop, the succession becomes normal from Devonian limestone to Visean flysch from bottom to top, respectively. This S-SE verging fold is called "Le Pin" anticline. Bedding-cleavage relationships with cleavage refraction in sandstone beds comply with the anticline geometry. A N70E composite lineation due to elongated nodules and goniatites, crenulation and intersection develops. Regionally, this para-autochthonous series is folded by two anticlines (Le Pin and le Lau folds) overturned to the South.

Back to the main road, the contact between the Para-autochthonous series and the Mont Peyroux nappe is marked by numerous quartz veins (no stop).

Stop D1.11:

Recumbent fold in Devonian limestone. Moulin de Graïs.

This famous outcrop (Color Plate 1, A) exposes a folded Late Devonian limestone (partly dolomitized). Bedding-cleavage relationships show a southward overturning. The horizontal part of the outcrop is the normal limb. To the south, radiolarian chert (Color Plate 1, B) and limestone are involved in isoclinal folds belonging to the same large-scale structure. It is worth noting that stretching lineation trends close to the fold axis and thus at high angle to the transport direction. In XZ section, pressure shadows indicate top-to-the-NE sense of shear.

Stop D1.11':

Optional. Landscape on the Vieussan antiform.

Turning left on D177, to Berlou, the large curve to the right provides a clear panorama of the northern limb of the Vieussan antiform, well marked in the Devonian limestones.

Stop D1.11»:

Basal thrust contact of the Pardailhan recumbent fold "Queue de cochon (pig's tail)".

Southward, the road goes through the Ordovician turbidite of the Mont Peyroux recumbent fold deformed both by isoclinal and upright folds. The contact between Ordovician turbidite belonging to the Mont Peyroux recumbent fold and the Devonian limestone boudins marking the basal thrust contact of the Pardailhan recumbent fold can be observed in the tight curve north of Escagnès. In spite of intense shearing, the limestone is weakly or undeformed. Back to Vieussan by the same road.

D1.12:

Ordovician/Devonian contact. N. of Vieussan.

Looking West, the hill slope shows several white masses corresponding to Devonian limestone boudins along the basal thrust contact of the Pardailhan recumbent fold (Fig. 18). The outcrop exposes inverted stratigraphic contact between Ordovician turbidite to the left and Early Devonian sandstone to the right. Isoclinal folds with curved hinges can be observed in the Ordovician sandstone. The angular unconformity between Ordovician and Devonian formations, and the lack of Late Ordovician-Silurian rocks in most of the Montagne Noire southern side can be interpreted as a sedimentary consequence of a remote tectonic-metamorphic event that took place more to the north in the internal zone of the Belt. It is worth noting that sedimentology of eo-Devonian rocks indicates a northern source of the terrigenous sediments. Detrital volcanic quartz grains, mica, garnet, zircon, rutile, tourmaline support a pre-Devonian metamorphic event occurring in the hinterland.

Stop D1.13:

Ordovician turbidite in the north part of the Mont Peyroux recumbent fold. N of Vieussan.

Looking to the north, the landscape presents the Axial Zone gneiss and the entrance of Gorges d'Heric visited on D2. The village of Tarassac is built on Devonian marbles (D1-14), the front view is Ordovician turbidite at the western pericline of the Vieussan antiform.

At the outcrop scale, the Ordovician rocks are black pelite and sandstone deformed by upright N80E trending folds (F2) and N50E isoclinal folds (F1). A few biotite grains can be observed in the vertical S2 foliation axial planar to F2.

Stop D1.14:

para-autochthonous Devonian marble. Tarassac, parking of VVP.

Muscovite bearing Devonian marble with pink calcite crystals corresponding to deformed crinoid stems exhibit a southward dip (70S60) and well marked subhorizontal mineral and stretching lineation. This marble is separated from the overlying Ordovician turbidite by a major thrust contact corresponding to the basal thrust surface of the Mont Peyroux recumbent fold. The Devonian marble and the underlying metapelites attributed to Ordovician (not seen here) are a normal sequence belonging to the Para-autochthonous Unit. ⁴⁰Ar/³⁹Ar date on muscovite gives 297 ± 3 Ma which is interpreted as the age of a Late Carboniferous gravity sliding event related to the formation of the Axial Zone (Maluski et al., 1991).

End of the 1st day. Overnight stay in Olargues

17). The foliation of micaschists and gneiss defines a NE-SW long axis elliptical dome whose western part is disturbed by the Eocene Mazamet thrust. Some authors argued that the Axial Zone metamorphic rocks correspond to the Precambrian basement of the Paleozoic series observed in the recumbent folds. In the present state of knowledge, there is no argument to support the existence of a Neo-Proterozoic (i. e. Cadomian) orogen in the Massif Central. Therefore, the reality of a Precambrian basement in the Montagne Noire Axial Zone is not supported by the data. The augen orthogneiss seen in the gorges d’Heric, are porphyritic granites intruding a Neo-Proterozoic to Paleozoic metasedimentary series of micaschists and gneiss and transformed into augen gneiss during Hercynian tectonics. Recent U/Pb dating supports an Early Paleozoic age for the magmatism. The presence of penninic style recumbent folds overturned to the north has also been assumed (Demange, 1975). Although possible, this interpretation cannot be demonstrated, mainly due to poor outcrop conditions.




Figure 18 - Panoramic view of the contact between the Pardailhan (top) and Mont-Peyroux nappe (bottom) marked by Devonian limestone boudins called “ pig’s tail ”.

DAY 2

Migmatite dome of the Montagne Noire Axial Zone

A. Geological setting

The Montagne Noire Axial Zone remains one of the most controversial area in the geology of Massif Central (cf extensive references in Soula et al., 2001). The Late Visean-Early Namurian recumbent folds examined during D1 are overprinted by metamorphic and structural features related to a granite-migmatite gneiss dome developed in the Axial Zone (Figs. 13,

The Axial Zone gneiss experienced a HT/LP type metamorphism up to partial melting giving rise to migmatites and anatectic cordierite granites (e. g. Laouzas granite). U/Pb dating on single grain zircon and monazite give a ca 330 age. Isograds of this HT/LP metamorphism define the same domal geometry as the foliation. Within the micaschist envelope, kyanite relics are locally found (K in Fig. 13). P-T paths for the gneiss core and metamorphic envelope have been proposed (e. g. Soula et al., 2001; Fig. 19). It is worth noting that some amphibolites included in the gneiss are retrogressed eclogites (E in Fig. 13) with estimated pressure and temperature around 9 ± 2




Figure 21 - D2 route map

the gneissic linear fabric. In the paragneiss, meter scale isoclinal folds are refolded by NE-SW trending upright folds.

Pegmatite dykes are also deformed by upright ptygmatic folds and veins parallel to the fold axes are boudinated. Along the stream, amphibolite boulders (containing garnet locally) including retrogressed eclogites, boulders of migmatites and migmatized augen gneiss, sometimes containing sillimanite nodules are widespread. Outcrop of migmatite will not be visited along this route.

In spite of a clear stretching and mineral lineation, the sigma-type porphyroclast systems exhibit both dextral and sinistral asymmetry at the outcrop scale. The ambiguous sense of shear might be due to superimposed deformation (namely doming overprinted upon low angle shearing) or to strain partitioning with a significant component of pure shear (Color Plate 1, C) during doming.

The bulk structure of the Héric section is a gneiss antiform overturned to the north (Figs. 13, 17). The antiform hinge zone is parallel to upright

folds in augen gneiss and meter to millimeter-scale crenulation.

Stop D2.2:

Augen gneiss and sheared pegmatite veins at the eastern part of the Caroux dome. Le Vernet .

From gorges d'Héric → east to Le Poujol → north (left) to Combes.

Fine grain augen gneiss (Sx 40SE 30) with a N70E trending stretching lineation at the eastern termination of the Caroux antiform. Sigmoidal K-feldspath indicates a down-dip, top-to-the East sense of shear. Pegmatite dykes cross cutting the foliation are also sheared to the East. Along the road next curves, many asymmetric pegmatite boudins can be observed within weathered gneiss (no stop).

Stop D2.3:

Biotite-Garnet-staurolite micaschists. Crossing of the road to Forêt des Ecrivains Combattants.

The gneiss-micaschist series experienced a high-

temperature and low-pressure metamorphism characterized by biotite, garnet, andalusite, staurolite tight isograds. The eastward dipping foliation (160E15) bears a composite crenulation, mineral and stretching lineation trending N70E along which top-to-the-E shear criteria develops. In thin section, quartz pressure shadows, helicitic garnet and staurolite, and shear bands indicate a top-to-the East shearing (Color Plate 1, D).

After the Col de Madale, the road runs within the Rosis synform which consists of crenulated and folded HT/LP micaschists between the Caroux antiform and the Espinouse dome.

Stop D2.4:

Migmatitic orthogneiss. Col de l'Ourtigas.

Partial melting develops at the expense of the Heric orthogneiss but MFK and gneissic fabric are still preserved. Migmatization is dated at 330 Ma. After the pass, the road crosses the northern border of the Vialais granite, however due to poor exposure quality and parking difficulties, the excursion will not stop there.

Optional

Stop D2.4':

Sheared augen gneiss and migmatite.

S. of Ginestet.

The migmatitic orthogneiss experiences a ductile shearing, the NW-SE trending foliation dips 50NE and bears a N60E trending slickenline corresponding to the Espinouse normal fault. $^{40}\text{Ar}/^{39}\text{Ar}$ dates on muscovite and biotite indicate 297 ± 3 Ma (Maluski et al., 1991).

North of Ginestet, begins the Montagne Noire Northern Side. In spite of globally poor exposure quality, Early Cambrian carbonates can be observed on top of the mountains.

Stop D2.5:

Stephanian conglomerate. Falaise d'Orques.

At the pass of Croix de Mounis, the road enters into the Stephanian Graissessac coal basin. In the northern landscape, the Early Cambrian limestone cliffs are exposed.

The Late Carboniferous conglomerate contains cm to m size blocks of Early Cambrian sandstone and limestone. Along the road, a decameter-scale Cambrian limestone block that is probably an olistolith can be observed too. Stephanian beds dipping 110S10 unconformably cover the Paleozoic (Cambrian-

Ordovician) sedimentary rocks of the Northern Side. Thus strictly speaking, the southern border fault of the Graissessac basin (Castanet fault) is reactivated after the deposition of the Stephanian rocks.

At the crossing with D22E road to Castanet-le-Haut, the landscape to the east shows the foliation of the Espinouse gneiss dome dipping north-eastward. The ductile dextral-normal Espinouse fault separates the Espinouse dome from the Stephanian Graissessac basin.

Stop D2.6:

Sheared rocks along the Espinouse fault.

N. of Nougayrols.

The Graissessac basin substratum (i. e. early Paleozoic rocks of the Northern Side) is sheared by the Espinouse fault. Quartz veins are well developed.

Stop D2.7:

Weakly metamorphosed schists. West of Castanet-le-Bas.

Weakly deformed and weakly metamorphosed greenish sandstone-pelite series correspond to the outermost part of the metamorphic rocks surrounding the Axial Zone. Centimeter- to meter-scale folds overturned to the SE can be observed at this outcrop. East of Verenoux, the road goes through unmetamorphosed and undeformed greenish sandstone and pelite corresponding to the Early Cambrian (grès de Marcory). This Unit (called Monts de St-Gervais Unit) belongs to the Northern Side, and forms an extensional allochthon emplaced upon the Axial Zone micaschists. No stop.

Stop D2.8:

Stephanian massive sandstone and coal mesures. East of La Mouline.

The Late Carboniferous rocks belong to the Graissessac coal basin. They are fluvial deposits dipping 120NE30 with coal intercalations. Decollement surfaces may develop along coal mesures. In the ancient open pit of the Graissessac mine (not visited) synsedimentary folds overturned to the South can be observed. These Late Carboniferous rocks are unconformably covered by Early Permian conglomerates, sandstones and pelites dipping 0E20, but unfortunately the contact cannot be observed there.

Stop D2.9:

Permian (Autunian) conglomerate.

West of La Tour-sur-Orb.

Continental conglomerates with quartz, sandstone pebbles.

Stop D2.10:

Permian extensional tectonics.

Mas d'Alary quarry.

In the ancient open pit formerly mined for uranium, continental red beds are cut by north dipping normal faults. Alike other early Permian basins in S. France (e. g. St-Affrique, Rodez), the Lodève basin is a half-graben bounded to its southern margin by normal faults.

End of the 2nd day. Drive to Millau (ca 60 km), overnight.

DAY 3

The stacking of Upper and Lower Gneiss Units and post-nappe crustal melting


Millau → Highway to La Canourgue Exit n° 39

Leaving Millau to the North, the road crosses the Jurassic limestone plateau called Causse de Sauveterre. Near Séverac-le-Château, Toarcian black shales were mined for oil. The Jurassic sedimentary series observed along the road, is deformed by decameter to kilometer-scale wave-length upright folds and reverse faults. These structures are related to the Eocene compression due to the Pyrenean orogeny.

A. Geological setting

The Marvejols area (Fig. 22) is a famous place in the geology of the French Massif Central since it is one of the first places where Variscan syn-metamorphic nappe thrusting has been documented on the basis of geochronology, metamorphism and tectonics (Pin, 1979). The metamorphic inversion with HP rocks of the Upper Gneiss Unit upon the Lower Gneiss Unit was used as an argument for tectonic superposition (Fig. 23). The contact between the two units is a high temperature mylonitic zone (Faure et al., 1979). Moreover, radiometric dates document the tectonic

Figure 22 - D3 route map




evolution of the area. The Lower Gneiss Unit, called locally the Lot Series, consists of a monotonous succession of metapelite and metagrauwacke intruded by various magmatic bodies:

1. Quartz-diorite orthogneiss is dated at 540 ± 12 Ma by U/Pb isotopic dilution method on zircon populations (Pin and Lancelot, 1978). This rock exhibits a well marked post-solidus planar and linear structure. $40\text{Ar}/39\text{Ar}$ dating on biotite gives 352 ± 1 Ma (Costa, 1989).
2. Pink K-feldspar orthogneiss of alkaline composition is not well dated due to inherited zircons. By comparison with the Mendic orthogneiss in the Montagne Noire northern side, an Early Cambrian age is generally accepted.
3. An acidic augen orthogneiss with mylonitic zones crops out immediately below the Upper Gneiss-Lower Gneiss Unit contact. An U/Pb age on zircon populations gives a lower intercept age of 346 ± 8 Ma (Pin, 1981). Due to its tectonic setting, and radiometric age this orthogneiss is interpreted as a synkinematic pluton coeval with the thrusting of the Upper Gneiss Unit. However, this conclusion does not comply with microstructural data and particularly with the NW-SE stretching lineation developed in the Lower Gneiss Unit (cf below). A preliminary chemical U/Th/Pb age of 370 ± 12 Ma is obtained on monazite (A. Joly unpublished data).
4. Other small gneiss masses are recognized in the Lower Gneiss Unit, some of them are considered as hypovolcanic granites or volcaniclastic metasediments (Pin, 1981).

As seen in the first steps of D3 day, the Lot Series of the Lower Gneiss Unit is characterized by a subhorizontal foliation and a conspicuous NW-SE trending mineral and stretching lineation. This ductile deformation is coeval with an intermediate temperature-intermediate pressure metamorphism. Biotite, garnet, staurolite, andalusite and muscovite are widespread in the metapelites. The Marvejols area is often taken as an example of the inverted metamorphism developed in the footwall of the Upper Gneiss Unit overthrust (Fig. 24). $^{40}\text{Ar}/^{39}\text{Ar}$ dates on biotite and muscovite from the Lot Series micaschist yields 351 ± 4 Ma and 342 ± 4 Ma respectively (Costa, 1989). The tectonic significance of the NW-SE trending stretching lineation which is widespread throughout the Massif

Figure 23 - General cross section from Marvejols to the Truyère area showing the tectonic superposition of the Upper Gneiss Unit upon the Lower Gneiss Unit and the shape of the Margeride pluton






Figure 24 - Metamorphic map of the Lower Gneiss Unit around the west part of the Margeride pluton (Marvejols, Truyère, Châtaigneraie) showing the inverted metamorphism (adapted from Burg et al., 1984).

Central is not clearly settled yet.

The mylonitic zone at the base of the Upper Gneiss Unit is characterized by a N-S trending mineral and stretching lineation. In spite of intense recrystallization and post-kinematic annealing, top-to-the-south shearing can be observed. $^{40}\text{Ar}/^{39}\text{Ar}$ dating of synfolial biotite gives 358 ± 4 Ma interpreted as the age of thrusting. Moreover, a late muscovite developed upon the early foliation is dated at 340 ± 4 Ma (Costa, 1989).

In the Marvejols area, the Upper Gneiss Unit consists of a lower part called "leptynite-amphibolite" sequence and an upper part with migmatitic gneiss and micaschist. The leptynite-amphibolite sequence contains metamorphosed mafic rocks of magmatic or sedimentary origin such as metagabbro, metabasalt, amphibolite rare serpentinites and acidic rocks (i. e. leptynites) Meter to centimeter scale acidic-mafic alternations are probably of volcani-clastic origin. Several U/Pb ages on zircon populations are available (Pin and Lancelot, 1982). Namely, an amphibolite boudin is dated at 487 ± 6 Ma, and a coronitic

metagabbro at 484 ± 7 Ma. An orthogneiss intrusive in the paragneiss gives 478 ± 6 Ma. These dates are interpreted as the evidence for an Ordovician magmatism related to the rifting of Armorica from Gondwana. Moreover, the 415 ± 6 Ma age of zircon populations from a high-pressure trondhjemite is considered as the age of melting coeval with eclogite facies metamorphism. Pressure and temperature constraints on this rock are 16 ± 4 kb and $800 \pm 50^\circ\text{C}$ respectively.

Upper and Lower Gneiss Units stacking was followed by huge crustal melting produced under distinct P-T conditions (Fig. 25). The second part of the D3 and the whole D4 days are devoted to the observation of some manifestations of the Middle to Late Carboniferous crustal melting. From older to younger, three stages are distinguished.

1. Pre-Velay Cévennes migmatites, dated between 333 to 324 Ma by chemical U/Th/Pb method on monazite (Be Mezème, 2002).
2. Namurian-Westphalian plutonism dated around

320-315 Ma. This magmatism is characterized by porphyritic monzogranite (Margeride or Pont-de-Montvert-Borne plutons) and also by leucogranite (Signaux and Rocles plutons). Although distinct massifs derived from different magmas, field relationships show that these two magmatic types are coeval.

3. Velay migmatites and cordierite granite dated around 300 Ma. This large massif will be examined during D4.

B. Stop description

Stop D3.1:

Cambrian quartz-diorite. N88, East of Pont des Ajustons, S. of Marvejols.

A fine-grained biotite-hornblende quartz-diorite originally intrudes micaschists (stop D3-2) belonging to the Lower Gneiss Unit. U/Pb dating on zircon populations gives a 540 ± 12 Ma age (Pin and Lancelot, 1978) for the emplacement of this pluton. $^{40}\text{Ar}/^{39}\text{Ar}$ dates on biotite provide a 352 ± 1 Ma age corresponding to the tectonic event (Costa, 1989). This well foliated and lineated rock experienced a NW-SE deformation.

Stop D3.2:

Lower Gneiss Unit of the Lot series.

The Lot series are composed of biotite-garnet \pm staurolite micaschists originally intruded by porphyric granite and metadiorite. The flat lying foliation bears a conspicuous NW-SE mineral and stretching lineation, indicating a top to NW sense of shear (Color Plate 1, E). Biotite and muscovite $^{40}\text{Ar}/^{39}\text{Ar}$ ages are respectively 351 and 342 ± 4 Ma (Costa, 1989). In the background, Early Jurassic limestones unconformably overly the metamorphic rocks.

Stop D3.3:

Augen orthogneiss within the Lot series. Pont de Pessil.

The Lower Gneiss Unit includes augen orthogneiss with mylonitic fabric. In sections perpendicular to the foliation and parallel to the NW-SE lineation, asymmetric K-feldspar augen indicate a top-to-the-SE sense of shear. U/Pb date on zircon populations gives a lower intercept age of 346 ± 8 Ma (Pin, 1981). An U/Th/Pb chemical age on monazite gives 370 ± 12 Ma (Joly, unpublished data).

Stop D3.4:

Coronitic metagabbro belonging to the Upper Gneiss Unit. Le Croisier.

The Upper Gneiss Unit outcrops North of




Figure 25 - P-T paths of pre-Velay and Velay events constructed from metamorphic rocks sampled in the south part of the Velay area (simplified from Montel et al., 1992).

Marvejols city. Metabasites, locally with mylonitic fabric, underwent the Eovariscan high-pressure metamorphism (Color Plate 1, F). However, due to the heterogeneity of deformation, magmatic textures are still preserved. This rock is dated by U/Pb method on zircon populations (upper intercept) at 484 ± 7 Ma (Pin and Lancelot, 1982).

Stop D3.5:

Leptynite-amphibolite complex

(Upper Gneiss Unit). Along the main road (N9).

This outcrop exposes a typical section of the "leptynite-amphibolite complex" made of alternations of mafic and acidic rocks considered as a volcaniclastic formation. The foliation exhibits meter to

amphibolite complex which becomes migmatized northward. The migmatite is not dated here. By comparison with other places in the Massif Central, a ca 380 Ma age can be inferred.

Stop D3.7:

Middle Carboniferous Margeride pluton.

Truc de Fortunio.

Drive to St-Amans → Estables

The Margeride massif is one of the largest granitic plutons in the French Massif Central (3200 km²). It consists mainly of a porphyritic monzogranite with large (up to 10cm) K-feldspar megacrysts. On the basis of biotite content, three facies, namely dark, intermediate and light facies, are distinguished




Figure 26 - Example of punctual dating on monazite single grain by U/Th/Pb chemical method (Be Mezème et al., 2003).

decameter-size amphibolite boudins (Color Plate 2, A). Zircon populations from an amphibolite give a U/Pb age of 487 ± 6 Ma age (upper intercept) interpreted as the rock formation age (Pin and Lancelot, 1982). The lower intercept at 340 ± 4 Ma is close to the thermal event (ca. 345-330 Ma) that overprints the tangential tectonics.


Stop D3.6:

Early Variscan migmatisation.

Gorges du Val d'Enfer.

The road runs from South to North in the leptynite-

(Couturier, 1977). Nevertheless, more than 80% of the massif is made of the intermediate facies. Moreover, muscovite-K-feldspar leucogranite intrude the monzogranite. Most of the leucogranites are meter-scale dykes, but east of the massif the Grandrieu leucogranite represents a kilometer-sized pluton. The monzogranite yields a Rb-Sr whole rock age of 323 ± 12 Ma (Couturier, 1977) and an isotopic



dilution U/Pb monazite age of 314 ± 3 Ma (Pin, 1979); the Grandrieu leucrogranite is dated at 305 ± 4 Ma (U/Pb method, Lafont and Respaut, 1988). The Margeride pluton is a subhorizontal of 4 to 8 km thick (Fig. 23). Petrostructural and AMS studies (J-Y. Talbot, work in progress, Fig. 27) show a complex pattern of the foliation trajectories. Although the foliation trajectories do not show a consistent shape, at the scale of the whole pluton, the foliation presents a flat lying attitude. The lincation pattern exhibits a well-defined NW-SE trend. It is worth noting that such a trend is widespread throughout the whole Massif Central and corresponds to the maximum stretching direction of the Namurian-Westphalian synorogenic extension (Faure, 1985).

Stop D3.8:

Augen orthogneiss below the Para-autochthonous Unit.

*Drive to Châteauneuf-de-Randon
→ Montbel → NE of Belvezet*

SE of the Margeride pluton, the Cévennes area exposes the para-autochthonous Unit of the Massif Central. Below the unconformity of the Mesozoic sedimentary rocks, an augen orthogneiss and its host rocks crop out in a tectonic window below the Para-autochthonous Unit (Cévennes micaschists). The augen orthogneiss exhibits a subhorizontal foliation and NE-SW trending stretching lincation. The age of the granitic magmatism is not determined here but assumed to be Early Cambrian (ca 550 Ma) by comparison with other orthogneiss in the Massif Central. The chemical U/Th/Pb age of 560 ± 18 Ma in the core of monazites grains from migmatitic orthogneiss supports this interpretation (cf. Stop D3-9)

Stop D3.9:

A few kilometers eastward, biotite, garnet (\pm cordierite) paragneiss and quartz-micaschist correspond to the orthogneiss host rock.

Stop D3.10:**Migmatitic orthogneiss. Barrage de Puylaurent.**

To the east, the orthogneiss experiences a partial melting giving rise to diatexites. Locally the orthogneiss fabric remains well preserved. A chemical U/Th/Pb dates on monazite give 560 ± 18 Ma and 324 ± 6 Ma for the grains core and rim respectively (Be Mezème, 2002, Cocherie et al., in press). The latter is interpreted as the age of migmatization.

Boudinaged dykes of leucogranites.

Pink granite dykes and leucogranite dykes intrude the metamorphic rocks. Muscovite dyke is dated by chemical U/Th/Pb method on monazite at 316 ± 5 Ma (Be Mezème et al., 2003; fig. 26).

In Prévencières, turn to the north (left) to Langogne. The road follows the brittle left-lateral Villefort fault of Permian age. South of La Bastide-Puylaurent, the road is located in the Para-autochthonous Unit (Schistes des Cévennes). From La Bastide Puylaurent, to Langogne, the migmatitic orthogneiss crops out again along the Allier river.

End of the 3rd day. Overnight in Langogne

DAY 4**The Velay dome (French Massif Central): melt generation and granite emplacement during orogenic evolution**

The generation of large granite-migmatite complexes by crustal melting during orogeny is a process still discussed in particular because of the deep, inaccessible location of their production sites (Clemens, 1990; Brown, 1994). Moreover, the development of a partially molten middle crust during collision tectonics implies a major change in the rheology of the thickened crust and largely control its behaviour during orogenic collapse (Vanderhaeghe and Teyssier, 2001). Thus, the Variscan belt which exposes numerous granitic intrusions and large migmatitic complexes is of great interest to study the role of partial melting during orogenic evolution (Brown and Dallmeyer, 1996; Gardien et al., 1997; Vanderhaeghe et al., 1999). The Velay migmatite-

granite dome located in the SE Massif Central (Fig. 28, 29, 30) offers a unique opportunity to examine the thermal conditions required for widespread crustal anatexis and the consequences of the presence of the generation of a large volume of partially molten rocks on the evolution of the Variscan orogenic crust.

The aim of the fourth day is to illustrate the melt generation and granite emplacement of the Velay dome in connection with the tectonic evolution of the Variscan belt, by showing:

- incipient stages of melting in the Late Neoproterozoic pre-tectonic granite and metasediments on the southern margin;
- the cordierite-bearing granites and migmatites;
- the relation between granite emplacement, extensional tectonics and formation of the St Etienne Stephanian basin on the northern flank of the Velay dome.

A. Geological setting

The Velay dome (Fig. 28,29), about 100 km in diameter, is composed of peraluminous granites (about 70%) characterized by abundance of nodular and prismatic cordierite and by enclaves of gneisses (25%) and granites (5%) of various nature and size (Didier, 1973; Dupraz and Didier, 1988). Previous work in this area provided the following results and models

- Montel et al. (1992) describe two successive stages of anatexis, first under water-saturated conditions with biotite stable followed by melting under biotite dehydration conditions.
- Burg and Vanderhaeghe (1993) proposed that the amplification of the Velay dome cored by migmatites and granites reflects gravitational instabilities within a partially molten middle crust during late-orogenic extension.
- Lagarde et al. (1994) suggested that the deformation pattern of the Velay dome records southward lateral expansion of the granites below the detachment zone of the Pilat, one of the major normal faults developed during the collapse of the Variscan belt (Malavielle et al., 1990)
- Geochemical and petrological data published by Williamson et al. (1992), Montel et al. (1992) and Barbey et al. (1999), indicate that the Velay dome has followed a clockwise P-T-time evolution overprinted by a thermal peak due to the underplating of mafic magmas (Fig. 31).



Figure 28 - Simplified geologic map of the eastern margin of the Massif Central, Day 4 and Day 5 localities




Figure 29 - Simplified cross section through the eastern margin of the Massif Central showing the main tectonic units and their structural relationships, Day 4 and Day 5 localities. Ages of some granites are indicated as specific geochronological markers.

According to Ledru et al., 2001, structural, petrologic and geochronological data indicate that the formation of the Velay migmatite-granite dome results from the conjunction of several phenomena.

- Partial melting of the thickened crust started at about 340 Ma, while thrusting in the hinterland of the Variscan belt was still active, and ended during collapse of the orogenic crust at ~300 Ma. Crustal anatexis responsible for the generation of the rocks forming the Velay dome hence lasted about 40 My.
 - Partial melting took place within a dominantly metasedimentary crustal layer dominated by fertile pelitic compositions. Melting reactions evolved from the water-saturated granitic solidus to destabilization of hydrous minerals and indicate that melting started at the end of the prograde metamorphic path and ended during decompression associated with exhumation of the migmatite-granite dome.
 - Thermal relaxation and increased radioactive heat production following crustal thickening likely caused a rise in temperature during the evolution of the Variscan orogenic crust. However, it is proposed that heat advection from mantle-derived magmas and also asthenospheric upwelling coeval with orogenic collapse have provided the extra heat source required to melt a large volume of the thickened crust and generate the migmatites and granites of the Velay dome.
- The formation of the Velay dome, coeval with the activation of crustal-scale detachments, potentially corresponds to flow of a partially molten crustal layer in response to gravitational collapse.

Four main structural zones, that will be partially illustrated during this fourth day, are defined (Fig. 28, 30):

1. The host rocks. The Velay granite-migmatite dome is hosted by gneissic units stacked during the collision history of the Variscan belt (Ledru et al., 1994a, b):
 - the Upper Gneiss Unit, that contains remnants of Early Paleozoic oceanic or marginal basins is presently in an upper geometric position, this unit



Figure 30 - Extension of Meso and Neovariscan metamorphism and foliation trends within the host rocks of the Velay migmatite-granite dome

contains dismembered basic-ultrabasic complexes at its base overlain by gneisses derived from granites, microgranites, acid and basic volcanics, tuffs and grauwackes. Numerous eclogitic relics are preserved within basic layers marking an Eovariscan stage of lithospheric subduction (450-400 Ma). Structural and radiometric data show these rocks were exhumed from 90 km at 420-400 Ma to less than 30 km at 360-380 Ma while subduction was still active (Lardeaux et al., 2001).

- The north Gondwana continental margin is represented by (a) a Lower Gneiss Unit composed of metasediments derived from pelites and argillites, and augen orthogneiss (the "Arc de Fix") originating from peraluminous porphyric granite dated at 528 ± 9 Ma (Rb-Sr whole rock, R'Kha Chaham et al., 1990), and (b) a mainly sedimentary parautochthonous sequence. This margin underwent a general medium-pressure metamorphism attributed to the thrusting of the Upper Gneiss Unit which occurred during Devonian, prior to 350 Ma in the internal zone

(Mesovariscan period). The Lower Gneiss Unit is intruded by syn tectonic granites precursor of the Velay dome emplaced between 335 - 315 Ma, including magnesio-potassic plutons, the so-called "vaugnerite". In the south (Fig. 2), the Cévennes micaschists are interpreted as the parautochthonous domain. Maximum P-T conditions during the metamorphic evolution are there estimated at 500 °C, 5 kbar, with the muscovite-chlorite-garnet parageneses being synchronous with southward thrusting and a thickening estimated at about 15 km (Arnaud and Burg, 1993; Arnaud, 1997). The closure of micas to Ar diffusion has been dated at 335-340 Ma ($^{40}\text{Ar}/^{39}\text{Ar}$, Caron et al., 1991).

2. The gneiss-migmatite zone, at the periphery and at the roof of the Velay dome. In the migmatites and in the gneissic hosts, the following melting reactions are identified (Fig. 31):

- The first melting stage developed under P-T conditions exceeding those for water-saturated quartz-feldspathic rocks, with biotite remaining stable: around

700 °C, 4 kbar within the metamorphic envelope, 5 kbar in the granitic core (M3 stage of Montel et al., 1992). The presence of corundum paragneiss enclaves confirms the initial presence of muscovite and the prograde character of this melting event (Aït Malek et al., 1995). A U-Pb monazite date indicates a minimum age of 314 ± 5 Ma (Mougeot et al., 1997). High-K magnesian monzodiorite, with mantellic affinity are also dated at 313 ± 3 and 314 ± 3 Ma (207Pb-206Pb and U-Pb respectively on zircon, Aït Malek, 1997). They contain peraluminous




Figure 31 - Pressure-temperature evolution from the gneisses to the Velay granite
 Mineral abbreviations: Ky = kyanite; Sil = sillimanite; And = andalusite; Cd = cordierite; Grt = garnet; Ms = muscovite; Qtz = quartz; Kfs = potassium feldspar; Pl = plagioclase; Bt = biotite; As = aluminium silicate; L = liquid. The transition from the M3 to the M4 metamorphic stage is indicated by arrows (1) in the Pilat micaschist (after Gardien et al., 1990), (2) in the migmatitic orthogneiss and granite in the southern part of the dome (oblique- and horizontal-ruled boxes, after Gardien et al., 1997, and Montel et al., 1992), (3) in the migmatitic part of the southern host rocks of the Velay granite (oblique-ruled boxes, after Montel et al., 1992)

xenoliths that record a first stage of isothermal decompression at 700-800 °C, 8-10 kb, consistent with a source located more than 30 km deep, followed by a stage at 5-6 kb (Montel, 1985). In view of the water-saturated conditions, it is unlikely that large quantities of granite (i.e. < 10-20%) were produced and extracted at this stage (Patiño Douce and Johnston, 1990).

- The second stage of melting is characterized by high-temperature metamorphism in the cordierite stability field, with biotite destabilized: 760-850 °C, 4.4-6.0 kbar (stage M4 of Montel et al., 1992). Leucosomes were dated at 298 ± 8 Ma based on Rb-Sr whole rock isochron (Caen-Vachette et al., 1982), and Rb-Sr whole rock-biotite isochrons yield ages between 305 and 276 Ma (Williamson et al., 1992). An age of 301 ± 5 Ma was obtained for the homogeneous parts of the granite using the U-Pb monazite method (Mougeot et al., 1997). Therefore, this second melting stage is considered to be generally synchronous with emplacement of the main cordierite-bearing granites. The volume of cordierite-bearing granites generated makes a case for massive partial melting at this stage, associated to destabilization of hydrous minerals.

3. The migmatite-granite domain. The various granites that appear in the Velay dome define a suite, with 3 main granite types distinguished according to age, structure, homogeneity, mineralogy and geochemistry:

- A heterogeneous banded biotite granite, found mainly on the western margin of the dome and locally on the southern and eastern margin. It corresponds to the first generated granite of the Velay suite. Foliation trajectories are in continuity with porphyric granites in the external rim of the dome suggesting continuity between these precursor granites and the development of the heterogeneous banded biotite granite.
- A main biotite-cordierite granite, in which several sub-types may be distinguished, in particular according to the cordierite habitus (Barbey et al., 1999).
 - a heterogeneous banded granite with abundant enclaves. Most of these enclaves represent incorporated and partly assimilated pieces of the Lower Gneiss unit and precursor plutons originating from the host rocks, although some enclaves with refractory composition or granulite facies metamorphism have a lower crustal origin

(Vitel, 1985). Cordierite may be prismatic, cockade-type or mimetic overprinting previous biotite - sillimanite assemblages. Most of the heterogeneous granites indicate mixing between melts of lower-crustal origin and melts from the para- and ortho-derived host rock (Williamson et al., 1992).

- a homogeneous leucocratic biotite-cordierite granite with mainly cockade-type cordierite. Its emplacement has been dated at 301 ± 5 Ma using the U-Pb method on monazite (Mougeot et al., 1997).
- a homogeneous granite with biotite and prismatic cordierite as a primary ferromagnesian phase, with few enclaves. The heterogeneous and homogeneous granites with prismatic cordierite, with a high Sr content, have a mixed isotopic signature between the host rocks and a lower-crustal origin. The deep source is considered to be the melting of the lower mafic/felsic plutonic crust (Williamson et al., 1992).
- a leucocratic granite with cockade-type cordierite, without enclaves. The cordierite-quartz aggregates postdate primary biotite bearing assemblages and probably prismatic cordierite.
- The late magmatic activity that includes:
 - homogeneous granite with K-feldspar porphyrocrysts and common prismatic cordierite, basic and micaceous inclusions (the Quatre Vios massif) (Fig. 10d). These granites are defined as late-migmatitic and are considered to originate from the melting of aluminous sediments at 4.5-5.5 kbar and 750-850 °C, under water-undersaturated conditions and have a significant mafic component (Montel et al., 1986; Montel and Abdelghaffar, 1993). Ages at 274 ± 7 Ma (Rb/Sr whole rock, Caen Vachette et al., 1984) are considered to be partially reset during Permian or Mesozoic hydrothermal event.
 - Stephanian leucogranites, microgranite and aplite-pegmatite dykes, Permian rhyolites. Microgranite dykes have been dated at 306 ± 12 and 291 ± 9 Ma and a Permian hydrothermal event at 252 ± 11 and 257 ± 8 Ma (microprobe dating of monazite, Montel et al., 2002).

4. The Stephanian intracontinental basin of St-Etienne. This basin is formed along the hanging wall of the Mont Pilat extensional shear zone (Malavielle et al., 1990). The Mont Pilat unit, attributed to the lower gneissic unit at the scale of the French Massif Central

consists of aluminous micaschists, metapelites, orthogneisses and amphibolites. This unit has a gently north-dipping foliation plane bearing a north-south stretching lineation. Numerous leucogranitic pods outcrop more or less parallel to this main foliation plane and have been dated at 322 ± 9 Ma (Rb/Sr whole rock, Caen-Vachette et al., 1984). Shear criteria, observed at different scales, are compatible with a top-north extension dated between 322 and 290 Ma ($^{39}\text{Ar}/^{40}\text{Ar}$, Malavieille et al., 1990). This event was coeval with the progressive development of low pressure - high temperature metamorphic conditions (3- 5 Kbar and 700 - 780 °C, Gardien et al., 1997).

B. Stop description (Fig. 28, 29)

Stop D4.1:

Meyras, Road from Le Puy-en-Velay to Aubenas, N102, Road cut.

Stop D4.1 shows the incipient stage of melting within the orthogneiss of the Lower Gneissic Unit. An augen orthogneiss (the "Arc de Fix"), originating from peraluminous porphyric granite dated at 528 ± 9 Ma (Rb-Sr whole rock, R'Kha Chaham et al., 1990), constitutes an almost continuous rim around the Velay granite-migmatite dome. The melting is marked by the segregation of cordierite-free melts along the main inherited foliation and locally discordant cordierite-bearing granitic patches (Color Plate 2, B). Large phenocrysts of K-feldspar attest of the porphyric type of the granite protolith while the foliation is marked by the elongation and crystallization of the quartzofeldspathic aggregates and biotite-rich melanosome. Magnesian-potassic dykes (the so-called "vaugnerite") are intrusive and boudinated within the orthogneiss.

The progressive development of the anatexis and textural evolution in the transition from subsolidus annealing to melting process has been studied in detail in this zone by Dallain et al. (1999). Anatexis first develops with the resorption of quartz along the existing foliation. The breakdown of muscovite is then accompanied by the growth of sillimanite. Quartz-plagioclase aggregates are replaced by assemblages that are in equilibrium with the granite eutectic point, although K-feldspar aggregates are preserved. The breakdown of biotite is responsible for the production of melt beyond 30-50 %, the value of the Rheological Critical Melt Percentage (Arzi, 1978). Leucosomes with cockade-type cordierite produced during this second melting stage tend to be discordant with the inherited structure. Structural orientations then

become more varied as the leucosome proportion increases, with folds becoming abundant and randomly oriented.

Stop D4.2:

Pont de Bayzan, Road from Le Puy-en-Velay to Aubenas, N102, River banks of the Ardèche river.

Stop D4.1 shows the incipient stage of melting within paragneiss of the Lower Gneissic Unit. The metagranite observed at the stop 4.1. is originally intrusive in sediments (pelites and argillites, including refractory quartz-rich and calcic layers) The location of early melting is controlled by foliation anisotropy (Macaudière et al., 1992) and folding (Barraud et al., 2003).

Numerous resistors from refractory layers preserve microstructure developed during the pre-migmatitic tectonic evolution that resulted in a composite foliation (named regionally S2) and polyphased folding (Color Plate 2, C). The outcrop is characterized by open folds that play an active role in the segregation of anatectic melts: cordierite-free leucosomes accumulate in saddle reef and axial planes of the folding that is attributed to S3 (Barraud et al., 2003).

Stop D4.3:

Ucel, Road from Aubenas to Mezilhac, Road-cut.

Stop D4.3 shows the unconformity of the Mesozoic sandstone over altered granite and biotite-sillimanite migmatitic paragneiss. The exhumation of the Velay granite-migmatite dome occurred during the Stephanian as boulders of granites and gneisses are found in the conglomerates of the Stephanian basin in the North and Prades basin in the South. Apatites in granite and migmatites yield a U-Pb age at 289 ± 5 Ma that is interpreted as a cooling age during the uplift of the Vealy region (Mougeot et al., 1997). Finally, a regional unconformity is characterized at the base of the Trias sandstone and conglomerate. A recent and fresh road cut provides a spectacular illustration of this unconformity.

Stop D4.4:

Volane river, Road from Aubenas to Mezilhac, D578, Road-cut.

Stop D4.4 shows an hololeucocratic granite, with cockade-type cordierite, that represents one of the petrographic type of the Velay migmatite-granite dome (Color Plate 2, D). Detailed observation indicates that cordierite is formed at the expense of biotite, in the presence of a melt phase. Cordierite is formed from the early phase of melting to the end of the magmatic

evolution of the Velay granite. However some garnets are present in cordierite nodules, indicating that, in that area, melting started in the garnet stability field.

Stop D4.5:

Volane river, Road from Aubenas to Mezilhac, D578, Road-cut.

Stop D4.5 shows a late-migmatitic homogeneous granite and relations with migmatitic gneisses and heterogeneous cordierite-bearing granite. In the upper part of the outcrop, a late-migmatitic granite, the Quatre-Vios granite, is intruding the heterogeneous banded granite: it is a coarse grained peraluminous granite with prismatic cordierite and frequently oriented feldspar phenocrysts. It contains abundant mafic microgranular enclaves and surmicaceous enclaves including biotite, garnet, cordierite, sillimanite, hercynite, ilmenite and rare plagioclase. This unusual mineralogy (absence of quartz and potash feldspar) and the corresponding chemical composition indicate that these enclaves are restites. P-T conditions calculated from this mineralogy yielded water-undersaturated conditions estimated at 4.5-5.5 kbar and 750-850 C. The mafic microgranular (biotite+plagioclase) enclaves correspond to frozen blobs of mafic magma. Locally, another type of late-migmatitic fine-grained granite, with typical acicular biotite and devoid of enclaves, crosscuts the Quatre-Vios granite.

In the lower part of the outcrop, another dyke of late-migmatitic granite is intrusive within migmatitic orthogneiss in which the second stage of melting is well marked by the biotite breakdown that produces Fe-rich garnet and cordierite. This zone is itself enclosed within the heterogeneous banded granite that contains a lot of enclaves of migmatitic paragneiss.

Stop D4.6:

Mont Gerbier-de-Jonc, Road from Mezilhac to Le Puy-en-Velay, D378, Sight as seen from the road.

The Velay volcanism is made up of an eastern chain of Mio-Pliocene basaltic to phonolitic volcanoes and a western Plio-Quaternary basaltic plateau (Mergoïl et al., 1993). The road to Le Puy-en-Velay shows nice sightseeing of this phonolitic chain that extends over more than 55 km with more than 180 points of extrusion, emplaced between 14 and 6 Ma.

The Mont Gerbier-de-Jonc is known as the spring of the Loire river. It is a phonolitic protrusion that displays a rough prismatic jointing.

Stop D4.7:

Moulin de Sezigneux, Road from St Chamond to Le Bessat, D2, Road cut.

The cordierite-andalusite-bearing micaschist of the Pilat Unit, seen at stop D4.7, displays extensional tectonics and HT-LP metamorphism within the Pilat series (Lower Gneissic Unit). This outcrop offers a good example of a low-pressure metamorphic unit, with micaschists and paragneisses intruded by syntectonic pegmatite dykes and pods. The main foliation plane is gently dipping to the North and shear criteria are well marked mainly around the pegmatitic pods indicating top to the North extensional tectonics. In the metamorphic rocks, the common mineralogy is: quartz + feldspars + biotite + andalusite and /or cordierite ± muscovite.

D4.8: Moulin de Sezigneux, Road from St Chamond to Le Bessat, D2, River banks.

Structure and metamorphism of the orthogneiss of the Lower Gneissic Unit are illustrated at stop D4.8. Ultramylonite and pseudotachylite textures are developed within the orthogneiss and S-C-C' structures indicate top to the North shear criteria compatible with extensional tectonics. Shear bands are underlined by syntectonic recrystallized biotites dated around 320-300 Ma (⁴⁰Ar/³⁹Ar). In the highly sheared parts of the outcrop ultramylonite bands and pseudotachylites are observable. The main foliation plane bears a north-south oriented stretching lineation. The metamorphic mineralogy is characterized by the association of quartz + feldspars + biotite + muscovite. Rare small-sized cordierites can also be observed.

End of the 4th day. Overnight stay in St Etienne

DAY 5

High to ultra-high pressure metamorphism and arc magmatism: records of subduction processes in the French Massif Central

The main goal of Day 5 is to present the geological and petrological records of subduction processes in the eastern Massif Central that preceded the development of the Velay migmatite-granite dome illustrated during Day 4. These records are: remnants of high to ultra-high pressure metamorphism in both crustal and mantle-derived lithologies in the Mont du Lyonnais unit, a partially preserved back-arc derived ophiolitic sequence, the Brévenne unit.




Figure 32 - Geological maps of the eastern margin of the Massif Central through time showing the progressive edification of the belt. The geodynamic cartoons show the possible position of the Monts du Lyonnais eclogites during the four critical periods illustrated (a to d).

A. Geological setting

1. Eclogites and garnet peridotite from The Monts du Lyonnais unit (Fig. 32a)

The Monts du Lyonnais unit belongs to the upper gneissic unit (Lardeaux, 1989; Ledru et al., 1994a). It comprises metasediments, orthogneisses (with protoliths of Ordovician age), leptynites (i.e. meta rhyolites), amphibolites and minor marbles. This unit also contains lenticular relics of either crustal (mafic and acid granulites, eclogites, Lasnier, 1968; Coffrant and Piboule, 1971; Dufour, 1985; Dufour et al., 1985; Lardeaux et al., 1989) or mantle origin (garnet and / or spinel bearing peridotites, Gardien et al., 1988, 1990). Eclogites outcrop, in close association with garnet-bearing peridotites, in the southernmost part of the Monts du Lyonnais unit. Eclogites and related garnet amphibolites also occur in a similar structural situation farther north (in the Morvan unit, Godard, 1990) and also southeast of the Monts du Lyonnais unit (in the Maclas-Tournon area, Gardien and Lardeaux, 1991). In the Monts du Lyonnais unit, three ductile strain patterns were distinguished (Lardeaux and Dufour, 1987; Feybesse et al., 1996) and related to high pressure and medium pressure metamorphic conditions:

- The relictual high-pressure structures
- A main deformation imprint, contemporaneous with amphibolite facies conditions, corresponds to a NW-SE crustal shortening with a finite NNE-SSW stretching direction (Fig. 32b)
- A deformation event developed under a transpressional regime dated between 335 and 350 Ma (Rb/Sr whole rock, Gay et al., 1981; $^{40}\text{Ar}/^{39}\text{Ar}$, Costa et al., 1993) which is correlated to the main deformation within the Brévenne ophiolite in relation with its overthrusting. (Fig. 32c)

With respect to this transpressive strain pattern, in the southern part of the Monts du Lyonnais unit, the eclogites outcrop exclusively in the strongly folded domains where they behave as rigid bodies in a deformed ductile matrix. We never found any eclogitic body within the shear zones.

2. The uppermost part of the magmatic arc: the Brévenne ophiolite (Fig. 32b)

The Devonian Brévenne ophiolitic unit consists of an association of metabasalts and metarhyolites together with intrusive intruded by trondhjemitic bodies (Peterlongo, 1970; Piboule et al., 1982, 1983). The ophiolitic unit was initially emplaced in a submarine environment (Pin et al., 1982; Delfour et al., 1989). These intercalations are cut and overlain by intrusive gabbros and dolerites and by submarine basaltic

lavas that, finally, are overlain by siltstones with pyroclastic intercalations (Milési et Lescuyer, 1989; Feybesse et al., 1996). A prograde greenschist to lower amphibolite facies metamorphism is recorded (Peterlongo, 1960; Fontailles, 1968; Piboule et al., 1982; Feybesse et al., 1988). The Brévenne ophiolite underwent a polyphase deformation. An early event, well developed in the northern part of the unit, is characterized by a NW-SE stretching lineation with top to the NW shearing (Leloix et al., 1999). During the second event, the ophiolitic unit is overthrusting the Monts du Lyonnais along a dextral transpressional zone in which syntectonic granites emplaced between 340 and 350 Ma (Fig. 32c, Gay et al., 1981; Feybesse et al., 1988; Costa et al., 1993). Subsequently, monzonitic granites of Namurian- Westphalian age and a contact metamorphism aureole postdate this tectonics (Delfour, 1989).

3. The development of the Velay migmatite granite dome and the collapse of the orogen (Fig. 32b)

The tectonic evolution of the eastern Massif Central is achieved during Westphalian and Stephanian. The formation of the Velay dome, coeval with the activation of crustal-scale detachments, potentially corresponds to flow of a partially molten crustal layer in response to gravitational collapse.

B. Stop description (Fig. 28, 29)

Stop D5.1:

The Bois des Feuilles, Road from St Symphorien-sur-Coise to Rive de Gier, D2.

This outcrop consists of garnet bearing peridotites and eclogites occurring as boudins within garnet sillimanite paragneisses. The coesite-bearing eclogite occurs in the southern part of the Monts du Lyonnais unit, near St Joseph in the Bozançon valley (1/50.000 geological map “ St Symphorien Sur Coise “, Feybesse et al., 1996) in association with “common” eclogites and serpentinites. In the whole area, eclogites are preserved in low-strain lenses (meter scale boudins) wrapped by amphibolites or amphibolite facies paragneisses. For practical reasons (difficult access), we shall observe only garnet-peridotites and “common eclogites”.

Well-preserved peridotites occur as metric to decametric scale bodies within the paragneisses. In the less retrogressed samples, garnets in equilibrium with olivine, clinopyroxenes and orthopyroxenes can be observed. Frequently, garnets contain inclusions of spinels and pyroxenes, while in some samples

fractionated members of a volcanic tholeiitic suite.

In the less retrogressed samples, the following mineral assemblages, representing the relicts of eclogite facies metamorphism, are recognized in the dark (1-2) and light (3-4) eclogites from the Monts du Lyonnais (Fig. 33):

- Garnet - omphacite - quartz - zoisite - rutile - apatite - sulfides,
- Garnet - omphacite - quartz - zoisite - colourless amphibole - rutile - sulfides,
- Garnet - omphacite - quartz (or coesite) - zoisite - kyanite - colorless amphibole - rutile,
- Garnet - omphacite - quartz - zoisite - kyanite - phengite - rutile.

Coesite and quartz pseudomorphs after coesite were exclusively detected as inclusions in two garnet grains within one sample of kyanite-bearing eclogite (Color Plate 2, E, F). SiO₂ polymorphs were distinguished optically, i.e. coesite was first positively identified relative to quartz by its higher refractive index, and then confirmed by Raman spectroscopy, by observation of the characteristic Raman lines 177, 271, 521 cm⁻¹. Only two coesite grains are preserved as relicts and, generally, coesite is otherwise completely transformed into polycrystalline radial quartz (palisade texture) or into polygonal quartz surrounded by radiating cracks. The extremely rare preservation of coesite in the Monts du Lyonnais eclogites is clearly the result of the high temperature conditions (near 750°C, see details in Dufour et al., 1985 and Mercier et al., 1991) reached during decompression as well as the consequence of fluid influx (hydration) during retrogression. Indeed, the kinetics of the coesite → quartz transformation are strongly temperature and fluid dependent (Gillet et al., 1984; Van der Molen and Van Roermund, 1986; Hacker and Peacock, 1995; Liou and Zhang, 1996) and consequently in the studied area, coesite has been almost entirely transformed into quartz.

Stop D5.2:

St-André-la-Côte, Road from St-André-en-Haut to Mornant.

Migmatites and granulitic rocks outcrop in the northern part of the Monts du Lyonnais. Near St-André-la-Côte village, mafic and acid granulite facies rocks are well exposed. In mafic granulites the following metamorphic assemblages are described (Dufour, 1985; Dufour et al., 1985):

- Garnet + plagioclase + orthopyroxene + ilmenite,
- Garnet + plagioclase + amphibole + ilmenite,

- Clinopyroxene + orthopyroxene + plagioclase ± amphibole + ilmenite,

- Garnet + clinopyroxene + orthopyroxene + plagioclase ± amphibole + ilmenite.

In acid granulites (Dufour, 1982; Lardeaux et al., 1989), the common mineralogy consists of quartz + plagioclase + K-feldspar + garnet + sillimanite ± spinel ± biotite. Kornerupine-bearing granulites have been also locally recognized in this outcrop (Lardeaux et al., 1989).

In this northern part of the Mont du Lyonnais unit, the metamorphic imprint is typical for Intermediate Pressure granulite facies and there is no trace of eclogitic high-pressure facies metamorphism.

Stop D5.3:

The Yzeron quarry, Road from Ste-Foy-l'Argentière to Craponne, D489.

Stop D5.3 shows the migmatitic orthogneisses from the northern Monts du Lyonnais. These rocks are metamorphosed and strongly foliated under amphibolite facies conditions. Their typical mineralogy is an association of quartz + plagioclase + K-feldspar + biotite ± sillimanite. Muscovite and chlorite are developed during retrogression. In the Yzeron quarry, superposed fold systems have been described in relation with progressive deformation under amphibolite and greenschist facies metamorphic conditions. The main foliation observed in the orthogneisses is folded and / or reworked in the ductile strike-slip shear zones related to the development of a regional scale transpressive regime.

The northern Monts du Lyonnais area, with intermediate-pressure granulite and amphibolite facies metamorphic rocks can be interpreted as remnants of the upper overriding continental crust on which the magmatic arc and the Brévenne back-arc were emplaced. In this model, the southern part of the Monts du Lyonnais unit corresponds to a subduction complex (remnant of subduction channel) located on the top of the lower plate (i.e. Pilat – Velay units).

Stop D5.4:

Brévenne valley, road cut: bimodal magmatic sequence.

Stop D5.4 shows different lithologies, like metabasalts, metarhyolites, metapyroclastites and metasediments, typical for the Brévenne ophiolitic unit. All the lithologies are metamorphosed under greenschist facies conditions. In mafic lithologies, the common mineralogy corresponds to an association of

plagioclase + actinolite + chlorite + sphene \pm calcite \pm quartz.

The different lithologies are also deformed and involved into a regional scale fold system with sub-vertical axial planes. These folds are related to the regional transpressive regime which affects also the Monts du Lyonnais unit at around 350 – 340 Ma.

Recent geochemical investigations support the origin of the Brévenne ophiolitic sequence, during Devonian, in a back-arc basin developed upon the upper plate of a subduction system.

End of the 5th day in Lyon (airport and railway station)

Acknowledgements

O. Monod is thanked for his review of the first draft of this guidebook. Mrs S. Matrat and D. Quiniou provided a great help during the final stages for preparing this guidebook.

References cited

- Arzi, A.A. (1978). Critical phenomena in the rheology of partially melted rocks. *Tectonophysics* 44, 173-184.
- Ait Malek, H. (1997). Pétrologie, Géochimie et géochronologie U/Pb d'associations acide-basiques: exemples du SE du Velay (Massif central français) et de l'anti-Atlas occidental (Maroc). Thèse doctorat de l'INPL, Univ. Nancy, 297 pp.
- Ait Malek, H., Gasquet, D., Marignac, C. and Bertrand, J.M. (1995). Des xénolites à corindon dans une vaugnèrite de l'Ardèche (Massif central français) : implications pour le métamorphisme ardéchois. *C. R. Acad. Sci. Paris* 321, 959-966.
- Arnaud, F. (1997). Analyse structurale et thermo-barométrique d'un système de chevauchements varisque : les Cévennes centrales (Massif Central français). Microstructures et mécanismes de déformation dans les zones de cisaillement schisteuses. Thèse 3^{ème} cycle, Institut National Polytechnique de Lorraine, Documents du BRGM, 286, 351 pp.
- Arnaud, F. and Burg, J.P. (1993). Microstructures des mylonites schisteuses: cartographie des chevauchements varisques dans les Cévennes et détermination de leur cinématique. *C. R. Acad. Sci. Paris* 317, 1441-1447.
- Arthaud, F. (1970). Etude tectonique et microtectonique comparée de deux domaines hercyniens : les nappes de la Montagne Noire (France) et l'anticlinorium de l'Iglesiente (Sardaigne). Thèse d'Etat, Univ. Montpellier, France, 175pp.
- Arthaud F. and Matte P. (1977) - Late Paleozoic strike slip faulting in southern Europe and northern Africa: result of right lateral shear zone between Appalachians and the Urals. *Geol. Soc. Am. Bull.* 88, p. 1305 -1320.
- Autran, A. and Cogné, J. (1980). La zone interne de l'orogénèse varisque dans l'Ouest de la France et sa place dans le développement de la chaîne hercynienne. (J. Cogné J. and M. Slansky M. Eds.), *Géologie de l'Europe du Précambrien aux bassins sédimentaires post-hercyniens*, 26^{ème} Cong. Géol. Int., Coll. C6, Paris 1980. *Ann. Soc. géol. Nord*, Lille XCIX, 90-111.
- Barbey, P., Marignac, C., Montel, J.M., Macaudière, J., Gasquet, D. and Jabbori, J. (1999). Cordierite growth texture and the conditions of genesis and emplacement of crustal granitic magmas: the Velay granite complex (Massif Central, France). *J. Petrology* 40, 1425-1441.
- Barraud, J., Gardien, V., Allemand, P. and Grandjean, P. (2003). Analog models of melt-flow in folding migmatites. *J. Struct. Geol.* in press.
- Be Mezème, E. (2002). Application de la méthode de datation à la microsonde électronique de monazite de migmatites et de granitoïdes tardi-hercyniens du Massif Central français. Master thesis, Univ. Orléans, 40 pp.
- Bernard-Griffiths, J., Cantagrel J.M. and Duthou J.L. (1977) - Radiometric evidence for an Acadian tectono-metamorphic event in western Massif Central français. *Contrib. Mineral. Petrol.* 61, 199-212.
- Berthé, D., Choukroune, P. and Jegouzo, P. (1978). Orthogneiss mylonite and non coaxial deformation of granite: the exemple of the South armorican shear zone. *J. Struct. Geol.* 1, 31-42.
- Bitri, A., Truffert, C., Bellot, J.P., Bouchot, V., Ledru, P., Milési, J.P. and Roig J.Y. (1999) - Imagerie des paléochamps hydrothermaux As-Sb d'échelle crustale et des pièges associés dans la chaîne varisque : sismique réflexion verticale (GéoFrance 3D : Massif central français). *C. R. Acad. Sci. Paris* 329, 771-777.
- Blanc, D. (1981). Les roches basiques et ultrabasiqes des monts du Lyonnais. Etude pétrographique, minéralogique et géochimique. Thèse Doctorat 3^{ème} cycle, Univ. Lyon 1, 152 p.
- Bouchez, J.L., and Jover, O. (1986). Le Massif Central : un chevauchement de type himalayen vers l'ouest-nord-ouest. *C. R. Acad. Sci. Paris* 302, 675-680.
- Boutin, R. and Montigny, R. (1993). Datation

- $^{39}\text{Ar}/^{40}\text{Ar}$ des amphibolites du complexe leptyno-amphibolique du plateau d'Aigurande : collision varisque à 390 Ma dans le Nord-Ouest du Massif central français. *C. R. Acad. Sci. Paris* 316, 1391-1398.
- Brown, M. (1994). The generation, segregation, ascent and emplacement of granite magma: the migmatite-to-crustally-derived granite connection in thickened orogens. *Earth Science Reviews* 36, 83-130.
- Brown, M. and Dallmeyer, R.D. (1996). Rapid Variscan exhumation and the role of magma in core complex formation: southern Brittany metamorphic belt. *J. metamorphic Geol.* 14, 361-379
- Burg, J.P., Leyreloup, A., Marchand, J. and Matte, P. (1984). Inverted metamorphic zonation and large-scale thrusting in the Variscan belt: an example in the French Massif Central. In: "Variscan tectonics of the North-Atlantic region" (D.H.W. Hutton, and D. J. Sanderson, Ed.), pp. 47-61, Spec. Publ. Geol. Soc. London, 14.
- Burg, J.P. Bale, P., Brun, J.P. and Girardeau, J. (1987). Stretching lineation and transport direction in the Ibero-Armorican arc during the siluro-devonian collision. *Geodinamica Acta* 1, 71-87.
- Burg J.P., Brun, J.P. and Van Den Driessche, J. (1991) - Le Sillon Houiller du Massif central français : faille de transfert pendant l'amincissement crustal de la chaîne varisque. *C. R. Acad. Sci. Paris* 311, II, 147-152.
- Burg, J.P. and Vanderhaeghe, O. (1993). Structures and way-up criteria in migmatites, with application to the Velay dome (French Massif central). *J. Struct. Geol.* 15, 1293-1301.
- Burg, J.P., Van Den Driessche, J. and Brun, J.P. (1994). Syn- to post-thickening extension: mode and consequences. *C. R. Acad. Sci. Paris* 319, 1019-1032.
- Caen Vachette, M., Couturié, J.P. and Didier, J. (1982). Age radiométrique des granites anatectiques et tardimigmatitiques du Velay (Massif Central français). *C. R. Acad. Sci. Paris* 294, 135-138.
- Caen Vachette, M., Gay, M., Peterlongo, J.M., Pitiot, P. and Vitel, G. (1984). Age radiométrique du granite syntectonique du gouffre d'Enfer et du métamorphisme hercynien dans la série de basse pression du Pilat (Massif Central Français). *C. R. Acad. Sci. Paris* 299, 1201-1204.
- Caron, C., Lancelot, J.R., and Maluski, H. (1991). A paired ^{40}Ar - ^{39}Ar and U-Pb radiometric analysis applied to the variscan Cévennes, french Massif central. EUG Strasbourg, *Terra abstracts* 3, 205.
- Clemens, J.D. (1990). The granulite-granite connexion. In: "Granulites and Crustal Evolution" (D. Vielzeuf and Ph. Vidal, Eds.), Kluwer Acad. Publ., pp. 25-36.
- Coffrant, D. (1974). Les éclogites et les roches basiques et ultrabasiques associées du massif de Sauviat-sur-Vige, Massif central français. *Bulletin de la Société Française de Minéralogie Cristallographie* 97, 70-78
- Coffrant, D. and Piboule, M. (1971). Les éclogites et les roches associées des massifs basiques de Saint-Joseph (Monts du Lyonnais, Massif Central français). *Bull. Soc. Géol. Fr.* 7, XIII, 283-291
- Couturier, J.P. (1969). Le massif granitique de la Margeride. Thèse d'Etat, Univ. Clermont-Ferrand, France, 190 pp.
- Costa, S. (1989). Age radiométrique $^{39}\text{Ar}/^{40}\text{Ar}$ du métamorphisme des séries du Lot et du charriage du groupe leptyno-amphibolique de Mavejols. *C. R. Acad. Sci. Paris* 309, 561-567.
- Costa S. (1992). East-West diachronism of the collisional stage in the French Massif Central: implications for the european variscan orogen. *Geodinamica Acta* 5, 51-68.
- Costa, S., Maluski, H. and Lardeaux, J.M. (1993). ^{40}Ar - ^{39}Ar chronology of Variscan tectono-metamorphic events in an exhumed crustal nappe: the Monts du Lyonnais complex (Massif Central, France). *Chem. Geol.* 105, 339-359.
- Dallain, C., Schulmann, K. and Ledru, P. (1999). Textural evolution in the transition from subsolidus annealing to melting process, Velay dome, French Massif Central. *J. Metamorphic Geol.* 17, 61-74.
- Delfour, J. (1989). Données lithostratigraphiques et géochimiques sur le Dévono-Dinantien de la partie sud du faisceau du Morvan (nord-est du Massif Central français). *Géologie de la France* 4, 49-77.
- Demange, M. (1975). Style pennique de la zone axiale de la Montagne Noire entre Saint-Pons et Murat-sur-Vèbre (Massif Central). *Bull. BRGM* 2, 91-139.
- Demange, M. (1985). The eclogite facies rocks of the Montagne Noire, France. *Chemical Geol.* 50, 173-188.
- Didier, J. (1973). Granites and their enclaves. The bearing of enclaves on the origin of granites, 2, Developments in *Petrology Series*, Amsterdam, Elsevier, 3, 37-56.
- Didier, J. and Lameyre, J. (1971). Les roches granitiques du Massif central. In: Symposium J. Jung: "Géologie, géomorphologie et structure profonde du Massif central français", pp. 17-32,

Clermont-Ferrand, Plein Air Service.

Dubuisson, G., Mercier, J.C.C., Girardeau, J. and Frison, J.Y. (1989). Evidence for a lost ocean in Variscan terranes of the western Massif Central, France. *Nature* 337, 23, 729-732.

Ducrot, J., Lancelot, J.R. and Marchand, J. (1983). Datation U-Pb sur zircons de l'éclogite de la Borie (Haut-Allier, France) et conséquences sur l'évolution anté-hercynienne de l'Europe Occidentale. *Earth Planet. Sci. Lett.* 18, 97-113.

Dufour, E. (1982). Pétrologie et géochimie des formations orthométamorphiques acides des Monts du Lyonnais (Massif Central français). Thèse Doctorat de 3^{ème} cycle, Univ. Lyon 1, 241 p.

Dufour, E. (1985). Granulite facies metamorphism and retrogressive evolution of the Monts du Lyonnais metabasites (Massif Central France) *Lithos* 18, 97-113

Dufour, E., Lardeaux, J.M. and Coffrant, D. (1985). Eclogites and granulites in the Monts du Lyonnais area: an eo-Hercynian plurifacial metamorphic evolution. *C. R. Acad. Sci. Paris* 300, 141-144

Dupraz, J. and Didier, J. (1988). Le complexe anatectique du Velay (Massif Central français) : structure d'ensemble et évolution géologique. *Géologie de la France* 4, 73-87.

Duthou, J.L., Cantagrel, J.M., Didier, J. and Vialette, Y. (1984). Paleozoic granitoids from the French Massif Central: age and origin studied by ⁸⁷Rb/⁸⁷Sr system. *Phys. Earth Planet. Int.* 35, 131-144.

Duthou, J-L., Chenevoy, M. and Gay M. (1994). Age Rb/Sr Dévonien moyen des migmatites à cordiérite du Lyonnais (Massif Central français). *C. R. Acad. Sci. Paris*. 319, 791-796.

Echtler, H. and Malavieille, J. (1990). Extensional tectonics, basement uplift and Stephano-Permian collapse basin in a Late Variscan metamorphic core complex (Montagne Noire, Southern Massif Central). *Tectonophysics* 177, 125-138.

Engel, W., Feist, R. and Franke, W. (1980). Le Carbonifère anté-stéphanien de la Montagne Noire : rapports entre mise en place des nappes et sédimentation. *Bull. BRGM* 2, 341-389.

Faure, M. (1995). Late orogenic carboniferous extensions in the Variscan French Massif Central. *Tectonics* 14, 132-153.

Faure, M., Pin, C. and Mailhé, D. (1979). Les roches mylonitiques associées au charriage du groupe leptyno-amphibolique sur les schistes du Lot dans la région de Marvejols (Lozère). *C. R. Acad. Sci. Paris* 288, 167-170.

Faure, M. and Cotterau, N. (1988). Données cinématiques sur la mise en place du dôme migmatitique carbonifère moyen de la zone axiale de la Montagne Noire (Massif Central, France). *C. R. Acad. Sci. Paris* 307, II, 1787-1794.

Faure, M., Leloix, C. and Roig, J.Y. (1997). L'évolution polycyclique de la chaîne hercynienne. *Bull. Soc. Geol. France* 168, 695-705.

Faure, M., Monié, P., Maluski, H., Pin, C. and Leloix, C. (2001). Late Visean thermal event in the northern part of the French Massif Central. New ⁴⁰Ar/³⁹Ar and Rb-Sr isotopic constraints on the Hercynian synorogenic extension. *Int. J. Geol.* 91, 53-75.

Feist, R. and Galtier, J. (1985). Découverte de flores d'âge namurien probable dans le flysch à olistolithes de Cabrières (Hérault). Implications sur la durée de la sédimentation synorogénique dans la Montagne Noire (France Méridionale), *C. R. Acad. Sci. Paris* 300, 207-212.

Feybesse, J.L., Lardeaux, J.M., Johan, V., Tegye, M., Dufour, E., Lemiere, B. and Delfour, J. (1988). La série de la Brévenne (Massif Central français): une unité dévonienne charriée sur le complexe métamorphique des Monts du Lyonnais à la fin de la collision varisque. *C. R. Acad. Sci. Paris* 307, 991-996.

Feybesse, J.L., Lardeaux, J.M., Tegye, M., Kerrien, Y., Lemiere, B., Mercier, F., Peterlongo, J.M. and Thieblemont, D. (1996). Carte géologique de France (1/50000), feuille St Symphorien-sur-Coise (721). BRGM Orléans.

Floc'h, J-P. (1983). La série métamorphique du Limousin central. Thèse d'Etat, Univ. Limoges, France, 445pp.

Fontelles, M. (1968). Contribution à l'analyse du processus de spilitisation. Etude comparée des séries volcaniques paléozoïques de la Bruche (Vosges) et de la Brévenne (Massif Central français). *Bull. BRGM* 2, (3), 1-54

Franke, W. (1989). Tectonostratigraphic units in the Variscan belt of central Europe. In "Terranes in the circum-Atlantic Paleozoic orogens" (R.D. Dallmeyer Ed.), pp. 67-90. Special paper, Geological Society of America, 230.

Franke, W. (2000). The mid-European segment of the Variscides: tectonostratigraphic units, terrane boundaries and plate tectonic evolution, in Orogenic Processes. In "Quantification and Modelling in the Variscan Belt" (W. Franke, V. Haak, O. Oncken, D. Tanner, Eds.) pp. 35-61. Special Publications, 179, Geological Society of London.

- Gardien, V., Lardeaux, J.M. and Misseri, M. (1988). Les péridotites des Monts du Lyonnais (Massif Central français) : témoins privilégiés d'une subduction de lithosphère paléozoïque. *C. R. Acad. Sci. Paris* 307, 1967-1972.
- Gardien, V. (1990). Reliques de grenat et de staurotite dans la série métamorphique de basse pression du Mont Pilat (Massif Central français): témoins d'une évolution tectonométamorphique polyphasée. *C. R. Acad. Sci. Paris* 310, 233-240.
- Gardien, V., Tegye, M., Lardeaux, J.M., Misseri, M. and Dufour, E. (1990). Crustal-mantle relationships in the french Variscan chain: the example of the Southern Monts du Lyonnais unit (eastern French Massif Central). *Journ. Metam. Geol.* 8, 477-492.
- Gardien, V. and Lardeaux, J.M. (1991). Découvertes d'éclogites dans la synforme de Maclas: extension de l'Unité Supérieure des Gneiss à l'Est du massif central. *C. R. Acad. Sci. Paris* 312, 61-68.
- Gardien, V., Lardeaux, J.M., Ledru, P., Allemand, P. and Guillot, S. (1997). Metamorphism during late orogenic extension: insights from the French Variscan belt. *Bull. Soc. Géol. Fr.* 168, 271-286.
- Gèze, B. (1949). Etude Géologique de la Montagne Noire et des Cévennes Méridionales. *Mem. Soc. Géol. France* 24, 215.
- Gay, M., Peterlongo, J.M. and Caen-Vachette, M. (1981). Age radiométrique des granites en massifs allongés et en feuillets minces syn-tectoniques dans les Monts du Lyonnais (Massif Central français). *C. R. Acad. Sci. Paris* 293, 993-996.
- Gillet, P., Ingrin, J. and Chopin, C. (1984). Coesite in subducted continental crust: *P-T* history deduced from an elastic model. *Earth Planet. Sci. Lett.* 70, 426-436
- Godard, G. (1990). Découverte d'éclogites, de péridotites à spinelle et d'amphibolite à anorthite, spinelle et corindon dans le Morvan. *C. R. Acad. Sci. Paris* 310, 227-232.
- Hacker, B.R. and Peacock, S.M. (1995). Creation, preservation, and exhumation of UHPM rocks. In: "Ultra-high-Pressure Metamorphism". (Coleman, Wang, Eds.), Cambridge University Press, Cambridge, pp. 159-181
- Lagarde, J.L., Dallain, C., Ledru, P. and Courrioux, G. (1994). Deformation localization with laterally expanding anatectic granites: Hercynian granites of the Velay, French Massif Central. *J. Struct. Geol.* 16, 839-852.
- Lardeaux, J.M. (1989). Les formations métamorphiques des Monts du Lyonnais *Bull. Soc. Géol. Fr.* 4, 688-690
- Lardeaux, J.M. and Dufour, E. (1987). Champs de déformation superposés dans la chaîne varisque. Exemple de la zone nord des Monts du Lyonnais (Massif Central français). *C. R. Acad. Sci. Paris* 305, 61-64.
- Lardeaux, J.M., Reynard, B. and Dufour, E. (1989). Granulites à kornéropine et décompression post-orogénique des Monts du Lyonnais. *C. R. Acad. Sci. Paris II* 308, 1443-1449
- Lardeaux, J.M., Ledru, P., Daniel, I. and Duchène, S. (2001). The variscan French Massif Central - a new addition to the ultra-high pressure metamorphic «club»: exhumation processes and geodynamic consequences. *Tectonophysics* 323, 143-167.
- Lasnier, B. (1968a). Découverte de roches éclogitiques dans le groupe leptyno-amphibolique des Monts du Lyonnais. *Bull. Soc. Géol. Fr.* 7, 179-185
- Ledru, P., Lardeaux, J.M., Santallier, D., Autran, A., Quenardel, J-M., Floc'h, J-P., Lerouge, G., Maillat, N., Marchand, J. and Ploquin, A. (1989). Où sont les nappes dans le Massif Central français ? *Bull. Soc. Géol. France* 8, 605-618.
- Ledru, P., Autran, A. and Santallier, D. (1994a). Lithostratigraphy of Variscan terranes in the French Massif Central. A basic for paleogeographical reconstruction. In: "Pre-Mesozoic geology in France and related areas", (J. D. Keppie, Ed.) pp. 276-288. Springer Verlag.
- Ledru, P., Costa, S. and Echtler, H. (1994b). Structure. In: "Pre-Mesozoic geology in France and related areas", (J. D. Keppie, Ed.) pp. 305-323, Springer Verlag.
- Ledru, P., Courrioux, G., Dallain, C., Lardeaux, J.M., Montel, J.M., Vanderhaeghe, O., and Vitel, G. (2001). The Velay dome (French Massif Central): melt generation and granite emplacement during orogenic evolution. *Tectonophysics* 332, 207-237.
- Leloix, C., Faure, M. and Feybesse, J.L. (1999). Hercynian polyphase tectonics in north-east French Massif Central : the closure of the Brévenne Devonian-Dinantian rift. *Int. J. Earth. Sci.* 88, 409-421.
- Liou, J.G. and Zaang, R.Y. (1996). Occurrences of intergranular coesite in ultrahigh-P rocks from the Sulu region, eastern China: implications of lack of fluid during exhumation. *Am. Mineralogist* 81, 1217-1221
- Macaudière, J., Barbey, P., Jabbori, J. and Marignac, C. (1992). Le stade initial de fusion dans le développement des dômes anatectiques : le dôme du Velay (Massif Central français). *C. R. Acad. Sci. Paris* 315, 1761-1767.

- Malavieille, J., Guihot, P., Costa, S., Lardeaux, J.M., and Gardien, V. (1990). Collapse of the thickened Variscan crust in the French Massif Central: Mont Pilat extensional shear zone and St Etienne upper Carboniferous basin. *Tectonophysics* 177, 139-149.
- Mattauer, M., Brunel, M. and Matte, P. (1988). Failles normales ductiles et grands chevauchements : une nouvelle analogie entre l'Himalaya et la chaîne hercynienne du Massif français. *C. R. Acad. Sci. Paris* 306, 671-676.
- Mattauer, M., Laurent P. and Matte P. (1996). Plissements hercyniens synschisteux post-nappe et étirement subhorizontal dans le versant Sud de la Montagne Noire. *C. R. Acad. Sci. Paris*. 322, 309-315.
- Mattauer, M. and Matte, P. (1998). Le bassin stéphanien de St-Etienne ne résulte pas d'une extension tardi-hercynienne généralisée : c'est un bassin pull-apart en relation avec un décrochement dextre. *Geodinamica Acta* 11, 23-31.
- Matte, P. (1991). Tectonics and plate tectonics model for the variscan belt of Europe. *Tectonophysics* 126, 329-374.
- Matte, P. (2001). The Variscan collage and orogeny (480-290 Ma) and the tectonic definition of the Armorica microplate : a review. *Terra Nova* 13, 122-1128.
- Matte, P., Lancelot, J-R. and Mattauer, M. (1998). La zone axiale hercynienne de la Montagne Noire n'est pas un "metamorphic core complex" extensif mais un anticlinal post-nappe à cœur anatectique. *Geodinamica Acta* 11, 13-22.
- Maluski, H., Costa, S. and Echler H. (1991). Late Variscan tectonic evolution by thinning of an earlier thickened crust. An $^{40}\text{Ar}/^{39}\text{Ar}$ study of the Montagne Noire, southern Massif central, France. *Lithos* 26, 287-304.
- Mercier, L., Lardeaux, J.M. and Davy, P. (1991). On the tectonic significance of the retromorphic P-T paths of the french Massif Central eclogites. *Tectonics* 10, 131-140.
- Milési, J.P. and Lescuyer J.L. (1989). The Chessy Zn-Cu-Ba massive sulphide deposit and the Devonian Brévenne volcano-sedimentary belt (eastern Massif Central, France). Project: identification of diagnostic markers of high-grade massive sulphide deposits of their enriched zones in France and in Portugal. CEE contrat MA IM-0030-F(D). Rapp. BRGM 89 DAM 010 DEX (final report)
- Mergoïl, J., Boivin, P., Blès, J.L., Cantagrel, J.M. and Turland M. (1993). Le Velay. Son volcanisme et les formations associées, notice de la carte à 1/100000. *Géologie de la France* 3, 3-96.
- Monié, P., Respaut, J.-P., Bricaud, S., Bouchot, V., Faure, M. and Roig, J.-Y. (2000). $^{40}\text{Ar}/^{39}\text{Ar}$ and U-Pb geochronology applied to Au-W-Sb metallogenesis in the Cévennes and Châtaigneraie districts (Southern Massif Central, France). In: "Orogenic gold deposits in Europe", (V. Bouchot, Ed.), pp. 77-79. Document BRGM 297, Bureau de Recherches Géologiques et Minières, Orléans.
- Montel, J.M., 1985. Xénolithes peralumineux dans les dolérites du Peyron, en Velay (Massif Central français). Indications sur l'évolution de la croûte profonde tardihercynienne. *C. R. Acad. Sci. Paris* 301, 615-620.
- Montel, J.M., Weber, C., Barbey, P. and Pichavant, M. (1986). Thermobarométrie du domaine anatectique du Velay (Massif Central français) et conditions de genèse des granites tardi-migmatitiques. *C. R. Acad. Sci. Paris* 302, 647-652.
- Montel, J.M. and Abdelghaffar, R. (1993). Les granites tardi-migmatitiques du Velay (Massif Central): principales caractéristiques pétrographiques et géochimiques. *Géologie de la France* 1, 15-28.
- Montel, J.M., Bouloton, J., Veschambre, M., Pellier, C. and Ceret, K. (2002). Age des microgranites du Velay (Massif Central Français). *Géologie de la France* 1, 15-20.
- Montel, J.M., Marignac, C., Barbey, P. and Pichavant, M. (1992). Thermobarometry and granite genesis : the Hercynian low-P, high-T Velay anatectic dome (French Massif Central). *J. Metam. Geol.* 10, 1-15.
- Mougeot, R., Respaut, J.P., Ledru, P. and Marignac, C. (1997). U-Pb chronology on accessory minerals of the Velay anatectic dome (French Massif Central). *Eur. J. Mineral.* 9, 141-156.
- Nicolas, A., Bouchez, J-L. Blaise, J. and Poirier, J-P. (1977). Geological aspects of deformation in continental shear zones. *Tectonophysics* 42, 55-73.
- Patiño Douce, A.E. and Johnston, A.D. (1990). Phase equilibria and melt productivity in the pelitic system: implications for the origin of peraluminous granitoids and aluminous granulites. *Contrib. Mineral. Petrol.* 107, 202-218.
- Peterlongo, J.M. (1960). Les terrains cristallins des monts du Lyonnais (Massif Central français). *Ann. Fac. Sci. Univ. Clermont-Ferrand* 4, (1), 187
- Peterlongo, J.M. (1970). Pillows-lavas à bordure variolitique et matrice basique dans la série métamorphique de la Brévenne (Rhône, Massif Central Français). *C. R. Acad. Sci. Paris* 2, 190-194


- Piboule, M., Briand, B. and Beurrier, M. (1982). Géochimie de quelques granites albitiques dévoniens de l'Est du Massif Central (France). *Neues Jb. Miner. Abh.*, 143, 279-308.
- Piboule, M., Beurrier, M., Briand, B. and Lacroix, P. (1983). Les trondhjemitites de Chindo et de St-Veran et le magmatisme kérotyphirique associé. Pétrologie et cadre géostructural de ce magmatisme Dévono-Dinantien. *Géologie de la France* 1, 2, (1-2), 55-72
- Piboule, M. and Briand, B. (1985). Geochemistry of eclogites and associated rocks of the southeastern area of the French Massif Central: origin of the protoliths. *Chem. Geol.* 50, 189-199
- Pin, C. (1979). Géochronologie U-Pb et microtectonique des séries métamorphiques anté-stéphaniennes de l'Aubrac et de la région de Marvejols (Massif Central). Thèse 3^o cycle, Univ. Montpellier, France, 220pp.
- Pin, C. (1981). Old inherited zircons in two synkinematic variscan granitoids: the "granite du Pinet" and the "orthogneiss de Marvejols" (southern French Massif Central). *N. Jb. Miner. Abh.* 142, 27-48.
- Pin, C. (1990). Variscan oceans: ages, origins and geodynamic implications inferred from geochemical and radiometric data. *Tectonophysics* 177, 215-227.
- Pin, C. and Lancelot, J. (1978). Un exemple de magmatisme cambrien dans le Massif Central : les métadiorites quartzites intrusives dans la série du Lot. *Bull. Soc. Géol. France* 7, 203-208.
- Pin, C. and Lancelot, J. (1982). U-Pb dating of an early paleozoic bimodal magmatism in the French Massif Central and of its further metamorphic evolution. *Contrib. Mineral Petrol.* 79, 1-12.
- Pin, C., Dupuy, C. and Peterlongo, JM. (1982). Répartition des terres rares dans les roches volcaniques basiques dévono-dinantiennes du nord-est du Massif central. *Bull. Soc. Géol. Fr.* 7, 669-676.
- Pin, C. and Vielzeuf, D. (1983). Granulites and related rocks in Variscan median Europe: a dualistic interpretation. *Tectonophysics* 93, 47-74.
- Pin, C. and Duthou, J.L. (1990). Sources of Hercynian granitoids from the French Massif Central: inferences from Nd isotopes and consequences for crustal evolution. *Chemical Geology* 83, 281-296.
- Pin, C. and Marini, F. (1993). Early Ordovician continental break-up in Variscan Europe: Nd-SR isotope and trace element evidence for bimodal igneous associations of the southern Massif Central, France. *Lithos* 29, 177-196.
- Pin, C. and Paquette, JL (1998). A mantle-derived bimodal suite in the Hercynian Belt: Nd isotope and trace element evidence for a subduction-related rift origin of the Late Devonian Brèvenne metavolcanics, Massif Central (France). *Contrib Mineral Petrol* 129, 222-238.
- Quénardel, J.M. and Rolin, P. (1984). Paleozoic evolution of the Plateau d'Aigurande (N-W. Massif Central, France). In "Variscan tectonics of the North Atlantic region" (D. Hutton and D. Sanderson, Eds.), pp. 63-77, Geol. Soc. London Spec. pub, 14.
- R'Kha Chaham, K., Couturié, J.P., Duthou, J.L., Fernandez, A. and Vitel, G. (1990). L'orthogneiss ocellé de l'Arc de Fix : un nouveau témoin d'âge cambrien d'un magmatisme hyper alumineux dans le Massif Central français. *C. R. Acad. Sci. Paris* 311, 845-850.
- Robardet, M., Verniers, J., Feist R. and Paris, F. (1994). Le Paléozoïque anté-varisque de France, contexte paléogéographique et géodynamique. *Géol. de la France* 3, 3-31.
- Robardet, M. (2003). The Armorica 'microplate': fact or fiction? Critical review of the concept and contradictory palaeobiogeographical data. *Palaeogeography, Palaeoclimatology, Palaeoecology* 195, 125-148.
- Roig, J-Y. and Faure M. (2000). La tectonique cisailante polyphasée du Sud-Limousin (Massif Central français) et son interprétation dans un modèle d'évolution polycyclique de la chaîne hercynienne. *Bull. Soc. Géol. Fr.* 171, 295-307.
- Roig, J-Y., Faure, M. and Truffert, C. (1998). Folding and granite emplacement inferred from structural, strain, TEM, and gravimetric analyses: the case study of the Tulle antiform, SW French Massif Central. *J. Struct. Geol.* 20, 1169-1189.
- Sider, J-M. and Ohnenstetter, M. (1986). Field and petrological evidence for the development of an ensialic marginal basin related to the Hercynian orogeny in the Massif Central, France. *Geol. Rundschau* 75, 421-443.
- Soula, J.C., Debat, P., Brusset, S., Bessière, G., Christophoul, F. and Déramond, J. (2001). Thrust related, diapiric and extensional doming in a frontal orogenic wedge: example of the Montagne Noire, southern French Hercynian Belt. *J. Struct. Geol.* 23, 1677-1699.
- Van den Driessche J. and Brun, J-P. (1991-92). Tectonic evolution of the Montagne Noire (French Massif Central): a model of extensional gneiss dome. *Geodinamica Acta* 5, 85-99.
- Vanderhaeghe, O., Burg, J.P. and Teyssier, C. (1999).

- Exhumation of migmatites in two collapsed orogens: Canadian Cordillera and French Variscides. In: "Exhumation processes: normal faulting, ductile flow and erosion" U. Ring, M.T. Brandon, G.S. Lister and S.D. Willett (Eds.) 181-204, *Geological Society, London, Special Publications*, 154,
- Vanderhaeghe, O. and Teyssier, C. (2001). Partial melting and flow of orogens. *Tectonophysics* 342, 451-472.
- Van der Molen, I. and Van Roermund, H.L.M. (1986) The pressure path of solid inclusions in minerals: the retention of coesite inclusions during uplift. *Lithos* 19, 317-324
- Vitel, G. (1985). La transition faciès granulite faciès amphibolite dans les enclaves basiques du Velay. *C. R. Acad. Sci. Paris* 300, 407-412.
- Williamson, B.J., Downes, H. and Thirlwall, M.F. (1992). The relationship between crustal magmatic underplating and granite genesis: an example from the Velay granite complex, Massif Central, France. *Trans. Royal Soc. Edinburgh, Earth Sciences* 83, 235-245.

Back Cover:
field trip itinerary

FIELD TRIP MAP

32nd INTERNATIONAL GEOLOGICAL CONGRESS



Edited by APAT