




ISPRA

Istituto Superiore per la Protezione
e la Ricerca Ambientale



Sistema Nazionale
per la Protezione
dell'Ambiente

Italian Emission Inventory 1990 -2020. Informative Inventory Report 2022



RA P P O R T I



ISPRA

Istituto Superiore per la Protezione
e la Ricerca Ambientale



Sistema Nazionale
per la Protezione
dell'Ambiente

Italian Emission Inventory 1990 -2020. Informative Inventory Report 2022

Legal Disclaimer

The Institute for Environmental Protection and Research (ISPRA), together with the 21 Regional Agencies (ARPA) and Provincial Agencies (APPA) for the protection of the environment, as of 14 January 2017 is part of the National Network System for the Protection of the Environment (SNPA), established by the Law June 28, 2016, n.132.

The Institute for Environmental Protection and Research, or persons acting on its behalf, are not responsible for the use that may be made of the information contained in this report.

ISPRA - Istituto Superiore per la Protezione e la Ricerca Ambientale
Via Vitaliano Brancati, 48 – 00144 Roma
www.isprambiente.gov.it

ISPRA, Rapporti 361/2022
ISBN 978-88-448-1105-1

Extracts from this document may be reproduced on the condition that the source is acknowledged

Graphic design:

Cover design: Franco Iozzoli

ISPRA – Communications Area

Cover drawing: Chiara Arcarese

Coordination of the online publication:

Daria Mazzella

ISPRA – Communications Area

April 2022

Authors

Ernesto Taurino, Antonella Bernetti, Antonio Caputo, Marco Cordella, Riccardo De Lauretis, Ilaria D'Elia (ENEA), Eleonora Di Cristofaro, Andrea Gagna, Barbara Gonella, Federica Moricci, Emanuele Peschi, Daniela Romano, Marina Vitullo

Contact: Riccardo De Lauretis
telephone +39 0650072543
e-mail riccardo.delawaretis@isprambiente.it

ISPRA- Institute for Environmental Protection and Research
Environmental Assessment, Control and Sustainability Department
Emissions, Prevention of Atmospheric Impacts and Climate Change Area
Air Emission Inventory Unit
Via V. Brancati, 48 00144 Rome ITALY

Text available on ISPRA website at <http://www.isprambiente.gov.it>

CONTENTS

| | |
|--|-----------|
| 1 INTRODUCTION | 9 |
| 1.1 BACKGROUND INFORMATION ON THE CONVENTION ON LONG-RANGE TRANSBOUNDARY AIR POLLUTION | 9 |
| 1.2 NATIONAL INVENTORY | 10 |
| 1.3 INSTITUTIONAL ARRANGEMENTS | 11 |
| 1.4 INVENTORY PREPARATION PROCESS | 12 |
| 1.5 METHODS AND DATA SOURCES | 13 |
| 1.6 KEY CATEGORIES | 15 |
| 1.7 QA/QC AND VERIFICATION METHODS | 19 |
| 1.8 GENERAL UNCERTAINTY EVALUATION | 31 |
| 1.9 GENERAL ASSESSMENT OF COMPLETENESS | 31 |
| 2 ANALYSIS OF KEY TRENDS BY POLLUTANT | 32 |
| 2.1 MAIN POLLUTANTS..... | 32 |
| 2.1.1 Sulphur dioxide (SO _x)..... | 32 |
| 2.1.2 Nitrogen oxides (NO _x)..... | 34 |
| 2.1.3 Ammonia (NH ₃)..... | 36 |
| 2.1.4 Non methane volatile organic compounds (NMVOC) | 38 |
| 2.1.5 Carbon monoxide (CO)..... | 40 |
| 2.2 PARTICULATE MATTER | 43 |
| 2.2.1 PM ₁₀ | 43 |
| 2.2.2 PM _{2.5} | 44 |
| 2.2.3 Black Carbon (BC) | 46 |
| 2.3 HEAVY METALS (PB, CD, HG) | 48 |
| 2.3.1 Lead (Pb) | 48 |
| 2.3.2 Cadmium (Cd)..... | 49 |
| 2.3.3 Mercury (Hg)..... | 50 |
| 2.4 PERSISTENT ORGANIC POLLUTANTS (POPS)..... | 52 |
| 2.4.1 Polycyclic aromatic hydrocarbons (PAH)..... | 52 |
| 2.4.2 Dioxins | 53 |
| 2.4.3 Hexachlorobenzene (HCB)..... | 54 |
| 2.4.4 Polychlorinated biphenyl (PCB)..... | 56 |
| 3 ENERGY (NFR SECTOR 1) | 57 |
| 3.1 OVERVIEW OF THE SECTOR | 57 |
| 3.2 ENERGY INDUSTRIES | 61 |
| 3.2.1 Methodological issues..... | 61 |
| 3.2.1.1 Public Electricity and Heat Production | 61 |
| 3.2.1.2 Refineries | 62 |
| 3.2.1.3 Manufacture of Solid Fuels and Other Energy Industries | 64 |
| 3.2.2 Time series and key categories | 64 |
| 3.2.2.1 Public Electricity and Heat production..... | 65 |
| 3.2.2.2 Refineries | 65 |
| 3.2.2.3 Manufacture of Solid Fuels and Other Energy Industries | 65 |
| 3.2.3 QA/QC and verification | 66 |
| 3.2.4 Recalculations..... | 69 |
| 3.2.5 Planned improvements..... | 69 |
| 3.3 MANUFACTURING INDUSTRIES AND CONSTRUCTION | 70 |
| 3.3.1 Methodological issues..... | 70 |
| 3.3.1.1 Iron and steel | 71 |
| 3.3.1.2 Non-ferrous metals..... | 71 |
| 3.3.1.3 Chemicals..... | 73 |
| 3.3.1.4 Pulp, paper and print | 73 |
| 3.3.1.5 Food processing, beverages and tobacco | 73 |
| 3.3.1.6 Non-metallic minerals..... | 74 |

| | | |
|----------|---|-----|
| 3.3.1.7 | Other | 75 |
| 3.3.2 | Time series and key categories | 75 |
| 3.3.2.1 | Iron and steel | 75 |
| 3.3.2.2 | Non-ferrous metals..... | 75 |
| 3.3.2.3 | Chemicals..... | 76 |
| 3.3.2.4 | Pulp, paper and print | 76 |
| 3.3.2.5 | Food processing, beverages and tobacco | 76 |
| 3.3.2.6 | Non-metallic minerals..... | 76 |
| 3.3.2.7 | Other | 76 |
| 3.3.3 | QA/QC and verification | 76 |
| 3.3.4 | Recalculations..... | 77 |
| 3.3.5 | Planned improvements..... | 77 |
| 3.4 | AVIATION (NFR SUBSECTOR 1.A.3.A) | 78 |
| 3.4.1 | Overview | 78 |
| 3.4.2 | Methodological issues..... | 78 |
| 3.4.3 | Time series and key categories | 79 |
| 3.4.4 | QA/QC and Uncertainty | 81 |
| 3.4.5 | Recalculations..... | 81 |
| 3.4.6 | Planned improvements..... | 81 |
| 3.5 | ROAD TRANSPORT (NFR SUBSECTOR 1.A.3.B)..... | 82 |
| 3.5.1 | Overview | 82 |
| 3.5.2 | Methodological issues..... | 82 |
| 3.5.2.1 | Exhaust emissions | 83 |
| 3.5.2.2 | Evaporative emissions..... | 84 |
| 3.5.2.3 | Emissions from automobile tyre and brake wear | 84 |
| 3.5.2.4 | Emissions from automobile road abrasion | 85 |
| 3.5.3 | Activity data | 85 |
| 3.5.4 | Time series and key categories | 92 |
| 3.5.5 | QA/QC and Uncertainty | 95 |
| 3.5.6 | Recalculation | 96 |
| 3.5.7 | Planned improvements..... | 97 |
| 3.6 | RAILWAYS (NFR SUBSECTOR 1.A.3.C) | 97 |
| 3.7 | NAVIGATION (NFR SUBSECTOR 1.A.3.D)..... | 99 |
| 3.7.1 | Overview | 99 |
| 3.7.2 | Methodological issues..... | 99 |
| 3.7.3 | Time series and key categories | 100 |
| 3.7.4 | QA/QC and Uncertainty | 101 |
| 3.7.5 | Recalculations..... | 101 |
| 3.7.6 | Planned improvements..... | 101 |
| 3.8 | PIPELINE COMPRESSORS (NFR SUBSECTOR 1.A.3.E)..... | 102 |
| 3.9 | CIVIL SECTOR: SMALL COMBUSTION AND OFF-ROAD VEHICLES (NFR SUBSECTOR 1.A.4 - 1.A.5)..... | 103 |
| 3.9.1 | Overview | 103 |
| 3.9.2 | Activity data | 103 |
| 3.9.3 | Methodological issues..... | 104 |
| 3.9.3.1 | NO _x emissions from gas powered plants in the civil sector | 106 |
| 3.9.3.2 | Emissions from wood combustion in the civil sector..... | 106 |
| 3.9.4 | Time series and key categories | 108 |
| 3.9.5 | QA/QC and Uncertainty | 109 |
| 3.9.6 | Recalculation | 109 |
| 3.9.7 | Planned improvements..... | 110 |
| 3.10 | FUGITIVE EMISSIONS (NFR SUBSECTOR 1.B) | 111 |
| 3.10.1 | Overview | 111 |
| 3.10.2 | Methodological issues..... | 111 |
| 3.10.2.1 | Fugitive emissions from natural gas distribution (1.B.2b)..... | 112 |
| 3.10.3 | Time series and key categories | 114 |
| 3.10.4 | QA/QC and Uncertainty | 114 |
| 3.10.5 | Recalculation | 114 |

| | | |
|----------|--|------------|
| 3.10.6 | Planned improvements..... | 114 |
| 4 | IPPU - INDUSTRIAL PROCESSES (NFR SECTOR 2) | 115 |
| 4.1 | OVERVIEW OF THE SECTOR..... | 115 |
| 4.2 | METHODOLOGICAL ISSUES..... | 115 |
| 4.2.1 | Mineral products (2A)..... | 116 |
| 4.2.2 | Chemical industry (2B)..... | 117 |
| 4.2.3 | Metal production (2C)..... | 118 |
| 4.2.4 | Other production (2G – 2H – 2I – 2L)..... | 121 |
| 4.3 | TIME SERIES AND KEY CATEGORIES..... | 122 |
| 4.3.1 | Mineral products (2A)..... | 122 |
| 4.3.2 | Chemical industry (2B)..... | 124 |
| 4.3.3 | Metal production (2C)..... | 124 |
| 4.3.4 | Other production (2G – 2H – 2I – 2L)..... | 126 |
| 4.4 | QA/QC AND VERIFICATION..... | 126 |
| 4.5 | RECALCULATIONS..... | 127 |
| 4.5.1 | Mineral industry (2A)..... | 127 |
| 4.5.2 | Chemical industry (2B)..... | 128 |
| 4.5.3 | Metal industry (2C)..... | 128 |
| 4.5.4 | Other product use (fireworks and tobacco) (2G)..... | 128 |
| 4.5.5 | Other industrial processes (2H)..... | 128 |
| 4.6 | PLANNED IMPROVEMENTS..... | 128 |
| 5 | IPPU - SOLVENT AND OTHER PRODUCT USE (NFR SECTOR 2) | 129 |
| 5.1 | OVERVIEW OF THE SECTOR..... | 129 |
| 5.2 | METHODOLOGICAL ISSUES..... | 130 |
| 5.2.1 | Domestic solvent use (2D3a)..... | 130 |
| 5.2.2 | Asphalt Roofing (2D3c)..... | 131 |
| 5.2.3 | Decorative coating (2D3d1)..... | 132 |
| 5.2.4 | Industrial coating (2D3d2)..... | 132 |
| 5.2.5 | Degreasing (2D3e)..... | 133 |
| 5.2.6 | Dry cleaning (2D3f)..... | 134 |
| 5.2.7 | Chemical products, manufacture and processing (2D3g)..... | 134 |
| 5.2.8 | Other product use (2D3i)..... | 135 |
| 5.3 | TIME SERIES AND KEY CATEGORIES..... | 135 |
| 5.4 | QA/QC AND VERIFICATION..... | 140 |
| 5.5 | RECALCULATIONS..... | 141 |
| 5.6 | PLANNED IMPROVEMENTS..... | 142 |
| 6 | AGRICULTURE (NFR SECTOR 3) | 143 |
| 6.1 | OVERVIEW OF THE SECTOR..... | 143 |
| 6.2 | METHODOLOGICAL ISSUES..... | 145 |
| 6.2.1 | Manure management (3B)..... | 145 |
| 6.2.1.1 | Dairy cattle (3B1a)..... | 146 |
| 6.2.1.2 | Swine (3B3)..... | 149 |
| 6.2.1.3 | Poultry (3B4g)..... | 150 |
| 6.3 | AGRICULTURAL SOILS (3D)..... | 154 |
| 6.4 | FIELD BURNING OF AGRICULTURAL RESIDUES (3F)..... | 157 |
| 6.5 | TIME SERIES AND KEY CATEGORIES..... | 158 |
| 6.6 | QA/QC AND VERIFICATION..... | 161 |
| 6.7 | RECALCULATIONS..... | 162 |
| 6.8 | PLANNED IMPROVEMENTS..... | 163 |
| 7 | WASTE (NFR SECTOR 5) | 164 |
| 7.1 | OVERVIEW OF THE SECTOR..... | 164 |
| 7.2 | METHODOLOGICAL ISSUES..... | 165 |
| 7.2.1 | Solid waste disposal on land (5A)..... | 165 |

| | | |
|-------------------|--|------------|
| 7.2.2 | <i>Biological treatment of waste (5B)</i> | 167 |
| 7.2.3 | <i>Waste Incineration (5C1a – 5C1b)</i> | 167 |
| 7.2.4 | <i>Cremation of corpses (5C1bv)</i> | 169 |
| 7.2.5 | <i>Small scale waste burning (5C2)</i> | 170 |
| 7.2.6 | <i>Wastewater treatments (5D)</i> | 172 |
| 7.2.7 | <i>Other waste (5E)</i> | 172 |
| 7.3 | TIME SERIES AND KEY CATEGORIES..... | 173 |
| 7.4 | RECALCULATIONS..... | 178 |
| 7.5 | PLANNED IMPROVEMENTS..... | 179 |
| 8 | RECALCULATIONS AND IMPROVEMENTS | 180 |
| 8.1 | RECALCULATIONS..... | 180 |
| 8.2 | PLANNED IMPROVEMENTS..... | 182 |
| 9 | PROJECTIONS | 183 |
| 9.1 | THE NATIONAL FRAMEWORK..... | 183 |
| 9.2 | INPUT SCENARIOS..... | 184 |
| 9.2.1 | <i>The energy scenario</i> | 184 |
| 9.2.2 | <i>The scenario of non-energy activities</i> | 184 |
| 9.2.3 | <i>The control strategy definition</i> | 186 |
| 9.3 | THE HARMONIZATION PROCESS..... | 188 |
| 9.4 | THE EMISSION SCENARIO..... | 191 |
| 9.5 | THE NEC EMISSION TARGETS..... | 196 |
| 10 | REPORTING OF GRIDDED EMISSIONS AND LPS | 197 |
| 10.1 | FUGITIVE..... | 198 |
| 10.2 | TRANSPORT..... | 198 |
| 10.3 | AGRICULTURE..... | 200 |
| 10.4 | LPS DATA..... | 201 |
| 11 | REFERENCES | 202 |
| 11.1 | INTRODUCTION..... | 202 |
| 11.2 | ANALYSIS OF KEY TRENDS BY POLLUTANT..... | 203 |
| 11.3 | ENERGY (NRF SECTOR 1)..... | 203 |
| 11.4 | IPPU - INDUSTRIAL PROCESSES (NRF SECTOR 2)..... | 206 |
| 11.5 | IPPU - SOLVENT AND OTHER PRODUCT USE (NRF SECTOR 2)..... | 207 |
| 11.6 | AGRICULTURE (NRF SECTOR 3)..... | 208 |
| 11.7 | WASTE (NRF SECTOR 5)..... | 210 |
| 11.8 | RECALCULATIONS AND IMPROVEMENTS..... | 212 |
| 11.9 | PROJECTIONS..... | 213 |
| 11.10 | REPORTING OF GRIDDED EMISSIONS AND LPS..... | 213 |
| APPENDIX 1 | SUMMARY INFORMATION ON CONDENSABLE IN PM | 214 |

EXECUTIVE SUMMARY

The *Italian Informative Inventory Report (IIR)* is edited in the framework of the *United Nations Economic Commission for Europe (UNECE) Convention on Long Range Transboundary Air Pollution (CLRTAP)*. It contains information on the Italian inventory up to the year 2020, including an explanation of methodologies, data sources, QA/QC activities and verification processes carried out during the inventory compilation, with an analysis of emission trends and a description of key categories.

The aim of the document is to facilitate understanding of the calculation of the Italian air pollutant emission data, hence providing a common mean for comparing the relative contribution of different emission sources and supporting the identification of reduction policies.

The Institute for Environmental Protection and Research (ISPRA) has the overall responsibility for the emission inventory submission to CLRTAP, as well as to the *United Nations Framework Convention on Climate Change (UNFCCC)*, and is in charge of all the work related to inventory compilation.

In particular, in compliance with the LRTAP Convention, Italy has to submit annually data on national emissions of SO_x, NO_x, NMVOC, CO and NH₃, and various heavy metals and POPs. The submission consists of the national emission inventory, communicated through compilation of the Nomenclature Reporting Format (NRF), and the informative inventory report (IIR) to ensure the properties of transparency, consistency, comparability, completeness and accuracy.

In the period 1990-2020, emissions from almost all the pollutants described in this report show a downward trend. Reductions are especially relevant for the main pollutants (SO_x -95%; NO_x -73%; CO -72%; NMVOC -56%), for BC (-67%), cadmium (-65%), mercury (-62%), lead (-96%) and hexachlorobenzene (-91%). The major drivers for the trend are reductions in the industrial and road transport sectors, due to the implementation of various European Directives which introduced new technologies, plant emission limits, the limitation of sulphur content in liquid fuels and the shift to cleaner fuels. Emissions have also decreased for the improvement of energy efficiency as well as the promotion of renewable energy.

The energy sector is the main source of emissions in Italy with a share of more than 80%, including fugitive emissions, for many pollutants (SO_x 93%; NO_x 89%; CO 94%; PM_{2.5} 86%; BC 93%; PAH 86%). The industrial processes sector is an important source of emissions specifically related to the iron and steel production, at least for particulate matter, heavy metals and POPs, whereas significant emissions of SO_x derive from carbon black and sulphuric acid production; on the other hand, the solvent and other product use sector is characterized by NMVOC emissions. The agriculture sector is the main source of NH₃ emissions in Italy with a share of 95% in national total. Finally, the waste sector, specifically waste incineration, is a relevant source for Cd (9%) and dioxins (17%).

Emission figures of the Italian emission inventory and other related documents are publicly available at <http://emissioni.sina.isprambiente.it/inventario-nazionale/>.

1 INTRODUCTION

1.1 BACKGROUND INFORMATION ON THE CONVENTION ON LONG-RANGE TRANSBOUNDARY AIR POLLUTION

The 1979 Geneva *Convention on Long-range Transboundary Air Pollution*, contributing to the development of international environmental law, is one of the fundamental international means for the protection of the human health and the environment through the intergovernmental cooperation.

The fact that air pollutants could travel several thousands of kilometres before deposition and damage occurred outlined the need for international cooperation.

In November 1979, in Geneva, 34 Governments and the European Community (EC) signed the Convention. The *Convention on Long-range Transboundary Air Pollution* was ratified by Italy in the year 1982 and entered into force in 1983. It has been extended by the following eight specific protocols:

- The 1984 Protocol on Long-term Financing of the Cooperative Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe (EMEP); 42 Parties. Entered into force on 28th January 1988.
- The 1985 Protocol on the Reduction of Sulphur Emissions or their Transboundary Fluxes by at least 30 per cent; 23 Parties. Entered into force on 2nd September 1987.
- The 1988 Protocol concerning the Control of Nitrogen Oxides or their Transboundary Fluxes; 31 Parties. Entered into force on 14th February 1991.
- The 1991 Protocol concerning the Control of Emissions of Volatile Organic Compounds or their Transboundary Fluxes; 22 Parties. Entered into force on 29th September 1997.
- The 1994 Protocol on Further Reduction of Sulphur Emissions; 27 Parties. Entered into force on 5th August 1998.
- The 1998 Protocol on Heavy Metals; 28 Parties. Entered into force on 29 December 2003.
- The 1998 Protocol on Persistent Organic Pollutants (POPs); 28 Parties. Entered into force on 23rd October 2003.
- The 1999 Protocol to Abate Acidification, Eutrophication and Ground-level Ozone; 23 Parties. Entered into force on 17th May 2005. (Guidance documents to Protocol adopted by decision 1999/1).
- The following table shows the dates of signature and ratification of Convention and Protocols for Italy.

Table 1.1 Dates of signature and ratification of the UNECE Convention and Protocols

| | SIGNATURE | RATIFICATION |
|---|------------|--------------|
| 1979 Convention | 14/11/1979 | 15/07/1982 |
| 1984 EMEP Protocol | 28/09/1984 | 12/01/1989 |
| 1985 Sulphur Protocol | 09/07/1985 | 05/02/1990 |
| 1988 NO _x Protocol | 01/11/1988 | 19/05/1992 |
| 1991 VOC Protocol | 19/11/1991 | 30/06/1995 |
| 1994 Sulphur Protocol | 14/06/1994 | 14/09/1998 |
| 1998 Heavy Metals Protocol | 24/06/1998 | |
| 1998 POPs Protocol | 24/06/1998 | 20/06/2006 |
| 1999 Multi-effect Protocol (reviewed in 2012) | 01/12/1999 | |

The following classes of pollutants should be included in the emission inventory:

Main Pollutants

- Sulphur oxides (SO_x), in mass of SO₂;
- Nitrous oxides (NO_x), in mass of NO₂;
- Non-methane volatile organic compounds (NMVOC);
- Ammonia (NH₃);
- Carbon monoxide (CO).

Particulate matter

- TSP, total suspended particulate;
- PM10, particulate matter less than 10 microns in diameter;
- PM2.5, particulate matter less than 2.5 microns in diameter;
- Black carbon.

Heavy Metals

- Priority Metals: Lead (Pb), Cadmium (Cd) and Mercury (Hg);
- Other metals: Arsenic (As), Chrome (Cr), Copper (Cu), Nickel (Ni), Selenium (Se) and Zinc (Zn).

Persistent organic pollutants (POPs)

- As specified in Annex II of the POPs Protocol, including Polychlorinated Biphenyls (PCBs);
- As specified in Annex III of the POPs Protocol: Dioxins (Diox), Polycyclic Aromatic Hydrocarbons (PAHs), Hexachlorobenzene (HCB).

1.2 NATIONAL INVENTORY

As a Party to the *United Nations Economic Commission for Europe (UNECE) Convention on Long Range Transboundary Air Pollution (CLRTAP)*, Italy has to submit annually data on emissions of air pollutants in order to fulfil obligations, in compliance with the implementation of Protocols under the Convention. Parties are required to report on annual national emissions of SO_x, NO_x, NMVOC, CO and NH₃, and various heavy metals and POPs according to the *Guidelines for Reporting Emission Data under the Convention on Long-range Transboundary Air Pollution (UNECE, 2008)*. The same data are submitted also in the framework of the National Emission Ceiling Directive of the European Union (EU, 2016).

Specifically, the submission consists of the national LRTAP emission inventory, communicated through compilation of the *Nomenclature Reporting Format (NRF)*, and the *Informative Inventory Report (IIR)*.

The Italian informative inventory report contains information on the national inventory for the year 2020, including descriptions of methods, data sources, QA/QC activities carried out and a trend analysis. The inventory accounts for anthropogenic emissions of the following substances: sulphur oxides (SO_x), nitrogen oxides (NO_x), ammonia (NH₃), non-methane volatile organic compounds (NMVOC), carbon monoxide (CO), total suspended particulate (TSP), particulate matter, particles of size <10 µm, (PM10), particulate matter, particles of size < 2.5µm, (PM2.5), black carbon (BC), lead (Pb), cadmium (Cd), mercury (Hg), arsenic (As), chromium (Cr), copper (Cu), nickel (Ni), selenium (Se), zinc (Zn), polychlorinated biphenyls (PCBs), polycyclic aromatic hydrocarbons (PAH), dioxins (Diox), hexachlorobenzene (HCB). Other pollutants are reported as not estimated; more in details polycyclic aromatic hydrocarbons have not been estimates for each compound for all the sectors and further investigation is planned for the reporting of these emissions.

Detailed information on emission figures of primary pollutants, particulate matter, heavy metals and persistent organic pollutants as well as estimation procedures are provided in order to improve the transparency, consistency, comparability, accuracy and completeness of the inventory provided.

The national inventory is updated annually in order to reflect revisions and improvements in the methodology and the availability of new information. Changes are applied retrospectively to earlier years, which accounts for any difference in previously published data.

Total emissions by pollutant from 1990 to 2020 are reported in Table 1.2.

Table 1.2 Emission time series by pollutant

| | | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|-----------------|-----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| SO _x | <i>Gg</i> | 1,784 | 1,322 | 756 | 411 | 222 | 126 | 120 | 117 | 109 | 105 | 82 |
| NO _x | <i>Gg</i> | 2,124 | 1,989 | 1,504 | 1,289 | 935 | 716 | 701 | 658 | 659 | 639 | 571 |
| NMVOC | <i>Gg</i> | 1,993 | 2,058 | 1,630 | 1,340 | 1,116 | 897 | 881 | 921 | 894 | 888 | 885 |
| NH ₃ | <i>Gg</i> | 469 | 454 | 457 | 421 | 379 | 357 | 370 | 364 | 351 | 349 | 363 |
| CO | <i>Gg</i> | 6,797 | 7,072 | 4,751 | 3,467 | 3,073 | 2,267 | 2,192 | 2,259 | 2,050 | 2,061 | 1,873 |
| As | <i>Mg</i> | 37 | 27 | 39 | 28 | 17 | 9 | 8 | 7 | 7 | 6 | 5 |
| Cd | <i>Mg</i> | 11 | 10 | 10 | 8 | 5 | 4 | 4 | 4 | 4 | 4 | 4 |
| Cr | <i>Mg</i> | 86 | 69 | 44 | 50 | 40 | 35 | 35 | 35 | 35 | 34 | 30 |

| | | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--------|--------|-------|-------|------|------|------|------|------|------|------|------|------|
| Cu | Mg | 193 | 216 | 221 | 230 | 203 | 189 | 170 | 162 | 165 | 164 | 136 |
| Hg | Mg | 15 | 14 | 15 | 13 | 9 | 7 | 6 | 7 | 7 | 6 | 6 |
| Ni | Mg | 114 | 110 | 107 | 112 | 41 | 30 | 30 | 30 | 29 | 28 | 26 |
| Pb | Mg | 4,278 | 1,993 | 961 | 295 | 216 | 197 | 174 | 181 | 183 | 176 | 156 |
| Se | Mg | 8 | 8 | 8 | 9 | 8 | 8 | 7 | 7 | 7 | 6 | 6 |
| Zn | Mg | 913 | 896 | 853 | 920 | 823 | 765 | 712 | 764 | 784 | 751 | 668 |
| TSP | Gg | 365 | 361 | 317 | 289 | 295 | 242 | 236 | 244 | 224 | 221 | 207 |
| PM10 | Gg | 302 | 297 | 257 | 232 | 240 | 195 | 190 | 196 | 178 | 176 | 166 |
| PM2.5 | Gg | 230 | 228 | 197 | 176 | 199 | 160 | 155 | 162 | 144 | 138 | 133 |
| BC | Gg | 48 | 47 | 43 | 39 | 32 | 23 | 21 | 21 | 19 | 18 | 16 |
| PAH | Mg | 90 | 92 | 60 | 64 | 87 | 70 | 70 | 74 | 67 | 65 | 60 |
| Dioxin | g ITeq | 529 | 511 | 434 | 361 | 342 | 310 | 312 | 330 | 311 | 307 | 280 |
| HCB | kg | 142 | 110 | 33 | 27 | 16 | 16 | 15 | 16 | 15 | 14 | 13 |
| PCB | kg | 154 | 166 | 157 | 179 | 133 | 114 | 120 | 122 | 121 | 116 | 103 |

The NRF files and other related documents can be found on website at the following address:

<http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/> .

1.3 INSTITUTIONAL ARRANGEMENTS

The Institute for Environmental Protection and Research (ISPRA) has the overall responsibility for the compilation of the national emission inventory and submissions to CLRTAP. The Institute is also responsible for the communication of pollutants under the NEC directive as well as, jointly with the Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), the development of emission scenarios, as established by the Legislative Decree n. 171 of 21st May 2004. Every four years, from 2017 with reference to 2015 emissions, ISPRA shall provide the disaggregation of the national inventory at provincial level as instituted by the Legislative Decree n. 81 of 30 May 2018. Moreover, ISPRA is the single entity in charge of the development and compilation of the national greenhouse gas emission inventory as indicated by the Legislative Decree n. 51 of 7th March 2008. The Ministry for the Environment, Land and Sea is responsible for the endorsement and for the communication of the inventory to the Secretariat of the different conventions.

The *Italian National System* currently in place is fully described in the document ‘*National Greenhouse Gas Inventory System in Italy*’ (ISPRA, 2018).

A specific unit of the Institute is responsible for the compilation of the *Italian Atmospheric Emission Inventory* and the *Italian Greenhouse Gas Inventory* in the framework of both the *Convention on Climate Change* and the *Convention on Long Range Transboundary Air Pollution*. The whole inventory is compiled by the Institute; scientific and technical institutions and consultants may help in improving information both on activity data and emission factors of specific activities. All the measures to guarantee and improve the transparency, consistency, comparability, accuracy and completeness of the inventory are undertaken.

ISPRA bears the responsibility for the general administration of the inventory, co-ordinates participation in review processes, publishes and archives the inventory results.

Specifically, ISPRA is responsible for all aspects of national inventory preparation, reporting and quality management. Activities include the collection and processing of data from different data sources, the selection of appropriate emissions factors and estimation methods consistent with the EMEP/EEA guidebook, the *IPCC 1996 Revised Guidelines*, the *IPCC Good Practice Guidance and Uncertainty management* and the *IPCC Good Practice Guidance for land use, land-use change and forestry*, and the *IPCC 2006 Guidelines*, the compilation of the inventory following the QA/QC procedures, the preparation of the *Informative Inventory Report* and the reporting through the *Nomenclature Reporting Format*, the response to review checks, the updating and data storage.

Different institutions are responsible for statistical basic data and data publication, which are primary to ISPRA for carrying out estimates. These institutions are part of the *National Statistical System* (Sistan), which provides national official statistics, and therefore are asked periodically to update statistics; moreover, the *National Statistical System* ensures the homogeneity of the methods used for official statistics data through a coordination plan, involving the entire public administration at central, regional and local levels.

The main Sistan products, which are primarily necessary for the inventory compilation, are:

- National Statistical Yearbooks, Monthly Statistical Bulletins, by ISTAT (National Institute of Statistics);
- Annual Report on the Energy and Environment, by ENEA (Agency for New Technologies, Energy and the Environment);
- National Energy Balance (annual), Petrochemical Bulletin (quarterly publication), by MSE (Ministry of Economic Development);
- Transport Statistics Yearbooks, by MIT (Ministry of Transportation);
- Annual Statistics on Electrical Energy in Italy, by TERNA (National Independent System Operator);
- Annual Report on Waste, by ISPRA;
- National Forestry Inventory, by MIPAAF (Ministry of Agriculture, Food and Forest Policies).

The national emission inventory itself is a Sistan product (ISPRA).

Other information and data sources are used to carry out emission estimates, which are generally referred to in Table 1.3 in the following section 1.5.

1.4 INVENTORY PREPARATION PROCESS

ISPRA has established fruitful cooperation with several governmental and research institutions as well as industrial associations, which helps improving information about some leading categories of the inventory. Specifically, these activities aim at the improvement of provision and collection of basic data and emission factors, through plant-specific data, and exchange of information on scientific researches and new sources. Moreover, when in depth investigation is needed and estimates are affected by a high uncertainty, sectoral studies are committed to *ad hoc* research teams or consultants.

ISPRA also coordinates with different national and regional authorities and private institutions for the cross-checking of parameters and estimates, as well as with *ad hoc* expert panels, in order to improve the completeness and transparency of the inventory.

The main basic data needed for the preparation of the national emission inventory are energy statistics, published by the Ministry of Economic Development (MSE) in the National Energy Balance (BEN), statistics on industrial and agricultural production, published by the National Institute of Statistics (ISTAT), statistics on transportation, provided by the Ministry of Transportation (MIT), and data supplied directly by the relevant professional associations.

Emission factors and methodologies used in the estimation process are consistent with the EMEP/EEA Guidebook, the IPCC Guidelines and Good Practice Guidance as well as supported by national experiences and circumstances.

For the industrial sector, emission data collected through the national Pollutant Release and Transfer Register (Italian PRTR), the Large Combustion Plant (LCP) Directive and in the framework of the European Emissions Trading Scheme have yielded considerable developments in the inventory of the relevant sectors. In fact, these data, even if not always directly used, are considered as a verification of emission estimates and improve national emissions factors as well as activity data figures.

In addition, final estimates are checked and verified also in view of annual environmental reports by industries.

For large industrial point sources, emissions are registered individually, when communicated, based upon detailed information such as fuel consumption. Other small plants communicate their emissions which are also considered individually.

Emission estimates are drawn up for each sector. Final data are communicated to the UNECE Secretariat filling in the NRF files.

The process of the inventory preparation is carried out annually. In addition to a new year, the entire time series is checked and revised during the annual compilation of the inventory. Recalculations are elaborated on account of changes in the methodologies used to carry out emission estimates, changes due to different allocation of emissions as compared to previous submissions and changes due to error corrections. The inventory may also be expanded by including categories not previously estimated if enough information on activity data and suitable emission factors have been identified and collected. Information on the major recalculations is provided in the sectoral chapter of the report.

All the reference material, estimates and calculation sheets, as well as the documentation on scientific papers and the basic data needed for the inventory compilation, are stored and archived at the Institute. After each reporting cycle, all database files, spreadsheets and electronic documents are archived as 'read-only-files'

so that the documentation and estimates could be traced back during the new year inventory compilation or a review process.

Technical reports and emission figures are publicly accessible on the web at the address <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>.

1.5 METHODS AND DATA SOURCES

An outline of methodologies and data sources used in the preparation of the emission inventory for each sector is provided in the following. In Table 1.3 a summary of the activity data and sources used in the inventory compilation is reported.

Table 1.3 *Main activity data and sources for the Italian Emission Inventory*

| SECTOR | ACTIVITY DATA | SOURCE |
|---|--|---|
| 1 Energy | | |
| 1A1 Energy Industries | Fuel use | Energy Balance - Ministry of Economic Development Major national electricity producers European Emissions Trading Scheme |
| 1A2 Manufacturing Industries and Construction | Fuel use | Energy Balance - Ministry of Economic Development Major National Industry Corporation European Emissions Trading Scheme |
| 1A3 Transport | Fuel use Number of vehicles Aircraft landing and take-off cycles and maritime activities | Energy Balance - Ministry of Economic Development Statistical Yearbooks - National Statistical System Statistical Yearbooks - Ministry of Transportation Statistical Yearbooks - Italian Civil Aviation Authority (ENAC) Maritime and Airport local authorities |
| 1A4 Residential-public-commercial sector | Fuel use | Energy Balance - Ministry of Economic Development |
| 1B Fugitive Emissions from Fuel | Amount of fuel treated, stored, distributed | Energy Balance - Ministry of Economic Development Statistical Yearbooks - Ministry of Transportation Major National Industry Corporation |
| 2 Industrial Processes | Production data | National Statistical Yearbooks- National Institute of Statistics International Statistical Yearbooks-UN European Emissions Trading Scheme European Pollutant Release and Transfer Register Sectoral Industrial Associations |
| 2D Solvent and Other Product Use | Amount of solvent use | National Environmental Publications - Sectoral Industrial Associations International Statistical Yearbooks - UN |
| 3 Agriculture | Agricultural surfaces Production data Number of animals Fertilizer consumption | Agriculture Statistical Yearbooks - National Institute of Statistics Sectoral Agriculture Associations |
| 4 Land Use, Land Use Change and Forestry | Forest and soil surfaces Amount of biomass Biomass burnt Biomass growth | Statistical Yearbooks - National Institute of Statistics State Forestry Corps National and Regional Forestry Inventory Universities and Research Institutes |
| 5 Waste | Amount of waste | National Waste Cadastre - Institute for Environmental Protection and Research |

Methodologies are consistent with the *EMEP/EEA Emission Inventory Guidebook, Revised 1996 and 2006 IPCC Guidelines*, and *IPCC Good Practice Guidance* (EMEP/CORINAIR, 2007; EMEP/EEA, 2009; EMEP/EEA, 2013; EMEP/EEA, 2016; EMEP/EEA, 2019; IPCC, 1997; IPCC, 2000; IPCC, 2006); national

emission factors are used as well as default emission factors from international guidebooks, when national data are not available. The development of national methodologies is supported by background documents.

The most complete document describing national methodologies used in the emission inventory compilation is the *National Inventory Report*, submitted in the framework of the *UN Convention on Climate Change* and the *Kyoto Protocol* (ISPRA, 2022 [a]).

Activity data used in emission calculations and their sources are briefly described here below.

In general, for the energy sector, basic statistics for estimating emissions are fuel consumption published in the national Energy Balance by the Ministry of Economic Development. Additional information for electricity production is provided by the major national electricity producers and by the major national industry corporation. On the other hand, basic information for road transport, maritime and aviation, such as the number of vehicles, harbour statistics and aircraft landing and take-off cycles are provided in statistical yearbooks published both by the National Institute of Statistics and the Ministry of Transportation. Other data are communicated by different category associations.

Data from the Italian Emissions Trading Scheme database (ETS) are incorporated into the national inventory whenever the sectoral coverage is complete; in fact, these figures do not always entirely cover the energy categories whereas national statistics, such as the national energy balance and the energy production and consumption statistics, provide the complete basic data needed for the Italian emission inventory. However, the analysis of data from ETS is used to develop country-specific emission factors and check activity data levels. In this context, ISPRA is also responsible for developing, operating and maintaining the national registry under Directive 2003/87/CE as instituted by the Legislative Decree 51 of March 7th 2008; the Institute performs this tasks under the supervision of the national Competent Authority for the implementation of directive 2003/87/CE, amended by Directive 2009/29/EC, jointly established by the Ministry for Environment, Land and Sea and the Ministry for Economic Development.

For the industrial sector, the annual production data are provided by national and international statistical yearbooks. Emission data collected through the national Pollutant Release and Transfer Register (Italian PRTR) are also used in the development of emission estimates or considered as a verification of emission estimates for some specific categories. Italian PRTR data are reported by operators to national and local competent authorities for quality assessment and validation. ISPRA collects facilities' reports and supports the validation activities at national and at local level. ISPRA communicates to the Ministry for the Environment, Land and Sea and to the European Commission within 31st March of the current year for data referring to two years earlier. These data are used for the compilation of the inventory whenever they are complete in terms of sectoral information; in fact, industries communicate figures only if they exceed specific releases thresholds; furthermore, basic data such as fuel consumption are not required and production data are not split by product but reported as an overall value. Anyway, the national PRTR is a good basis for data checks and a way to facilitate contacts with industries which supply, under request, additional information as necessary for carrying out sectoral emission estimates.

In addition, final emissions are checked and verified also considering figures reported by industries in their annual environmental reports.

Both for energy and industrial processes, emissions of large industrial point sources are registered individually; communication also takes place in the framework of the European Directive on Large Combustion Plants, based upon detailed information such as fuel consumption. Other small plants communicate their emissions which are also considered individually.

For the other sectors, i.e. for solvents, the amount of solvent use is provided by environmental publications of sector industries and specific associations as well as international statistics.

For agriculture, annual production data and number of animals are provided by the National Institute of Statistics and other sectoral associations.

For waste, the main activity data are provided by the Institute for Environmental Protection and Research.

In case basic data are not available proxy variables are considered; unpublished data are used only if supported by personal communication and confidentiality of data is respected.

All the material and documents used for the inventory emission estimates are stored at the Institute for Environmental Protection and Research. The inventory is composed by spreadsheets to calculate emission estimates; activity data and emission factors as well as methodologies are referenced to their data sources.

A 'reference' database has also been developed to increase the transparency of the inventory; at the moment, it is complete as far as references to greenhouse gas emissions are concerned.

1.6 KEY CATEGORIES

A key category analysis of the Italian inventory is carried out according to the Approach 1 method described in the EMEP/EEA Guidebook (EMEP/EEA, 2019). According to these guidelines, a key category is defined as an emission category that has a significant influence on a country's inventory in terms of the absolute level in emissions. Key categories are those which, when summed together in descending order of magnitude, add up to over 80% of the total emissions.

National emissions have been disaggregated into the categories reported in the National Format Report; details vary according to different pollutants in order to reflect specific national circumstances. Results are reported in the following tables for the year 1990 (Table 1.4) and 2020 (Table 1.5) by pollutant.

The trend analysis has also been applied considering 1990 and 2020. The results are reported in Table 1.6.

Table 1.4 Key categories for the Italian Emission Inventory in 1990

| | Key categories in 1990 | | | | | | | | | | | | | Total (%) |
|-----------------------|------------------------|----------------------------------|---------------------|--------------------|--------------------|--------------------|--------------------|-------------------|-------------------|-------------------|-------------------|----------------|----------------|-----------|
| SO_x | 1A1a (43.1%) | 1A1b (10.8%) | 1A2c (7.2%) | 1A3d ii (4.4%) | 1A4b i (4.1%) | 1B2a iv (3.8%) | 1A2gviii (3.7%) | 1A2f (3.6%) | | | | | | 80.6 |
| NO_x | 1A3b i (27.8%) | 1A1a (19.2%) | 1A3b iii (16.0%) | 1A2f (5.7%) | 1A4c ii (4.8%) | 1A3d ii (4.5%) | 1A3b ii (2.9%) | | | | | | | 80.9 |
| NH₃ | 3Da2a (20.3%) | 3B1a (19.9%) | 3B1b (18.9%) | 3Da1 (14.9%) | 3B3 (7.8%) | | | | | | | | | 81.7 |
| NMVOC | 1A3b i (21.6%) | 2D3d (13.6%) | 1A3b iv (8.8%) | 1A3b v (6.0%) | 2D3a (5.8%) | 1A4b i (5.0%) | 2D3g (3.9%) | 1A4c ii (3.5%) | 2D3i (3.3%) | 1B2a v (3.0%) | 2D3e (2.6%) | 3B1a (2.5%) | 3B1b (2.3%) | 81.9 |
| CO | 1A3b i (60.7%) | 1A4b i (11.6%) | 1A3b iv (7.3%) | 1A4c ii (4.1%) | | | | | | | | | | 83.6 |
| PM10 | 1A4b i (22.2%) | 1A1a (12.5%) | 1A3b i (6.2%) | 1A4c ii (5.3%) | 1A3b iii (4.5%) | 3Dc (4.2%) | 1A3b ii (3.4%) | 1A2f (3.4%) | 1A3d ii (3.1%) | 3B4g ii (2.7%) | 1A3b vi (2.5%) | 2C1 (2.4%) | 2A1 (2.3%) | |
| | 2A5b (2.2%) | 1A2a (2.1%) | 1A2c (1.9%) | | | | | | | | | | | 80.9 |
| PM2.5 | 1A4b i (28.9%) | 1A1a (10.9%) | 1A3b i (8.2%) | 1A4c ii (6.9%) | 1A3b iii (5.9%) | 1A3b ii (4.5%) | 1A3d ii (4.0%) | 1A2f (3.3%) | 2C1 (2.5%) | 1A2a (2.2%) | 1A3bvi (1.8%) | 1A1b (1.7%) | | 80.9 |
| BC | 1A3b i (21.6%) | 1A4c ii (18.8%) | 1A3b iii (14.2%) | 1A3b ii (11.9%) | 1A4b i (10.9%) | 1A2g vii (4.9%) | | | | | | | | 82.2 |
| Pb | 1A3b i (77.8%) | 1A3b iv (5.2%) | | | | | | | | | | | | 83.1 |
| Cd | 1A2b (26.2%) | 1A2a (19.2%) | 2C1 (12.0%) | 1A4b i (9.2%) | 1A2f (5.2%) | 2G (4.5%) | 5C2 (3.9%) | 1A4a i (3.5%) | 2B10a (3.2%) | | | | | 83.0 |
| Hg | 1B2d (22.2%) | 2B10a (18.4%) | 2C1 (15.0%) | 1A2b (10.6%) | 1A2f (8.3%) | 1A1a (6.5%) | | | | | | | | 81.1 |
| PAH | 2C1 (53.8%) | 1A4b i (35.3%) | | | | | | | | | | | | 89.1 |
| Dioxin | 1A4a i (19.6%) | 1A2a (15.4%) | 1A4b i (13.2%) | 2C1 (12.7%) | 5C1a (8.1%) | 5E (5.9%) | 5C1b i (5.9%) | | | | | | | 80.7 |
| HCB | 3Df (83.4%) | | | | | | | | | | | | | 83.4 |
| PCB | 2C1 (59.4%) | 1A2a (25.6%) | | | | | | | | | | | | 85.1 |
| | 1 Energy | 2 IPPU - Solvent and product use | 5 Waste | | | | | | | | | | | |
| | 2 IPPU - Industry | 3 Agriculture | | | | | | | | | | | | |

Table 1.5 Key categories for the Italian Emission Inventory in 2020

| | Key categories in 2020 | | | | | | | | | | | | | Total (%) |
|-----------------------|------------------------|----------------------------------|---------------------|---------------------|--------------------|-------------------|-------------------|-------------------|-----------------|------------------|-------------------|----------------|------------------|-----------|
| SO_x | 1A2f (22.9%) | 1B2a iv (13.7%) | 1A3d ii (9.8%) | 1A2g viii (6.7%) | 1A2a (6.7%) | 1A4b i (6.6%) | 1A1a (6.1%) | 1A4a i (5.3%) | 2B10a (5.0%) | | | | | 82.8 |
| NO_x | 1A3b i (19.1%) | 1A3d ii (15.0%) | 1A3b iii (11.7%) | 1A4b i (6.8%) | 1A3b ii (6.2%) | 1A4a i (5.5%) | 1A2f (5.0%) | 1A4c ii (4.3%) | 1A1a (4.2%) | 3Da1 (4.0%) | | | | 81.7 |
| NH₃ | 3Da2a (18.4%) | 3B1b (18.2%) | 3Da1 (16.9%) | 3B1a (13.7%) | 3B3 (8.5%) | 3B4g ii (3.7%) | 3Da2c (2.7%) | | | | | | | 82.1 |
| NMVOC | 2D3d (15.2%) | 1A4b i (15.1%) | 2D3a (14.3%) | 2D3g (5.7%) | 1A3b v (5.2%) | 3B1a (4.2%) | 3B1b (4.2%) | 1A4a i (3.2%) | 2H2 (3.2%) | 2D3i (3.2%) | 1A3b iv (3.0%) | 2D3h (2.0%) | 1A3b i (1.9%) | 80.4 |
| CO | 1A4b i (62.3%) | 1A3b i (9.5%) | 1A3b iv (4.7%) | 1A2a (3.1%) | 1A3d ii (2.8%) | | | | | | | | | 82.5 |
| PM10 | 1A4b i (53.1%) | 3Dc (6.2%) | 1A3b vi (4.2%) | 1A3d ii (3.9%) | 1A2f (2.7%) | 5E (2.5%) | 1A3bvii (2.3%) | 2C1 (2.1%) | 2A1 (1.9%) | 1A3b i (1.6%) | | | | 80.3 |
| PM2.5 | 1A4b i (65.2%) | 1A3d ii (4.8%) | 5E (3.1%) | 1A2f (3.0%) | 1A3b vi (2.9%) | 2C1 (2.2%) | | | | | | | | 81.1 |
| BC | 1A4b i (49.7%) | 1A3b i (13.3%) | 1A3d ii (6.9%) | 5C2 (5.6%) | 1A3b iii (4.0%) | 1A4c ii (3.6%) | | | | | | | | 83.1 |
| Pb | 2C1 (37.8%) | 1A2f (36.4%) | 1A4b i (5.5%) | 1A3b vi (5.5%) | | | | | | | | | | 85.3 |
| Cd | 2C1 (22.4%) | 1A2f (14.2%) | 1A2a (11.0%) | 1A4b i (9.5%) | 2G (8.8%) | 5C2 (8.7%) | 1A3b i (5.6%) | | | | | | | 80.3 |
| Hg | 2C1 (44.2%) | 1A2b (9.9%) | 1A2f (9.4%) | 1A2a (6.9%) | 1A1a (6.4%) | 1B2d (5.9%) | | | | | | | | 82.8 |
| PAH | 1A4b i (79.0%) | 2C1 (11.8%) | | | | | | | | | | | | 90.8 |
| Dioxin | 1A4b i (33.2%) | 2C1 (27.0%) | 1A2b (15.3%) | 5E (14.7%) | | | | | | | | | | 90.3 |
| HCB | 1A2b (37.3%) | 3Df (18.5%) | 1A4b i (11.5%) | 1A4a i (11.1%) | 5C1b iii (5.6%) | | | | | | | | | 84.0 |
| PCB | 2C1 (71.2%) | 1A4b i (14.5%) | | | | | | | | | | | | 85.7 |
| 1 Energy | | 2 IPPU - Solvent and product use | | | 5 Waste | | | | | | | | | |
| 2 IPPU - Industry | | 3 Agriculture | | | | | | | | | | | | |

Table 1.6 Key categories for the Italian Emission Inventory in trend 1990-2020

| | Key categories in trend | | | | | | | | | | | | Total (%) | |
|-------------------------|----------------------------------|--------------------|--------------------|-------------------|--------------------|--------------------|-------------------|------------------|------------------|-------------------|----------------|------------------|-----------|------|
| SO_x | 1A1a (31.1%) | 1A2f (16.2%) | 1B2a iv (8.3%) | 1A2c (5.6%) | 1A1b (4.9%) | 1A2a (4.7%) | 1A3d ii (4.6%) | 1A4a i (4.4%) | 1B2c (3.4%) | | | | | 83.2 |
| NO_x | 1A1a (21.3%) | 1A3dii (14.9%) | 1A3b i (12.4%) | 1A4a i (7.0%) | 1A4b i (6.1%) | 1A3b iii (6.1%) | 1A3b ii (4.7%) | 3Da1 (3.7%) | 3Da2a (3.3%) | 1A4c i (3.1%) | | | | 82.6 |
| NH₃ | 3B1a (25.9%) | 3Da2c (10.4%) | 3Da1 (8.8%) | 3Da2a (7.9%) | 3B4a (6.4%) | 1B2d (6.3%) | 3B4g ii (4.5%) | 3B4g i (4.5%) | 1A3b i (4.3%) | 5B2 (3.2%) | | | | 82.1 |
| NMVOC | 1A3b i (26.3%) | 1A4b i (13.5%) | 2D3a (11.3%) | 1A3b iv (7.7%) | 1A4a i (4.1%) | 1A4c ii (3.7%) | 3B1b (2.5%) | 2D3g (2.4%) | 3B1a (2.3%) | 2D3d (2.2%) | 2H2 (2.2%) | 1B2a v (2.0%) | | 80.4 |
| CO | 1A3b i (43.0%) | 1A4b i (42.6%) | | | | | | | | | | | | 85.6 |
| PM₁₀ | 1A4b i (35.8%) | 1A1a (14.2%) | 1A4c ii (5.4%) | 1A3b i (5.4%) | 1A3biii (4.5%) | 1A3b ii (3.5%) | 3Dc (2.3%) | 1A2a (2.2%) | 1A1b (2.0%) | 1A3b vi (1.9%) | 1A2c (1.7%) | 5E (1.7%) | | 80.8 |
| PM_{2.5} | 1A4b i (40.9%) | 1A1a (12.0%) | 1A3b i (7.0%) | 1A4c ii (7.0%) | 1A3b iii (5.8%) | 1A3b ii (4.5%) | 1A2a (2.3%) | 5E (1.9%) | | | | | | 81.3 |
| BC | 1A4b i (36.8%) | 1A4c ii (14.5%) | 1A3b iii (9.6%) | 1A3b ii (8.2%) | 1A3b i (7.9%) | 1A2gvii (4.1%) | | | | | | | | 81.1 |
| Pb | 1A3b i (42.6%) | 2C1 (19.9%) | 1A2f (18.9%) | | | | | | | | | | | 81.4 |
| Cd | 1A2b (28.3%) | 2C1 (13.8%) | 1A2f (11.8%) | 1A2a (10.9%) | 5C2 (7.9%) | 2G (5.7%) | 1A3b i (4.8%) | | | | | | | 83.2 |
| Hg | 2C1 (37.7%) | 2B10a (23.8%) | 1B2d (21.0%) | | | | | | | | | | | 82.5 |
| PAH | 1A4b i (46.1%) | 2C1 (44.3%) | | | | | | | | | | | | 90.4 |
| Dioxin | 1A4b i (17.8%) | 1A4a i (16.5%) | 1A2a (13.3%) | 2C1 (12.8%) | 1A2b (9.1%) | 5E (7.9%) | 5C1a (7.2%) | | | | | | | 84.6 |
| HCB | 3Df (43.9%) | 1A2b (23.8%) | 1A4b i (7.4%) | 1A4a i (6.9%) | | | | | | | | | | 82.0 |
| PCB | 1A2a (42.7%) | 2C1 (25.3%) | 1A4b i (18.2%) | | | | | | | | | | | 86.2 |
| 1 Energy | 2 IPPU - Solvent and product use | | | 5 Waste | | | | | | | | | | |
| 2 IPPU - Industry | 3 Agriculture | | | | | | | | | | | | | |

1.7 QA/QC AND VERIFICATION METHODS

ISPRA has elaborated an inventory QA/QC procedures manual which describes specific QC procedures to be implemented during the inventory development process, facilitates the overall QA procedures to be conducted, as far as possible, on the entire inventory and establishes quality objectives (ISPRA, 2014). Specific QA/QC procedures and different verification activities implemented thoroughly in the current inventory compilation are figured out in the annual QA/QC plans (ISPRA, 2022 [b]).

Quality control checks and quality assurance procedures together with some verification activities are applied both to the national inventory as a whole and at sectoral level. Future planned improvements are prepared for each sector by the relevant inventory compiler; each expert identifies areas for sectoral improvement based on his own knowledge and in response to different inventory review processes.

In addition to *routine* general checks, source specific quality control procedures are applied on a case by case basis, focusing on key categories and on categories where significant methodological and data revision have taken place or new sources.

Checklists are compiled annually by the inventory experts and collected by the QA/QC coordinator. These lists are also registered in the ‘reference’ database.

General QC procedures also include data and documentation gathering. Specifically, the inventory analyst for a source category maintains a complete and separate project archive for that source category; the archive includes all the materials needed to develop the inventory for that year and is kept in a transparent manner.

Quality assurance procedures regard different verification activities of the inventory.

Feedbacks for the Italian inventory derive from communication of data to different institutions and/or at local level. Emission figures are also subjected to a process of re-examination once the inventory, the inventory related publications and the national inventory reports are posted on website, specifically www.isprambiente.gov.it.

The preparation of environmental reports where data are needed at different aggregation levels or refer to different contexts, such as environmental and economic accountings, is also a check for emission trends. At national level, for instance, emission time series are reported in the Environmental Data Yearbooks published by the Institute, in the Reports on the State of the Environment by the Ministry for the Environment, Land and Sea and, moreover, figures are communicated to the National Institute of Statistics to be published in the relevant Environmental Statistics Yearbooks as well as used in the framework of the EUROSTAT NAMEA Project.

Technical reviews of emission data submitted under the CLRTAP convention are undertaken periodically for each Party. Specifically, an in-depth review of the Italian inventory was carried out in 2010 and 2013 (UNECE, 2010; UNECE, 2013). A summary of the main findings of the last review can be found in the relevant technical report at the address

http://www.ceip.at/fileadmin/inhalte/emep/pdf/2013_s3/ITALY-Stage3ReviewReport-2013.pdf.

Moreover, under the European National Emission Ceiling Directive (NECD), an in-depth review has been conducted in 2017, 2018, 2019, 2020 and in 2021 (EEA, 2017 [a]; EEA 2018; EEA 2019; EEA 2020; EEA 2021). The main resulting findings and how the recommendations were addressed are reported in the following table.

Recommendations from TERT, considering revised estimates (RE) and technical corrections (TC)

| Observation | Key Category | NFR, Pollutant(s), Year(s) | Recommendation | RE or TC | Implementation |
|-------------------|--------------|---|---|----------|----------------|
| IT-2A5b-2021-0001 | Yes | 2A5b Construction and Demolition, PM10, 1990-2019 | For 2A5b Construction and Demolition for PM10 for 1990-2019 the TERT noted that there may be an under-estimate of emissions. In response to a question raised during the review Italy explained that it has collected statistical activity data on residential and non-residential building construction permits (from 1995 on) that will allow to apply a Tier 1 approach while for road construction it is still evaluating the proper indicator to calculate the annual activity data because only | No | Implemented |

| Observation | Key Category | NFR, Pollutant(s), Year(s) | Recommendation | RE or TC | Implementation |
|---------------------|--------------|---|--|----------|-----------------|
| | | | <p>economic statistics are available. The TERT is not able to determine whether the issue is above or below the threshold of significance for a technical correction.</p> <p>The TERT recommends that Italy includes all the available estimates in its next submission, including a description of the method, AD and EF in the IPPU chapter of the IIR.</p> | | |
| IT-2D3c-2021-0001 | Yes | 2D3c Asphalt Roofing, CO, 1990-2019 | <p>For 2D3c Asphalt Roofing for CO for 1990-2019 the TERT noted that the notation key 'NA' (not applicable) is used while a Tier 1 method is available in the 2019 EMEP/EEA Guidebook. In response to a question raised during the review Italy explained that CO emissions will be included in its next submission. The TERT noted that the issue is below the threshold of significance for a technical correction.</p> <p>The TERT recommends that Italy includes the emissions in its next submission and describes the method, the AD and EF in the IPPU chapter of the IIR.</p> | No | Implemented |
| IT-5C1bii-2021-0002 | No | 5C1bii Hazardous Waste Incineration, SO ₂ , NO _x , NMVOC, PM _{2.5} , BaP, PAHs, PCBs, HCB, Cd, Hg, Pb, PCDD/F, PM ₁₀ , CO, BC, TSP, 2008-2019 | <p>For 5C1bii Hazardous Waste Incineration and all pollutants and years 2008-2019 the TERT noted that there is a lack of transparency regarding the different use of the notation keys in the NFR tables between the emissions part ('NA' is used) and the part of the activity data for 'other activity' ('NO' is used) and that no explanation on the use of the notation keys for this source is given in the IIR. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained for this source that indeed there is no incineration of hazardous waste in the period 2008-2019 and that suggested for the emissions to use 'NO' for this period. The TERT recommends that Italy uses the notation key 'NO' for both emissions and activity data when the process is no longer occurring and explains this in the IIR in the 2022 submission.</p> | No | Not implemented |
| IT-1A3c-2021-0001 | No | 1A3c Railways, PM _{2.5} , 1990-2009 | <p>For 1A3c, PM_{2.5}, years 1990-2019, the TERT noted PM_{2.5} emissions are equal to PM₁₀ whilst the 2019 EMEP/EEA Guidebook provides different emission factors for PM₁₀ and PM_{2.5} in Table 3.1. In response to a question raised during the review the Italy stated that they will revise the estimates in the next submission. The TERT noted that the issue is below the threshold of significance for a technical correction.</p> <p>The TERT recommends that in the 2022 submission Italy applies the methodology for PM_{2.5} emissions for 1A3c from the 2019 EMEP/EEA Guidebook or update the IIR to explain the methodology used.</p> | No | Implemented |

| Observation | Key Category | NFR, Pollutant(s), Year(s) | Recommendation | RE or TC | Implementation |
|--------------------|--------------|---|---|----------|------------------------|
| IT-1A4ci-2021-0001 | No | 1A4ci Agriculture/Forestry/Fishing: Stationary, PM2.5, 2000-2019 | <p>For category 1A4ci and pollutants PM10 and PM2.5 the TERT noted that there is a lack of transparency regarding the reason why PM10 and PM2.5 emissions are the same despite significant use of non-gaseous fuels in recent years. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that it intends to review emission factors used for biomass.</p> <p>The TERT recommends that Italy completes the review and, implements and reports the results in the 2022 submission.</p> | No | Implemented |
| IT-2A1-2019-0001 | Yes | 2A1 Cement Production, PM2.5, 1990-2019 | <p>For 2A1 cement production for PM2.5 for 1990-2019 the TERT noted that there is a lack of transparency regarding the method AD and EF used for the estimate. This does not relate to an over- or under-estimate of emissions. This was raised during the 2019 and 2020 NECD inventory review. In response to a question raised during the review, Italy explained that the relevant paragraph of the IIR will be updated in the next submission and that the correct information about the PM10 recalculation has been reported in the relevant paragraph 4.5.1.</p> <p>The TERT recommends that Italy includes in its next IIR the description for the method, AD and EF for this category and pollutant.</p> | No | Implemented |
| IT-2A5a-2017-0001 | No | 2A5a Quarrying and Mining of Minerals Other Than Coal, PM2.5, 1990-2019 | <p>For 2A5a Quarrying and Mining of Minerals Other Than Coal for PM2.5 for 1990-2019 the TERT noted that there is a lack of transparency regarding the further investigations that Italy planned and the timeframe for those. This does not relate to an over- or under-estimate of emissions. This was raised during the 2017, 2018, 2019 and 2020 NECD inventory review. In response to a question raised during the review, Italy explained that it plans different checks and comparison with bottom-up inventories and an in-depth study is expected in 2022. Consequently, the estimates at national level should be ready in 2023.</p> <p>The TERT recommends that Italy includes in its next IIR the progress of these investigations and confirms or updates the time frame accordingly.</p> | No | Not fully implemented. |
| IT-2A5b-2017-0001 | No | 2A5b Construction and Demolition, PM2.5, 1990-2019 | <p>For 2A5b Construction and Demolition for PM2.5 for 1990-2019 the TERT noted that there may be an under-estimate of emissions. This was raised during the 2017, 2018, 2019 and 2020 NECD inventory review. In response to a question raised during the review Italy explained more in detail about the further investigations that are under way (IIR p. 114). Italy explained that it has collected statistical activity data on residential and non-residential building construction permits (from 1995 on) that will allow to apply a Tier 1 approach while</p> | No | Implemented |

| Observation | Key Category | NFR, Pollutant(s), Year(s) | Recommendation | RE or TC | Implementation |
|--------------------|--------------|--|---|----------|----------------|
| | | | <p>for road construction it is still evaluating the proper indicator to calculate the annual activity data because only economic statistics are available. The TERT is not able to determine whether the issue is above or below the threshold of significance for a technical correction.</p> <p>The TERT recommends that Italy includes all the available estimates in its next submission, including a description of the method, AD and EF in the IPPU chapter of the IIR.</p> | | |
| IT-2D3e-2017-0001 | No | 2D3e Degreasing, NMVOC, 1990-2019 | <p>For 2D3e Degreasing NMVOC emissions for 1990-2019 the TERT noted that there is a lack of transparency regarding the implementation of the recommendation to improve the methodology by obtaining the information on the composition of cleaning products regarding the different NMVOC compounds or (in case it would be not possible) to take into account the whole amount of used cleaning products. This does not relate to an over- or under-estimate of emissions. This was raised during the 2017, 2018, 2019 and 2020 NECD inventory review. In response to a question raised during the review, Italy explained that it used as activity data the amount of solvent used and that in absence of information on the abatement technologies, as suggested for Tier 2, the assumption is that the system is uncontrolled, and all solvent is emitted. Italy will check this assumption with the relevant industrial association for the next submission.</p> <p>The TERT recommends that Italy clearly describes the method, EF and AD in the IPPU chapter in its next submission.</p> | No | Implemented |
| IT-2D3g-2021-0001 | No | 2D3g Chemical Products, PM2.5, 1990-2019 | <p>For 2D3g Chemical Products for 1990-2019 the PM2.5 estimate is equal to the estimate for PM10. The TERT would expect that for this category, PM10 estimates are higher than (rather than equal to) PM2.5 estimates. The TERT noted that there is a lack of transparency regarding the reason for this. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that PM emissions come from polyester and polyvinylchloride processing and EFs have been provided by the relevant industry and that no PM EFs are provided in the 2019 EMEP/EEA Guidebook for this activity.</p> <p>The TERT recommends that Italy checks with the relevant industry on the method for determining the EFs, why the EF for PM2.5 is the same as that for PM10 and provides information on this in its next IIR.</p> | No | Implemented |
| IT-5C1bi-2021-0001 | No | 5C1bi Industrial Waste Incineration, | <p>For 5C1bi (Industrial Waste Incineration) and PM2.5 and years 1990-2007 the TERT noted that the PM2.5 and PM10 emissions reported</p> | No | Implemented |

| Observation | Key Category | NFR, Pollutant(s), Year(s) | Recommendation | RE or TC | Implementation |
|---------------------|--------------|---|--|----------|----------------|
| | | PM2.5, 1990-2009 | <p>in the NFR tables are equal. In response to a question raised during the review Italy explained that the PM10-emissions relate to a personal communication (in 1999) in the period that only PM10 needed to be considered and now uses the same emission factor as conservative estimate for PM2.5. In their answer Italy also suggested that they could use the EMEP/EEA Guidebook PM10/PM2.5 ratio to correct this. The TERT notes that using PM2.5 as being a fraction of PM10 is in line with the EMEP/EEA Guidebook where in fact this is also the case as the units for PM10 and PM2.5 are the same. The TERT noted that the issue is below the threshold of significance for a technical correction.</p> <p>The TERT recommends that Italy in the 2022 submission of the NFR and IIR explains the personal communication for using the PM10 emissions factor for the period 1990-2007 and calculates the PM2.5 emissions over that period with the PM10/PM2.5 ratio from the EMEP/EEA Guidebook.</p> | | |
| IT-5C1bii-2021-0001 | No | 5C1bii Hazardous Waste Incineration, PM2.5, 1990-2007 | <p>For 5C1bii Hazardous Waste Incineration and PM2.5 and years 1990-2007 the TERT noted that the PM2.5 and PM10 emissions reported in the NFR tables are equal. In response to a question raised during the review Italy explained that the PM10 emissions relate to a personal communication (in 1999) in the period that only PM10 needed to be considered and now uses the same emission factor as conservative estimate for PM2.5. In their answer Italy also suggested that they could use the EMEP/EEA Guidebook PM10/PM2.5 ratio to correct this. The TERT notes that using PM2.5 as being a fraction of PM10 is in line with the EMEP/EEA Guidebook where in fact this is also the case as the units for PM10 and PM2.5 are the same. The TERT noted that the issue is below the threshold of significance for a technical correction.</p> <p>The TERT recommends that Italy in the 2022 submission of the NFR and IIR explains the personal communication for using the PM10 emissions factor for the period 1990-2007 and calculates the PM2.5 emissions over that period with the PM10/PM2.5 ratio from the EMEP/EEA Guidebook.</p> | No | Implemented |
| IT-5C1biv-2021-0001 | No | 5C1biv Sewage Sludge Incineration, PM2.5, 1990-2009 | <p>For 5C1biv (Sewage Sludge Incineration) and PM2.5 and years 1990-2009 the TERT noted that the PM2.5 and PM10 emissions reported in the NFR tables are equal. In response to a question raised during the review Italy explained that the PM10 emissions relate to a personal communication (in 1999) in the period that only PM10 needed to be considered and now uses the same emission factor as conservative estimate for PM2.5. In their answer Italy also suggested that they could use the EMEP/EEA Guidebook PM10/PM2.5 ratio to correct this. The TERT notes that using PM2.5 as being a fraction of PM10 is in line with the EMEP/EEA</p> | No | Implemented |

| Observation | Key Category | NFR, Pollutant(s), Year(s) | Recommendation | RE or TC | Implementation |
|-------------------|--------------|---|---|----------|-----------------|
| | | | <p>Guidebook where in fact this is also the case as the units for PM10 and PM2.5 are the same. The TERT noted that the issue is below the threshold of significance for a technical correction.</p> <p>The TERT recommends that Italy in the 2022 submission of the NFR and IIR explains the personal communication for using the PM10 emissions factor for the period 1990-2009 and calculates the PM2.5 emissions over that period with the PM10/PM2.5 ratio from the EMEP/EEA Guidebook.</p> | | |
| IT-1A2b-2019-0001 | No | 1A2b Stationary Combustion in Manufacturing Industries and Construction: Non-Ferrous Metals, PCBs, HCB, 1990-2018 | <p>For category 1A2b Stationary Combustion in Manufacturing Industries and Construction: Non-Ferrous Metals and pollutants HCB and PCB for all years the TERT noted that the notation key 'NA' (Not Applicable) was used in the 2021 submission. This was raised during 2019 and 2020 NECD inventory review. In response to a question raised during the review, Italy explained that emissions had been calculated but had been omitted from the 2021 submission. Italy provided a revised estimate for all years and stated that it will be included in the next submission. The TERT agreed with the revised estimate provided by Italy.</p> <p>The TERT recommends that Italy include the revised estimate in its 2022 NFR and IIR submission.</p> | RE | Implemented |
| IT-5E-2019-0001 | No | 5E Other Waste, PM2.5, Cd, Hg, Pb, PCDD/F, PM10, TSP, 1990-2019 | <p>For 5E Other Waste the TERT noted from the NFR tables and IIR pages 157-158 that there may be an under-estimate of emissions for the years 1990-2019. This was raised during 2019 and 2020 NECD inventory review. In response to a question raised during the review, Italy explained that the emission factors reported in the EMEP/EEA Guidebook are inadequate to the Italian reality and, in general, highly uncertain. However, Italy also provided revised estimates for the years 1990-2019 based on the EMEP/EEA Guidebook Tier 2 methodology and including all accidental building fires for PM10, PM2.5 and PCDD/F and BC based on the IIASA, 2004 (Interim Report IR-04-079). The TERT noted that in the estimates an EF for PM2.5 of 27.33 kg/fire is used instead of the EF from the EMEP/EEA Guidebook of 27.23 kg/fire. This relates possibly to a typing error. The TERT agreed with the revised estimate provided by Italy.</p> <p>The TERT recommends that Italy checks the EF used for PM2.5 and explains in the IIR in more detail the origin of the EF used for BC and include the revised estimates and the emissions from heavy metals from this source in its 2022 NFR and IIR submission.</p> | RE | Implemented |
| IT-0A-2020-0002 | No | 0A National Total - National total for the entire territory - Based on fuel sold/fuel used, | The TERT noted that no information on uncertainty was provided for BC, PM10 and CO. The TERT recommends that the guidance given in "Part A 5 Uncertainties 2019" of the 2019 EMEP/EEA Guidebook is used to assess the uncertainty by pollutant | No | Not implemented |

| Observation | Key Category | NFR, Pollutant(s), Year(s) | Recommendation | RE or TC | Implementation |
|------------------|--------------|--|---|----------|----------------|
| | | SO ₂ , NO _x , NH ₃ , NMVOC, PM _{2.5} , BaP, PAHs, PCBs, HCB, Cd, Hg, Pb, PCDD/F, PM ₁₀ , CO, 2018 | and sector. The TERT further recommends that the methods used for the assessment of the uncertainty are described in the IIR transparently along with a summary of uncertainties by pollutant and sector as requested in the “Recommended Structure for Informative Inventory Report” (Annex II_v2018 to the revised 2014 reporting guidelines). | | |
| IT-1A1-2019-0001 | No | 1A1 Energy Production, BaP, PAHs, PCBs, HCB, Cd, Hg, Pb, PCDD/F, 1990 - 2017 | For category 1A1 Energy Production and pollutants heavy metals and persistent organic pollutants the TERT noted that there is a lack of transparency regarding whether Italy has reviewed current country-specific emission factors against those in the EEA Guidebook 2019 and plant-specific data (where available). The TERT notes that improvements in transparency have been provided in the IIR and in the ISPRA spreadsheet of emission factors (referenced in the IIR). This was raised during the 2019 and 2020 NECD inventory review and does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that work has commenced on reviewing emission factors for 1A1 (also 1A2 and 1A4) and transparency has been improved by colour-coding of emission factors in supplementary documentation for the IIR. Furthermore, Italy has indicated that there will be a reorganisation to the IIR structure for the energy activities for the 2022 submission to improve transparency. The TERT recommends that the reorganisation of the IIR is completed for the 2022 submission and that progress on reviewing the emission factors is documented in the IIR. | No | Implemented |
| IT-1A2-2019-0001 | Yes | 1A2 Stationary Combustion in Manufacturing Industries and Construction, BaP, PAHs, PCBs, HCB, Cd, Hg, Pb, PCDD/F, 1990-2018 | For category 1A2 Stationary Combustion in Manufacturing Industries and Construction and pollutants heavy metals and persistent organic pollutants the TERT noted that there is a lack of transparency regarding whether Italy has reviewed current country-specific emission factors against those in the EEA Guidebook 2019 and plant-specific data (where available). The TERT notes that improvements in transparency have been provided in the IIR and in the ISPRA spreadsheet of emission factors (referenced in the IIR). This was raised during the 2019 and 2020 NECD inventory review and does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that work has commenced on reviewing emission factors for 1A2 (also 1A1 and 1A4) and transparency has been improved by colour-coding of emission factors in supplementary documentation for the IIR. Furthermore, Italy has indicated that there will be a reorganisation to the IIR structure for the energy activities for the 2022 submission to improve transparency. | No | Implemented |

| Observation | Key Category | NFR, Pollutant(s), Year(s) | Recommendation | RE or TC | Implementation |
|-------------------|--------------|--|--|----------|-----------------|
| | | | The TERT recommends that the reorganisation of the IIR is completed for the 2022 submission and that progress on reviewing the emission factors is documented in the IIR. | | |
| IT-1A2a-2019-0001 | Yes | 1A2a Stationary Combustion in Manufacturing Industries and Construction: Iron and Steel, PCBs, HCB, Cd, 2005, 2016, 2017 | For category 1A2a Stationary Combustion in Manufacturing Industries and Construction: Iron and Steel and pollutants Cd and HCB the TERT noted that the IEF ratios are outliers when compared to other Member States. This was raised during the 2019 NECD inventory review. In response to a question raised during the review, Italy explained that for HCB, the emission factors used by the inventory are based on measurements from 2007 and the derived emission factors are applied across the time series as no other emission factors are available. For Cd, Italy has made changes to estimation methodology in the 2021 submission based on a revised estimate from the 2020 NECD inventory review. Italy has also commenced a review of HCB emission factors. The TERT recommends that Italy reviews the methodology for HCB emission estimates and reports in the 2022 submission. | No | Implemented |
| IT-1A2b-2019-0002 | Yes | 1A2b Stationary Combustion in Manufacturing Industries and Construction: Non-Ferrous Metals, PCDD/F, 1990-2018 | For 1A2b Stationary Combustion in Manufacturing Industries and Construction: Non-Ferrous Metals and pollutant PCDD/F the TERT noted that there is a lack of transparency regarding allocation of emissions between activity 1A2b and 2C activities. This was raised during the 2019 and 2020 NECD inventory review and does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained it intends to explore the allocation of PCDD/F emissions between 1A2b and 2C activities alongside a wider review of emission factors used in 1A2. The TERT recommends that Italy completes this improvement and reports results in the 2022 submission. | No | Not implemented |
| IT-2C1-2021-0002 | Yes | 2C1 Iron and Steel Production, PAHs, 1990, 2005, 2016, 2017, 2018, 2019 | For emissions of PAHs for years 1990, 2005, 2016-2019 for 2C1 Iron and Steel Production the TERT noted that there is a lack of transparency regarding whether or not emissions from blast furnace charging, basic oxygen furnace and sinter production are estimated. From the description in the IIR it is not clear since the IIR (on p. 106) does not include an EF for these subcategories while there exists a method end EF in the EMEP 2019 Guidebook. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that further information is given in the combustion chapter and that emissions are estimated on the basis of country specific emission factors at activity level, especially referring to sinter plants production, as provided by the main national operators and it is not possible to split up between combustion and process. For | No | Implemented |

| Observation | Key Category | NFR, Pollutant(s), Year(s) | Recommendation | RE or TC | Implementation |
|------------------|--------------|---|--|----------|---------------------|
| | | | blast furnaces, results from measurements learn that the EF used for pig iron tapping covers also blast furnace charging. Emissions from BOF are negligible. Based on the data collected data in the context of the environmental authorisation of the largest Italian (and European) plant, Italy is reviewing the emissions of all pollutants starting from 2015 based on the measurement data for each section of the installation and for each chimney. The revision of the estimates is foreseen for the next submission. The TERT recommends that Italy includes a clear explanation for the estimate of PAH emissions in the IPPU chapter or a clear reference to the relevant chapters elsewhere and that Italy includes the reviewed emissions from 2015 on, including for a clear explanation on the method and data used. | | |
| IT-2C3-2020-0001 | No | 2C3 Aluminium Production, PM _{2.5} , HCB, PCDD/F, PM ₁₀ , BC, 1990-2018 | For 2C3 Aluminium Production for PM _{2.5} , HCB, PCDD/F, PM ₁₀ and BC for 1990-2018 the TERT noted that there is a lack of transparency regarding the use of the notation key 'IE'. This does not relate to an over- or under-estimate of emissions. This was raised during the 2020 NECD inventory review. In response to a question raised during the review, Italy provided further clarification of how emissions are estimated in 1A2b by providing the AD and the country specific EF. The TERT recommends that Italy reports these emissions under 2C3 in its next submission and reports on the method, AD and EF in the IPPU section of the IIR. | No | Partly implemented. |
| IT-5C-2019-0001 | Yes | 5C Waste Incineration, PCBs, HCB, 1990-2018 | For 5C (Waste incineration) and pollutants PCBs and HCB and years 1990-2019 the TERT noted that there may be an over-estimate of emissions. This was raised during the 2019 and 2020 NECD inventory review. This over-estimate does not have an impact on total emissions that is above the threshold of significance. In the 2020 review the TERT reiterated the recommendation from the 2019 review to take in account the implementation of abatement technology over the years for those sources where this is not considered yet (for instance 5C1bv). The 2021 review noted that in the IIR (pg 153) is stated that the results of this will be available in the 2022 submission. In reply to a question during the review Italy confirmed that this is still the plan. The TERT reiterates the recommendation to take in account the implementation of abatement technology over the years for those sources where this is not considered yet (for instance 5C1bv). The TERT additionally, reiterates the recommendation that Italy further improves the transparency in the IIR and to use the emission factors from the survey on incineration of sewage sludge in the 2022 submission. | No | Partly implemented |

| Observation | Key Category | NFR, Pollutant(s), Year(s) | Recommendation | RE or TC | Implementation |
|----------------------|--------------|--|--|----------|---|
| IT-LPS-GEN-2020-0003 | | General, 2015 | The TERT noted that in 2015, the same pair of longitude and latitude coordinates was assigned to more than one differently named LPS in 53 cases. The TERT recommends that Italy ensures that it provides unique longitude and latitude coordinates for each LPS in its future submissions. | No | Implemented |
| IT-LPS-GEN-2020-0001 | | General, PM _{2.5} , HCB, 2015 | For the LPS reporting, the TERT noted that emissions of several pollutants were not reported, including HCB and PM _{2.5} . In response to a question raised during the review Italy explained that LPS reporting is being prepared based on E-PRTR data which does not cover HCB and PM _{2.5} but only total particles. The TERT noted that the issue is below the threshold of significance for a technical correction. The TERT recommends that Italy clarifies whether any LPS have HCB and PM _{2.5} emission above the reporting threshold and report any emissions above the reporting threshold in the next submission. | No | Implemented |
| IT-LPS-A-2020-0002 | | A Public Power, PM _{2.5} , 2015 | The TERT recommends that Italy consider how to include PM _{2.5} emissions for the LPS where PM ₁₀ emissions are reported for the next submission. | No | Implemented |
| IT-LPS-A-2020-0001 | | A Public Power, SO ₂ , NO _x , NH ₃ , NMVOC, PM _{2.5} , BaP, PAHs, PCBs, HCB, Cd, Hg, Pb, PCDD/F, PM ₁₀ , CO, BC, 2015 | The TERT recommends that Italy provide information to summarise the LPS dataset and how it was developed. Italy may wish to consider the recommended structure for an Informative Inventory Report contained in the 2014 Guidelines for Estimating and Reporting Emission Data | No | Implemented, more info in the IIR, chap. 10 |
| IT-LPS-D-2020-0001 | | D Fugitive, 2015 | For LPS-D and NFR 1B2aiv Oil: refining/storage there may be a case on noncompliant reporting. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that emissions from refineries were reported in a different sector and that in the next submission they will report LPS data according to the different categories involved at plant level. Emission will be provided separately in the industry or fugitive sector according to their NFR codes and point source emission data which are available at NFR category level. The TERT recommends that Italy amends the reporting of fugitive and process emissions to be consistent with LPS requirements for the 2021 submission. | No | Implemented |
| IT-LPS-K-2020-0001 | | K Agriculture Livestock, NH ₃ , 2015 | The TERT notes that for the year 2015, emissions are reported for 14 agricultural facilities in the E-PRTR database (v18) at a far higher level than they are reported in the LPS submission. In response to the review Italy indicated that this was an error in the LPS submission, and that the E-PRTR facility data is correct. The TERT recommends that Italy corrects this in the next LPS submission and provides any analytical comparison of E-PRTR facility | No | Implemented |

| Observation | Key Category | NFR, Pollutant(s), Year(s) | Recommendation | RE or TC | Implementation |
|-----------------------|--------------|--|---|----------|-------------------------------|
| | | | data and national inventory estimates in its IIR description of LPS and Gridded estimates in future submissions. | | |
| IT-GRID-GEN-2020-0001 | | General, NH ₃ , NMVOC, NO _x , PM _{2.5} , 2015 | The TERT recommends that Italy makes improvements to the gridding methods for the 2021 submission with distributions that reflect activities in all relevant GNFR sector for all pollutants and also provides a chapter in the IIR outlining the methods used to generate all of the gridded estimates, in accordance with the requirements outlined in the 2019 EMEP/EEA Guidebook. For example, regarding the F_RoadTransport sector Italy should follow methodology in section '3.4.3 Line sources to grids', the line sources such as road network should be intersected with the grid cells and allocate the emissions to the relevant grid cells proportionally. | No | More info in the IIR, chap 10 |
| IT-GRID-B-2020-0001 | | B Industry, SO ₂ , NH ₃ , NMVOC, PM _{2.5} , PCBs, Cd, Hg, Pb, PCDD/F, PM ₁₀ , CO, 2015 | For the GNFR category B_Industry the TERT noted that there is a lack of transparency regarding a misallocation of emissions between B_Industry and D_Fugitive GNFR sectors and an allocation of B_Industry emissions to the sector name 'B_IndustrialComb' which is not consistent with the guidelines. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that in the next submission Italy will report gridded data according to the last available reporting guidelines and the relevant codes. The TERT recommends that Italy report gridded data according to the last available reporting guidelines and the relevant NFR codes in the next submission. | No | Implemented |
| IT-GRID-F-2020-0002 | | F Road Transport, NO _x , NMVOC, PM _{2.5} , PM ₁₀ , CO, 2015 | The TERT recommends that Italy report road transport and rail emissions in the relevant separate GNFR sector categories in the next submission. | No | Implemented |
| IT-GRID-F-2020-0001 | | F Road Transport, SO ₂ , NH ₃ , Cd, Pb, PCDD/F, PAHs, 2015 | The TERT recommends that Italy report road transport and rail emissions in the relevant separate GNFR sector categories in the next submission. | No | Implemented |

A bilateral independent review between Italy and Spain was undertaken in the year 2012, with a focus on the revision of emission inventories and projections of both the Parties. With regard to the emission inventory the Italian team revised part of the energy sector of Spain, specifically the public power plants, petroleum refining plants, road transport and off-road categories, whereas the Spanish team revised the Industrial processes and solvent and other product use, and the LULUCF sectors of Italy. Results of these analyses are reported in a technical report. Aim of the review was to carry out a general quality assurance analysis of the inventories in terms of methodologies, EFs and references used, as well as analysing critical cross cutting issues such as the details of the national energy balances and comparison with international data (EUROSTAT and IEA) and use of plant specific information.

In addition, an official independent review of the entire Italian inventory was undertaken by the Aether consultants in 2013. Main findings and recommendations are reported in a final document, and regard mostly the transparency in the NIR, the improvement of QA/QC documentation and some pending issues in the LULUCF sector. These suggestions were considered in the implementation of the following inventories.

Comparisons between national activity data and data from international databases are usually carried out in order to find out the main differences and an explanation to them. Emission intensity indicators among countries (e.g. emissions per capita, industrial emissions per unit of added value, road transport emissions per passenger car, emissions from power generation per kWh of electricity produced, emissions from dairy cows per tonne of milk produced) can also be useful to provide a preliminary check and verification of the order of magnitude of the emissions. Additional comparisons between emission estimates from industrial sectors and those published by the industry itself in the Environmental reports are carried out annually in order to assess the quality and the uncertainty of the estimates.

The quality of the inventory has also improved by the organization and participation in sector specific workshops.

A specific procedure undertaken for improving the inventory regards the establishment of national expert panels (in particular, in road transport, land use change and forestry and energy sectors) which involve, on a voluntary basis, different institutions, local agencies and industrial associations cooperating for improving activity data and emission factors accuracy.

Furthermore, activities in the framework of the improvement of local inventories are carried out together with local authorities concentrating on the comparison between top down and bottom up approaches and identifying the main critical issues. In 2021, ISPRA has finalised the provincial inventory at local scale for the years 1990, 1995, 2000, 2005, 2010, 2015 and 2019 applying a top down approach. Methodologies and results were checked out by regional and local environmental agencies and authorities, and figures are available at ISPRA web address [Serie Storiche Emissioni – EMISSIONI \(isprambiente.it\)](https://www.isprambiente.it/serie-storiche-emissioni). Methodologies used for a previous reporting cycle are described in a related publication (ISPRA, 2009).

This work is also relevant to carry out regional scenarios, for the main pollutants, within the Gains Italy project implemented by ENEA and supported by ISPRA and the regional authorities.

In addition to these expert panels, ISPRA participates in technical working groups within the National Statistical System. These groups, named *Circoli di qualità* (“Quality Panels”), coordinated by the National Institute of Statistics, are constituted by both producers and users of statistical information with the aim of improving and monitoring statistical information in specific sectors such as transport, industry, agriculture, forest and fishing. These activities should improve the quality and details of basic data, as well as enable a more organized and timely communication.

Other specific activities relating to improvements of the inventory and QA/QC practices regard the progress on management of information collected in the framework of different European obligations, Large Combustion Plant, E-PRTR and Emissions Trading, which is gathered together in an informative system thus highlighting the main discrepancies among data, detecting potential errors and improving the time series consistency. ISPRA collects these data from the industrial facilities and the inventory team manages the information and makes use of it in the preparation of the national inventory. The informative system is based on identification codes to trace back individual point sources in different databases and all the figures are considered in an overall approach and used in the compilation of the inventory.

A proper archiving and reporting of the documentation related to the inventory compilation process is also part of the national QA/QC programme.

All the material and documents used for the inventory preparation are stored at the Institute for Environmental Protection and Research.

Information relating to the planning, preparation, and management of inventory activities are documented and archived. The archive is organised so that any skilled analyst could obtain relevant data sources and spreadsheets, reproduce the inventory and review all decisions about assumptions and methodologies undertaken. A master documentation catalogue is generated for each inventory year and it is possible to track changes in data and methodologies over time. Specifically, the documentation includes:

- electronic copies of each of the draft and final inventory report, electronic copies of the draft and final NFR tables;
- electronic copies of all the final, linked source category spreadsheets for the inventory estimates (including all spreadsheets that feed the emission spreadsheets);
- results of the reviews and, in general, all documentation related to the corresponding inventory year submission.

After each reporting cycle, all database files, spreadsheets and electronic documents are archived as ‘read-only’ mode.

A ‘reference’ database is also compiled every year to increase the transparency of the inventory. This database consists of a number of records that references all documentation used during the inventory

compilation, for each sector and submission year, the link to the electronically available documents and the place where they are stored as well as internal documentation on QA/QC procedures.

1.8 GENERAL UNCERTAINTY EVALUATION

An overall uncertainty analysis for the Italian inventory related to the pollutants described in this report has not been assessed yet. Nevertheless, different studies on uncertainty have been carried out (Romano et al., 2004) and a quantitative assessment of the Italian GHG inventory is performed by the Tier 1 method defined in the 2006 IPCC Guidelines (IPCC, 2006) which provides a calculation based on the error propagation equations. Details on the results of the GHG inventory uncertainty figures can be found in the *National Inventory Report 2022* (ISPRA, 2022 [a]).

It should be noted that different levels of uncertainty pertain to different pollutants. Estimates of the main pollutants are generally of high level, but PM emissions, especially those of small particle sizes, heavy metal and POP estimates are more uncertain. For this reason, even though not quantified in terms of uncertainty, improvements are planned especially for the specified pollutants.

Nevertheless, since quantitative uncertainty assessments constitute a mean to either provide the inventory users with a quantitative assessment of the inventory quality or to direct the inventory preparation team to priority areas, a planned improvement for next submissions is the completion of such analysis.

1.9 GENERAL ASSESSMENT OF COMPLETENESS

The inventory covers all major sources, as well as all main pollutants, included in the UNECE reporting guidelines (UNECE, 2014). NFR sheets are complete as far as the details of basic information are available.

Allocation of emissions is not consistent with the guidelines only where there are no sufficient data available to split the information. For instance, emissions from category 1.A.5.a other stationary are reported and included under category 1.A.4.a i commercial and institutional emission estimates. Mobile commercial and institutional emission estimates (1.A.4.a ii) are included in 1.A.3 sector. PM and HMs emissions from 2.A.3 glass production are included in 1.A.2.f combustion category source as well as those from lead, zinc and copper production are included in 1.A.2.b category. HCB, PCB and Dioxin emissions from aluminium production are included in 1.A.2.b category as well as PM emissions from secondary aluminium production while HCB from iron and steel are included in 1.A.2.a category. NO_x, SO_x and NH₃ from 1.B.1.b, fugitive emissions from solid fuel transformation, are included in the 1.A.2.a category and HMs and POPs from 1.B.2.a iv are included in 1.A.1.b category. PM emissions from storage, handling and transport of mineral products (2.A.5.c) are included under the relevant emission categories.

There are a few emission sources not assessed yet: HMs, PAH, dioxin and PCB non-exhaust emissions from 1.A.3.b vii, road abrasion, PAH, dioxin and PCB emissions from 1.A.3.b v gasoline evaporation, dioxin and PCB emissions from 1.A.3.b vi, automobile tyre and brake wear, NH₃ emissions from 1.A.3.a domestic and international aviation LTO cycle, NH₃ from 1.A.3.e i, pipeline transportation, NO_x and NH₃ from 3.D.a iv, crop residues applied to soils, and 3.D.b, indirect emissions from managed soils. Emission factors for these categories, when available in the Guidebook (EMEP/EEA, 2019), need further assessment for the applicability to the national circumstances. PAH emissions are not detailed in the four indicator compounds for all the categories; we should still estimate them for categories 1.A.1, 1.A.2 stationary, 1.A.3.a, 1.A.3.d ii, 1.A.4.c iii, 2.C.1, and 2.C.3, because for some categories emission factors are not fully available by compound. PM and black carbon emissions from the categories reported in the NFR under 2.A.5 a, quarrying and mining of minerals other than coal, are not estimated because no information on activity data is still available. Emissions of PAH from asphalt blowing, 2.D.3g, are also under further investigation and reported as NE, although according to the relevant industrial association PAH emissions are negligible because all the asphalt blowing plants have abatement filter system of PM and afterburners of gas. Moreover, these plants should respect national environmental legislation not exceeding at the stack more than 0.1mg/Nm³ for total PAH.

Further investigation will be carried out about these source categories and pollutants in order to calculate and improve figures.

2 ANALYSIS OF KEY TRENDS BY POLLUTANT

2.1 MAIN POLLUTANTS

In the following sections, Italian emission series of sulphur oxides, nitrogen oxides, non-methane volatile organic compounds, carbon monoxide and ammonia are presented.

2.1.1 Sulphur dioxide (SO_x)

The national atmospheric emissions of sulphur oxides have significantly decreased in recent years, as occurred in almost all countries of the UNECE.

Figure 2.1 and Table 2.1 show the emission trend from 1990 to 2020. Figure 2.1 also illustrates the share of SO_x emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.1 SO_x emissions trend, percentage share by sector and variation 1990-2020

Table 2.1 SO_x emission trend from 1990 to 2020 (Gg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|---------|-------|-------|-------|------|------|------|------|------|------|
| | Gg | | | | | | | | | |
| Combustion in energy and transformation industries | 1,000.8 | 776.4 | 466.8 | 187.0 | 77.1 | 29.6 | 19.1 | 16.9 | 13.4 | 10.9 |
| Non industrial combustion plants | 82.3 | 32.5 | 25.0 | 22.7 | 12.1 | 10.3 | 10.1 | 10.4 | 10.0 | 9.8 |
| Combustion - Industry | 324.1 | 230.7 | 119.8 | 91.4 | 60.0 | 35.4 | 33.8 | 30.2 | 31.9 | 31.5 |
| Production processes | 136.0 | 115.5 | 38.8 | 46.4 | 35.9 | 24.3 | 26.8 | 23.5 | 20.0 | 16.7 |

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|------------------------------------|----------------|----------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|-------------|
| | <i>Gg</i> | | | | | | | | | |
| Solvent and other product use | 0.009 | 0.009 | 0.032 | 0.040 | 0.032 | 0.017 | 0.013 | 0.013 | 0.013 | 0.008 |
| Road transport | 129.3 | 71.6 | 11.9 | 2.2 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.3 |
| Other mobile sources and machinery | 98.3 | 84.1 | 83.9 | 50.6 | 29.1 | 21.2 | 21.5 | 23.3 | 25.0 | 8.5 |
| Waste treatment and disposal | 12.9 | 11.6 | 10.0 | 10.7 | 7.0 | 4.5 | 4.7 | 4.3 | 3.8 | 4.0 |
| Agriculture | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Total | 1,783.6 | 1,322.4 | 756.4 | 411.0 | 221.7 | 125.7 | 116.5 | 109.1 | 104.6 | 81.9 |

Figures show a general decline of SO_x emissions during the period, from 1,784 Gg in 1990 to 82 Gg in 2020. The national target of SO_x emissions, set by the National Emission Ceilings Directive at 475 Gg for 2010 (EC, 2001) was reached and continues to be respected after this year revision of the time series. The targets established for 2020 in the framework of the UNECE/CLRTAP Convention and in the framework of 2030 of the revised National Emission Ceiling Directive (EU, 2016), equal for Italy respectively to 65% and 29% of 2005 emissions, has been already reached.

The decreasing trend is determined mainly by the reduction in emissions from *combustion in energy* (-99%) and in *industry* (-90%), representing in 2020 about 13%, and 39% of the total, respectively. Emissions deriving from *non-industrial combustion plants* and *road transport* show a strong decrease too (-88% and -100%, respectively), but these emissions represent about 12% and 0.4% of the total in 2020. *Production processes* and *other mobile sources and machinery* also present a significant decreasing trend, showing an influence on the total of 20% and 10% and dropping by about -88% and -91%, respectively. SO_x emissions from agriculture and from solvent and other product use are also estimated but their contribution is irrelevant.

An explanation of the sectoral decreasing trend is outlined more in details in the following.

Combustion in energy and transformation industries

The trend of emissions of this sector shows a reduction in the early eighties mainly due to the use of natural gas in place of coal in the energy production and to the implementation of the Directive EEC 75/716 (EC, 1975) which introduces more restrictive constraints in the sulphur content of liquid fuels.

During the years 1985-1990, there was an increase of energy consumption that, not sufficiently hampered by additional measures, led to an increase in the emissions of the sector and consequently of total SO_x levels.

However in the nineties, there was an inverse trend due to the introduction of two regulatory instruments: the DPR 203/88 (Decree of President of the Republic of 24th May 1988), laying down rules concerning the authorisation of plants, and the Ministerial Decree of 12th July 1990, which introduced plant emission limits. Also the European Directive 88/609/EEC (EC, 1988) concerning the limitation of specific pollutants originated from large combustion plants, transposed in Italy by the DM 8th May 1989 (Ministerial Decree of 8th May 1989) gave a contribution to the reduction of emissions in the sector.

Finally, in recent years, a further shift to natural gas in place of fuel oil and a further reduction in the use of coal fuels have contributed to a decrease in emissions.

Non industrial combustion plants

The declining of the emissions occurred mainly as a result of the increase in natural gas and LPG as alternative fuel to coal, diesel and fuel oil for heating; furthermore, several European Directives on the sulphur content in fuels were adopted. In accordance with national legislation, the sulphur content allowed in diesel fuel has decreased from 0.8% in 1980 to 0.2% in 1995 and 0.1% in 2008, while in fuel oil for heating from 3% in 1980 to 0.3% in 1998. Moreover, coal is not more allowed for residential and commercial heating.

Combustion in industry

Emissions from this sector show the same trend of reduction as the category previously analysed, as both in the scope of the same rules.

Production processes

Emissions from refineries have been reduced as a result of compliance with the DM 12th July 1990 (Ministerial Decree of 12th July 1990), which introduces limit values. The reduction of emissions from chemical industry is due to the drop off of the sulphuric acid production and to the decrease of emissions in the production of carbon black.

Road transport

The reduction of emissions is mainly due to the introduction of European Directives regulating the sulphur content in liquid fuels.

Other mobile sources and machinery

As regards off roads, emissions mainly derive from maritime transport, which show a decrease due to the introduction of European Directives regulating the sulphur content in fuels.

2.1.2 Nitrogen oxides (NO_x)

The national atmospheric emissions of nitrogen oxides show a decreasing trend in the period 1990-2020, from 2,125 Gg to 671 Gg. Figure 2.2 and Table 2.2 show emission figures from 1990 to 2020. Figure 2.2 also illustrates the share of NO_x emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.2 NO_x emission trend, percentage share by sector and variation 1990-2020

Table 2.2 *NO_x emission trend from 1990 to 2020 (Gg)*

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|----------------|----------------|----------------|----------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | Gg | | | | | | | | | |
| Combustion in energy and transformation industries | 457.4 | 344.3 | 172.6 | 117.9 | 81.3 | 52.4 | 45.6 | 41.6 | 38.7 | 34.0 |
| Non industrial combustion plants | 64.2 | 65.5 | 64.8 | 74.9 | 85.5 | 86.2 | 87.3 | 86.4 | 85.9 | 82.9 |
| Combustion - Industry | 250.6 | 182.4 | 154.0 | 155.5 | 99.7 | 60.2 | 54.4 | 53.3 | 52.2 | 45.3 |
| Production processes | 29.9 | 31.0 | 9.2 | 16.0 | 10.7 | 9.5 | 10.7 | 10.5 | 10.5 | 9.3 |
| Solvent and other product use | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Road transport | 996.1 | 1039.7 | 777.3 | 628.9 | 422.1 | 327.6 | 283.3 | 286.1 | 270.9 | 213.3 |
| Other mobile sources and machinery | 261.5 | 258.5 | 260.1 | 233.0 | 183.1 | 127.3 | 122.9 | 129.0 | 129.8 | 128.3 |
| Waste treatment and disposal | 2.9 | 3.1 | 2.6 | 2.9 | 2.6 | 2.4 | 2.4 | 2.3 | 2.3 | 2.4 |
| Agriculture | 61.7 | 64.1 | 63.3 | 59.6 | 49.5 | 49.9 | 51.6 | 49.5 | 48.9 | 55.0 |
| Total | 2,124.5 | 1,988.6 | 1,504.1 | 1,288.9 | 934.7 | 715.7 | 658.4 | 658.9 | 639.3 | 570.6 |

Total emissions show a reduction of about 73% from 1990 to 2020, with a marked decrease between 1995 and 2000, especially in the road transport and energy combustion sectors. The target value of emissions, fixed for 2010 by the National Emission Ceilings Directive (EC, 2001) at 990 Gg has been reached and continues to be respected. In 2015, in the framework of the UNECE/CLRTAP Convention, and in particular the Multieffects Protocol, a new target has been established for Italy equal to 60% of 2005 emissions in 2020 and it has been reached. Moreover, the revised National Emission Ceiling Directive (EU, 2016), established a target for Italy equal to 35% of 2005 emissions in 2030.

The main source of emissions is *road transport* (about 37% in 2020), which shows a reduction of 79% between 1990 and 2020; *other mobile sources and machinery* in 2020 contributes to the total emissions for 22% and have reduced by 51% from 1990. *Combustion in energy* and in *industry* shows a decrease of about 93% and 82%, respectively, having a share on the total of about 6% and 8% in 2020, respectively. Among the sectors concerned, the only one which highlights an increase in emissions is *non-industrial combustion plants* showing an increase by 29%, accounting for 15% of the total.

Details on the sectoral emission trend and respective variation are outlined in the following sections, starting from the early eighties.

Combustion in energy and transformation industries

Emissions from this sector show an upward trend until 1988 due to an increase in energy consumption, not prevented by reduction measures. From 1988 onwards, emissions present a gradual reduction due, mainly, to the introduction of the two regulatory instruments already mentioned for sulphur dioxide: the DPR 203/88 (Decree of President of the Republic of 24th May 1988), laying down rules for the authorization of facilities and the Ministerial Decree of 12th July 1990, which introduces plant emission limits. The adoption of these regulations, as the Ministerial Decree of 8th May 1989 on large combustion plants, has led to a shift in energy consumption from oil with high sulphur content to oil with lower sulphur content and to natural gas.

In recent years, the conversion to the use of natural gas to replace fuel oil has intensified, thanks to incentives granted for the improvement of energy efficiency. Furthermore, a significant reduction in the use of coal fuels for energy production has been recorded in the last years. These measures, together with those of promoting renewable energy and energy saving, have led to a further reduction of emissions in the sector.

In addition, in the last years, more stringent emission limits to the new plants have been established during the authorisation process with the aim to prevent air quality issues at local level.

Non industrial combustion plants

The increase in emissions is explained by the growing trend of energy consumption during the period considered. This is because in the last twenty years all the new buildings are equipped with heating system and old buildings have been modernized.

A national survey on energy consumption of households, conducted by the National Institute of Statistics (ISTAT, 2014), has supplied the amount of biomass burned to heating. Estimated values of biomass burnt are about 80% higher than previous estimates reported in the National Energy Balance (MSE, several years) and derived from regional or incomplete surveys. From 2013 this new biomass figures are reported in the National Energy Balance. In 2015 the reconstruction backwards of the time series has been finalised, with the collaboration of ISTAT and GSE (Energy Services Manager), and official data have been communicated to Eurostat. Furthermore, the continuous improvement of appliances for biomass combustion has led to a significant reduction in emissions for different pollutants (PM, COVNM, PAH) but on the other hand an increase of NO_x EFs because of the optimization of combustion process.

Combustion in industry

Emissions from this sector show a decreasing trend, motivated by the same reasons as the energy industry, having undergone the same legislation.

Road transport

The decrease is the result of two opposing trends: an increase in emissions in the early years of the historical series, with a peak in 1992, due to the increase in the fleet and in the total mileage of both passengers and goods transported by road, and a subsequent reduction in emissions. This decrease is, once more, the result of two opposing trends: on one hand, the growth of both the fleet and the mileage, on the other hand the introduction of technologies to reduce vehicle emissions, as the catalytic converter, provided by European Directives, in particular the Directives 91/441/EC (EC, 1991), 94/12/EC (EC, 1994) and 98/69/EC (EC, 1998) on light vehicles.

To encourage the reduction of emissions, different policies have also been implemented, including incentives to renew the public and private fleet and for the purchase of electric vehicles, promotion for the integrated expansion of rail, maritime and urban transport system, and programmes of sustainable mobility. In 2020 pandemic and relevant lockdown measures affected 2020 emission levels especially for this sector.

Other mobile sources and machinery

From 1980 emissions have a slightly rising trend until 1998 and then decrease slightly until arriving in 2017 at lower levels. Emissions in the sector are characterized predominantly by maritime transport, by machinery used in agriculture and industry.

Regarding mobile machinery used in agriculture and industry, these sectors were not governed by any legislation until the Directive 97/68/EC (EC, 1997 [a]), which provides for a reduction in NO_x limits from 1st January 1999, and Directive 2004/26/EC (EC, 2004) which provide further reduction stages with substantial effects from 2011, with a following decreasing trend particularly in recent years.

2.1.3 Ammonia (NH₃)

The national atmospheric emissions of ammonia show a slight decline in the period 1990-2020, from 467 Gg to 363 Gg. Figure 2.3 and Table 2.3 report the emission figures from 1990 to 2020. Figure 2.3 also illustrates the share of NH₃ emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.

According to the National Emission Ceilings Directive, the target value of emissions for 2010 amounts to 419 Gg which was achieved. The target established for 2020 in the framework of the UNECE/CLRTAP Convention and relevant protocol is equal for Italy to 95% of 2005 emissions and has been reached. Moreover, the revised national emission Ceiling Directive (EU, 2016) introduced a ceiling equal to 84% of 2005 emissions for 2030.




Figure 2.3 NH₃ emission trend, percentage share by sector and variation 1990-2020

Table 2.3 NH₃ emission trend from 1990 to 2020 (Gg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | Gg | | | | | | | | | |
| Combustion in energy and transformation industries | 0.3 | 0.2 | 0.2 | 0.3 | 0.3 | 0.3 | 0.2 | 0.2 | 0.1 | 0.1 |
| Non industrial combustion plants | 1.1 | 1.1 | 1.0 | 1.0 | 1.8 | 1.6 | 1.7 | 1.3 | 1.3 | 1.2 |
| Combustion - Industry | 0.5 | 0.6 | 0.6 | 4.0 | 1.6 | 1.0 | 1.2 | 1.1 | 1.1 | 1.0 |
| Production processes | 0.9 | 0.5 | 0.5 | 0.6 | 0.6 | 0.5 | 0.6 | 0.6 | 0.5 | 0.4 |
| Solvent and other product use | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 |
| Geothermal production | 8.4 | 9.0 | 12.3 | 13.3 | 6.0 | 4.1 | 5.5 | 2.9 | 1.1 | 1.1 |
| Road transport | 0.8 | 5.3 | 20.5 | 15.7 | 9.9 | 6.4 | 5.6 | 5.8 | 5.7 | 4.5 |
| Other mobile sources and machinery | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Waste treatment and disposal | 5.2 | 6.5 | 7.4 | 7.6 | 7.1 | 8.6 | 8.5 | 8.5 | 8.6 | 9.0 |
| Agriculture | 451.7 | 430.6 | 414.0 | 378.1 | 351.5 | 334.6 | 339.8 | 330.4 | 330.4 | 345.0 |
| Total | 469.2 | 454.2 | 457.0 | 421.0 | 379.1 | 357.4 | 363.6 | 351.1 | 349.2 | 362.6 |

In 2020 agriculture is the main source of emissions, with a 95% contribution out of the total NH₃ emissions; from 1990 to 2020 emissions from this sector show a decrease of about 24%. Emissions from road transport show a strong increase, but the share on the total is 1.2%. Emissions from waste treatment and disposal, accounting also only for 2.5% of the total, show an increase of about 71% because of the increase of NH₃ emissions from anaerobic digestion at biogas facilities. Emissions from non industrial combustion plants show a relevant increase, equal to 15%, but in 2020 the contribution to total emissions is 0.3%. Emissions from combustion in energy and transformation industries as emissions from combustion in industry are not relevant

accounting for 0.03% and 0.3% respectively. Emissions from *production processes* show a reduction of about 55%, but also this contribution is irrelevant as well as emissions from *solvent and other product use*. Finally, emissions from *geothermal production* contribute in 2020 for 0.3% of total national emissions.

Specifically, emissions from *agriculture* have decreased because of the reduction in the number of animals and the trend in agricultural production, and the introduction of abatement technologies due to the implementation of the EU IPPC Directive (EC, 1996). In the last years further emissions reduction result from the implementation of the European Union Rural Development Programs which provide incentives to the introduction of good practice and technologies for the environmental protection and mitigation of GHG and ammonia emissions. Emissions from *road transport* have increased as a result of the introduction of catalytic converter but during the last years a decrease is observed due to the introduction of more stringent limits in the new vehicles. Emissions from *geothermal production* have decreased because of the introduction of control and abatement systems in the production plants. *Waste sector* trend is driven by the increase of biogas facilities due to the incentives for energy production by renewable sources.

2.1.4 Non methane volatile organic compounds (NMVOC)

The national atmospheric emissions of NMVOC show a decreasing trend in the period 1990-2020. Figure 2.4 and Table 2.4 illustrate the emissions values from 1990 to 2020. Figure 2.4 also illustrates the share of NMVOC emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.4 NMVOC emission trend, percentage share by sector and variation 1990-2020

The total emission trend shows a reduction of about 56% between 1990 and 2020, from 1,993 Gg to 885 Gg.

In the framework of the National Emission Ceilings Directive (EC, 2001), the target value of NMVOC for 2010 fixed at 1,159 Gg was reached. The target established in the framework of the UNECE/CLRTAP Convention for 2020, equal to 65% of 2005 emission level, has been reached taking in account that it does not include emissions from the agriculture sector apart those from combustion of agriculture residues. In the framework of the European National Emission Ceiling Directive (EU, 2016) a target has been established for Italy equal to 54% of 2005 emissions in 2030.

Solvent and other product use is the main source of emissions, contributing to the total with 42% and showing a decrease of about 39%. The main reductions relate to the *road transport* sector (-88%), accounting for 10% of the total and to the sector of *extraction and distribution of fossil fuels/geothermal energy* (-66%),

accounting only for 3%. Emissions from *agriculture* decrease of about 16%, accounting for 14% of the national total. Emissions from *other mobile sources and machinery*, accounting for 3% of the total, decrease of about 83%. Emissions from *non industrial combustion plants* show the largest increase (60%), accounting for 19%. Emissions from *waste treatment and disposal* and *combustion in industry* show a decrease of about 3% and 14%, respectively, but both these sources account only for about 1%.

Details on the sectoral emission trend and respective variation are outlined in the following sections.

Table 2.4 NMVOC emission trend from 1990 to 2020 (Gg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|----------------|----------------|----------------|----------------|----------------|--------------|--------------|--------------|--------------|--------------|
| | <i>Gg</i> | | | | | | | | | |
| Combustion in energy and transformation industries | 7.6 | 7.4 | 6.1 | 5.6 | 4.9 | 3.7 | 4.0 | 3.7 | 3.7 | 3.4 |
| Non industrial combustion plants | 102.8 | 112.9 | 113.3 | 121.8 | 216.3 | 188.7 | 200.5 | 173.9 | 171.3 | 164.3 |
| Combustion - Industry | 7.2 | 8.0 | 8.4 | 8.4 | 6.7 | 6.7 | 6.8 | 6.8 | 6.7 | 6.2 |
| Production processes | 113.8 | 103.3 | 88.5 | 92.3 | 74.5 | 57.1 | 55.3 | 53.4 | 54.5 | 56.1 |
| Solvent and other product use | 90.9 | 103.7 | 56.6 | 53.9 | 48.6 | 37.7 | 38.5 | 35.5 | 31.7 | 30.5 |
| Geothermal production | 610.8 | 560.1 | 495.7 | 480.2 | 393.4 | 316.6 | 344.4 | 355.3 | 360.6 | 375.5 |
| Road transport | 767.1 | 878.7 | 609.2 | 359.3 | 184.7 | 127.0 | 113.5 | 108.2 | 103.1 | 91.7 |
| Other mobile sources and machinery | 133.4 | 122.0 | 97.6 | 73.7 | 51.0 | 29.0 | 24.9 | 24.6 | 23.4 | 22.4 |
| Waste treatment and disposal | 11.3 | 13.1 | 12.8 | 13.4 | 12.2 | 11.0 | 10.7 | 10.6 | 10.5 | 10.9 |
| Agriculture | 148.5 | 149.1 | 141.7 | 131.2 | 124.0 | 119.6 | 122.5 | 122.2 | 122.2 | 124.3 |
| Total | 1,993.4 | 2,058.3 | 1,629.9 | 1,339.8 | 1,116.1 | 897.0 | 921.1 | 894.4 | 887.7 | 885.4 |

Solvent and other product use

Emissions from this sector stem from numerous activities such as painting (both domestic and industrial), degreasing and dry cleaning, manufacturing and processing of chemicals, other use of solvents and related activities including the use of household products that contain solvents, such as cosmetics, household products and toiletries.

Significant reductions occurred in the nineties by the introduction in the market of products with low solvent content in paints, and the reduction of the total amount of organic solvent used for metal degreasing and in glues and adhesives; furthermore, in many cases, local authorities have imposed abatement equipment in the industrial painting sector and forced the replacement of open loop with closed loop laundry machines even before the EU Directive 99/13/EC (EC, 1999) came into force. In 2020 due to pandemic a strong increase of household products have been registered.

Road transport

The trend of emissions in this sector is characterized by a first stage of reduction in the early eighties, which occurred despite the increase of consumption and mileage because of the gradual adjustment of the national fleet to the European legislation, ECE Regulation 15 and subsequent amendments, introducing stricter emission limits for passenger cars. Subsequently, in the early nineties, an increase in emissions is observed, with a peak in 1992, due to a high increase in gasoline consumption not efficiently opposed by the replacement of the fleet. With the introduction of Directive 91/441/EC (EC, 1991) and following legislation, which provide the use of catalytic device to reduce exhaust and evaporative emissions from cars, NMVOC emissions gradually reduced.

A different explanation of the emission trend pertains to the nineties. In fact, in this period an increase of the fleet and the mileage is observed in Italy, especially for the emergent use of mopeds for urban mobility, which, until 1999, were not subject to any national emission regulation. Thereafter, various measures were introduced in order to facilitate the reduction of NMVOC emissions, including incentives for replacement of both the fleet of passenger cars and of mopeds and motorcycles with low-emission vehicles; incentives were also provided for the use of fuels different from gasoline, such as LPG and natural gas. In addition, funds were allocated for the implementation of urban traffic plans, for the establishment of restricted traffic areas and car-free days, for checks on exhaust pipes of cars, for the implementation of voluntary agreements with manufacturers of mopeds and motorcycles in order to anticipate the timing provided by the European Directive 97/24/EC (EC, 1997 [b]) as regards the placing on the market of mopeds with low emissions.

Non industrial combustion plants

The increasing emission trend is driven by the increase of wood biomass fuel consumption for residential heating. The 2013 consumption value reported in the national energy balance results from a detailed survey conducted by the national institute of statistics in 2014 (ISTAT, 2014) and is much higher than previous estimates. In 2015 the reconstruction backwards of the time series of wood combustion has been finalised, with the collaboration of ISTAT and GSE (Energy Services Manager), and official data have been communicated to Eurostat.

Other mobile sources and machinery

The reduction in emissions is explained by the reduction of gasoline consumption in the sector, largely for two-stroke engines used in agriculture and in maritime activities.

Agriculture

NMVOC emissions from agriculture, mainly depend on activity data about different livestock categories. These emissions became significant because of the implementation of the 2016 Guidebook EMEP/EEA emission factors. For the compliance with the established targets these emissions could be subtracted by the total according to the National emission Ceiling Directive (EU, 2016) due to their uncertainty.

As regards the other sectors, a decrease in emissions from production processes is observed, mainly in the food industries, in the chemical sector and in the processes in the refineries. The emissions concerning the extraction and distribution of fuels, even in the presence of an increase in quantity treated, have been reduced as a result of the application of the DM 16th May 1996 (Ministerial Decree 16 May 1996), concerning the adoption of devices for the recovery of vapours and of the applications of measures on deposits of gasoline provided by the DM 21st January 2000 (Ministerial Decree 21 January 2000).

Emissions from the other sectors are not subject to specific regulations.

2.1.5 Carbon monoxide (CO)

The national CO emissions show a decreasing trend in the period 1990-2020, from 6,797 Gg to 1,873 Gg. The emission figures from 1990 to 2020 are shown in Figure 2.5 and Table 2.5. Figure 2.5 also illustrates the share of CO emissions by category in 1990 and 2020, as well as the total and sectoral variation from 1990 to 2020.





Figure 2.5 CO emission trend, percentage share by sector and variation 1990-2020

Table 2.5 CO emission trend from 1990 to 2020 (Gg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| | Gg | | | | | | | | | |
| Combustion in energy and transformation industries | 58.9 | 54.1 | 54.4 | 53.9 | 34.5 | 39.9 | 44.5 | 39.6 | 38.6 | 39.0 |
| Non industrial combustion plants | 795.1 | 894.0 | 913.1 | 930.4 | 1,664.9 | 1,395.4 | 1,475.5 | 1,289.1 | 1,267.7 | 1,204.6 |
| Combustion - Industry | 305.6 | 410.9 | 314.6 | 326.0 | 233.6 | 92.8 | 81.9 | 80.8 | 112.2 | 87.9 |
| Production processes | 223.7 | 139.8 | 129.2 | 143.6 | 105.0 | 63.6 | 71.8 | 70.8 | 69.6 | 60.0 |
| Solvent and other product use | 5.1 | 5.1 | 5.7 | 5.3 | 5.1 | 4.4 | 4.3 | 4.2 | 4.0 | 3.9 |
| Road transport | 4,874.5 | 5,106.1 | 2,973.8 | 1,681.2 | 776.3 | 479.3 | 403.6 | 386.5 | 389.6 | 294.0 |
| Other mobile sources and machinery | 480.5 | 402.5 | 302.9 | 263.4 | 193.9 | 132.0 | 118.7 | 122.9 | 123.7 | 126.6 |
| Waste treatment and disposal | 40.7 | 46.9 | 45.4 | 50.5 | 47.2 | 47.0 | 46.0 | 44.3 | 44.0 | 44.9 |
| Agriculture | 12.5 | 12.2 | 12.1 | 13.1 | 12.5 | 12.7 | 12.4 | 12.2 | 12.0 | 12.1 |
| Total | 6,796.5 | 7,071.6 | 4,751.1 | 3,467.4 | 3,073.0 | 2,267.2 | 2,258.6 | 2,050.5 | 2,061.5 | 1,872.8 |

The decrease in emissions (-72%) is mostly due to the trend observed for the transport sector (including road, railways, air and maritime transport) which shows a total reduction from 1990 to 2020 of about 92%. Specifically, by sector, emissions from *road transport* and *other mobile sources and machinery*, accounting in 2020 respectively for 16% and 7% of the total, show a decrease from 1990 to 2020 of about 94% and 74% respectively. On the other hand, emissions from *non industrial combustion plants*, representing about 64% of the total in 2020, show a strong increase between 1990 and 2020, equal to 51% due to the increase of wood combustion for residential heating.

Figures show an increase in emissions from *waste treatment and disposal* too (10%), whose share is 2% of the total and a slight decrease (-3%) for *agriculture* which accounts for less than 1% of the total.

2.2 PARTICULATE MATTER

2.2.1 PM10

The national atmospheric emissions of PM10 show a decreasing trend in the period 1990-2020, from 302 Gg to 166 Gg. Figure 2.6 and Table 2.6 illustrate the emission trend from 1990 to 2020. Figure 2.6 also illustrates the share of PM10 emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.6 PM10 emission trend, percentage share by sector and variation 1990-2020

Table 2.6 PM10 emission trend from 1990 to 2020 (Gg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|------|------|------|------|-------|-------|-------|------|------|------|
| | Gg | | | | | | | | | |
| Combustion in energy and transformation industries | 44.8 | 39.6 | 18.4 | 5.9 | 2.8 | 1.2 | 0.9 | 0.8 | 0.7 | 0.6 |
| Non industrial combustion plants | 67.8 | 71.2 | 68.6 | 68.6 | 123.1 | 106.8 | 113.0 | 95.1 | 94.0 | 89.9 |
| Combustion - Industry | 27.6 | 25.1 | 18.6 | 17.9 | 12.4 | 7.7 | 7.7 | 7.9 | 7.3 | 6.7 |
| Production processes | 30.1 | 29.1 | 26.0 | 27.6 | 20.3 | 13.7 | 13.2 | 13.6 | 13.6 | 12.3 |
| Extraction and distribution of fossil fuels | 0.7 | 0.6 | 0.6 | 0.8 | 0.7 | 0.6 | 0.5 | 0.4 | 0.3 | 0.2 |
| Solvent and other product use | 2.8 | 2.8 | 3.8 | 3.8 | 3.4 | 2.5 | 2.3 | 2.3 | 2.2 | 2.0 |
| Road transport | 58.3 | 57.1 | 52.1 | 46.1 | 33.0 | 23.8 | 20.3 | 19.8 | 19.4 | 15.5 |
| Other mobile sources and machinery | 31.6 | 32.1 | 30.5 | 25.1 | 15.9 | 9.7 | 8.9 | 9.0 | 8.8 | 8.8 |
| Waste treatment and disposal | 5.4 | 5.6 | 5.5 | 5.8 | 5.3 | 5.8 | 6.5 | 6.3 | 6.5 | 6.5 |
| Agriculture | 33.5 | 34.2 | 33.0 | 30.2 | 22.9 | 23.1 | 23.2 | 23.2 | 23.2 | 23.2 |

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | <i>Gg</i> | | | | | | | | | |
| Total | 302.5 | 297.4 | 257.1 | 231.5 | 239.8 | 195.0 | 196.5 | 178.4 | 176.1 | 165.7 |

From 1990 to 2020 the trend shows a reduction of about 45%. A considerable amount of emissions is mostly to be attributed to *non industrial combustion plant* (54% in 2020) which is increasing its emissions, about 33%, due to the increase of wood combustion for residential heating.

Agriculture sector, accounting for 14% of total emissions in 2020, reduced its emissions by 31% in 2020 respect to 1990, due to a reduction in emissions from livestock, which fall by about 42%, and a reduction in emissions from crops, which fall by about 19% due to a reduction in the area of arable land in 2020 compared to 1990.

Road transport accounts for 12% of total emissions in 2020 and decrease by 74% due to the introduction of the relevant European Directives controlling and limiting PM emissions at the car exhaust pipe.

In 2020 *other mobile sources and machinery*, accounting for 5% of the total, shows a reduction of about 72% in consideration of the implementation of the relevant European Directives on machinery. Emissions from *combustion in industry* account for about 4% of the total and decrease by about 76%. Emissions from *production processes* accounting for 7% of the total in 2020 decrease of about 59% between 1990 and 2020. The largest decrease (-99%) is observed in emissions deriving from *combustion in energy and transformation industries*, whose contribution to total emissions is almost irrelevant in 2020 and lower than 1%. The reduction in the energy and industrial sectors is mainly due to the introduction of two regulatory instruments, already mentioned for other pollutants, the DPR 203/88 (Decree of President of the Republic of 24th May 1988), laying down rules for the authorization of facilities and the Ministerial Decree of 12th July 1990, which introduces plant emission limits.

2.2.2 PM2.5

The trend of the national atmospheric emissions of PM2.5 is decreasing between 1990 and 2020, with a variation from 230 Gg to 133 Gg. Figure 2.7 and Table 2.7 illustrate the emission trend from 1990 to 2020. Figure 2.7 also illustrates the share of PM2.5 emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.7 PM2.5 emission trend, percentage share by sector and variation 1990-2020

Table 2.7 *PM2.5 emission trend from 1990 to 2020 (Gg)*

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | <i>Gg</i> | | | | | | | | | |
| Combustion in energy and transformation industries | 30.1 | 27.8 | 12.7 | 3.7 | 1.8 | 0.8 | 0.6 | 0.6 | 0.5 | 0.4 |
| Non industrial combustion plants | 66.9 | 70.6 | 67.9 | 67.9 | 121.8 | 105.6 | 111.7 | 93.9 | 92.8 | 88.8 |
| Combustion - Industry | 19.9 | 18.3 | 14.0 | 13.6 | 9.8 | 6.3 | 6.4 | 6.5 | 6.1 | 5.6 |
| Production processes | 14.2 | 13.6 | 11.5 | 12.1 | 9.7 | 6.9 | 6.3 | 6.4 | 6.3 | 5.6 |
| Extraction and distribution of fossil fuels | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 |
| Solvent and other product use | 2.6 | 2.6 | 3.3 | 3.2 | 2.9 | 2.3 | 2.1 | 2.1 | 2.0 | 1.8 |
| Road transport | 52.9 | 51.0 | 45.5 | 39.0 | 26.5 | 17.5 | 14.7 | 13.9 | 13.5 | 10.6 |
| Other mobile sources and machinery | 31.5 | 32.0 | 30.4 | 25.0 | 15.9 | 9.7 | 8.8 | 8.9 | 8.8 | 8.8 |
| Waste treatment and disposal | 5.0 | 5.2 | 5.2 | 5.4 | 4.9 | 5.5 | 6.1 | 6.0 | 2.5 | 6.2 |
| Agriculture | 7.1 | 7.0 | 6.9 | 6.5 | 5.3 | 5.4 | 5.4 | 5.3 | 5.3 | 5.3 |
| Total | 230.4 | 228.2 | 197.4 | 176.4 | 198.6 | 159.9 | 162.2 | 143.7 | 137.8 | 133.2 |

In the framework of the revision of the Multieffect protocol of the UNECE/CLRTAP Convention, a target has been established for this pollutant. Italy should reduce in 2020 their PM2.5 emissions by 10% with respect the 2005 emission level and it has been reached. Moreover, in the national emission Ceiling Directive a target has been established for 2030 equal to 60% of 2005 emissions.

Total emissions show a global reduction from 1990 to 2020 of about 42%. Specifically, emissions from *road transport*, accounting for 8% of total emissions, decrease of about 80%. Emissions from *other mobile sources and machinery* show a reduction of 72%, accounting in 2020 for 7% of total emissions. Emissions from *non industrial combustion plants* and from *combustion in industry* account for 67% and 4% of the total respectively, but while the former shows an increase of about 33%, the latter decreases by about 72%.

Agriculture sector, accounting for 4% of total emissions in 2020, reduced its emissions by 24% in 2020 respect to 1990. Emissions from *waste treatment and disposal*, accounting for 5% of the total in 2020, show an increase of about 24%. The largest decrease is observed for *combustion in energy and transformation industries* (-99%), being the influence on the total in 2020 lower than 1%.

For the explanation of the trends see what already reported for PM10.

2.2.3 Black Carbon (BC)

Black Carbon emissions have been estimated as a fraction of PM_{2.5}. National BC atmospheric emissions are decreasing between 1990 and 2020, with a variation from 48 Gg to 16 Gg. Figure 2.7 and Table 2.7 illustrate the emission trend from 1990 to 2020. Figure 2.7 also illustrates the share of BC emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.8 BC emission trend, percentage share by sector and variation 1990-2020

Table 2.8 BC emission trend from 1990 to 2020 (Gg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | Gg | | | | | | | | | |
| Combustion in energy and transformation industries | 1.66 | 1.57 | 0.78 | 0.30 | 0.11 | 0.04 | 0.04 | 0.03 | 0.03 | 0.03 |
| Non industrial combustion plants | 5.39 | 5.66 | 5.56 | 5.73 | 10.33 | 9.08 | 9.60 | 8.43 | 8.37 | 8.03 |
| Combustion - Industry | 0.73 | 0.62 | 0.48 | 0.54 | 0.43 | 0.27 | 0.28 | 0.29 | 0.27 | 0.25 |
| Production processes | 0.49 | 0.42 | 0.28 | 0.29 | 0.23 | 0.14 | 0.12 | 0.12 | 0.12 | 0.10 |
| Extraction and distribution of fossil fuels | 0.06 | 0.05 | 0.05 | 0.06 | 0.06 | 0.05 | 0.04 | 0.04 | 0.03 | 0.02 |
| Road transport | 23.99 | 22.21 | 20.66 | 19.56 | 13.79 | 7.94 | 6.27 | 5.46 | 5.11 | 3.82 |
| Other mobile sources and machinery | 14.10 | 14.50 | 13.41 | 10.61 | 5.81 | 3.39 | 2.89 | 2.81 | 2.58 | 2.44 |
| Waste treatment and disposal | 1.37 | 1.46 | 1.44 | 1.53 | 1.42 | 1.52 | 1.63 | 1.59 | 0.96 | 0.98 |
| Agriculture | 0.12 | 0.12 | 0.11 | 0.12 | 0.11 | 0.12 | 0.11 | 0.11 | 0.11 | 0.11 |
| Total | 47.91 | 46.60 | 42.77 | 38.74 | 32.30 | 22.56 | 20.99 | 18.89 | 17.58 | 15.79 |

Total emissions show a global reduction from 1990 to 2020 of about 67%. Specifically, emissions from *road transport*, accounting for 24% of total emissions, decrease of about 84%. Emissions from *other mobile sources and machinery* show a reduction of 83%, accounting in 2020 for 15% of total emissions. Emissions from *non industrial combustion plants* and from *combustion in industry* account for 51% and 2% of the total respectively, but while the former shows an increase of about 49%, the latter decreases by about 66%.

Industrial processes, accounting for less than 1% in 2020, decrease of 79%. Emissions from *waste treatment and disposal*, accounting for 6% of the total in 2020, show a decrease of about 29%. The largest decrease is observed for *combustion in energy and transformation industries* (-98%), being the influence on the total in 2020 less than 1%.

For the explanation of the trends refer to previous paragraph.

2.3 HEAVY METALS (Pb, Cd, Hg)

This section provides an illustration of the most significant developments between 1990 and 2020 of lead, cadmium and mercury emissions.

2.3.1 Lead (Pb)

The national atmospheric emissions of lead show a strong decreasing trend (-96%) between 1990 and 2020, varying from 4,278 Mg to 156 Mg. Figure 2.9 and Table 2.9 illustrate the emission trend from 1990 to 2020. Figure 2.9 also illustrates the share of Pb emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.9 Pb emission trend, percentage share by sector and variation 1990-2020

Table 2.9 Pb emission trend from 1990 to 2020 (Mg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|---------|---------|-------|-------|-------|------|------|------|------|------|
| | | | | Mg | | | | | | |
| Combustion in energy and transformation industries | 4.0 | 4.0 | 3.6 | 3.8 | 3.1 | 2.7 | 2.3 | 2.0 | 1.7 | 1.4 |
| Non-industrial combustion plants | 14.5 | 16.6 | 22.4 | 46.3 | 16.5 | 15.1 | 15.2 | 14.4 | 14.1 | 13.6 |
| Combustion - industry | 263.1 | 234.9 | 153.6 | 141.9 | 104.7 | 95.2 | 79.4 | 78.9 | 77.2 | 67.7 |
| Production processes | 63.7 | 68.2 | 67.3 | 74.2 | 69.5 | 66.1 | 69.6 | 72.0 | 68.1 | 61.4 |
| Solvent and other product use | 2.4 | 2.4 | 8.4 | 10.3 | 8.4 | 4.3 | 3.4 | 3.4 | 3.4 | 2.1 |
| Road transport | 3,782.3 | 1,617.6 | 690.0 | 13.2 | 12.3 | 12.2 | 10.1 | 10.5 | 10.5 | 8.6 |
| Other mobile sources and machinery | 142.2 | 44.2 | 13.3 | 1.1 | 1.1 | 1.1 | 1.1 | 1.2 | 1.3 | 0.7 |
| Waste treatment and disposal | 5.9 | 5.5 | 2.8 | 4.1 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 |

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--------------|----------------|----------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | | | | Mg | | | | | | |
| Agriculture | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 |
| Total | 4,278.2 | 1,993.5 | 961.4 | 294.9 | 215.8 | 197.0 | 181.4 | 182.5 | 176.5 | 155.8 |

In 2020 emissions from *combustion in industry* have the most significant impact on the total (43%) and show a reduction of about 74%; this reduction is to be attributed primarily to *processes with contact*, which contribute with 81% to the sectoral reduction and account for almost the total share of the sector. Emissions from *production processes* and, in particular, from processes in iron and steel industries and collieries, decreased by about 4%, and represent 39% of the total. Emissions from *non industrial combustion plants* show a 6% decrease and represent, in 2020, 9% of the total. As to emissions from *transport* activities, because of changes occurred in the legislation regarding fuels, trends show a sharp reduction in emissions from 2002 onwards.

2.3.2 Cadmium (Cd)

The national atmospheric emissions of cadmium show a decreasing trend. Figure 2.10 and Table 2.10 illustrate the emission trend from 1990 to 2020. Figure 2.10 also illustrates the share of Cd emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.10 Cd emission trend, percentage share by sector and variation 1990-2020

Table 2.10 Cd emission trend from 1990 to 2020 (Mg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|--------------|--------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | <i>Mg</i> | | | | | | | | | |
| Combustion in energy and transformation industries | 0.19 | 0.20 | 0.17 | 0.16 | 0.12 | 0.10 | 0.09 | 0.08 | 0.07 | 0.06 |
| Non-industrial combustion plants | 1.51 | 1.20 | 1.73 | 2.61 | 0.66 | 0.55 | 0.55 | 0.50 | 0.48 | 0.46 |
| Combustion - industry | 5.61 | 5.56 | 4.99 | 2.47 | 1.56 | 1.15 | 1.23 | 1.24 | 1.24 | 1.16 |
| Production processes | 2.01 | 1.78 | 1.42 | 1.52 | 1.35 | 1.14 | 1.14 | 1.16 | 1.10 | 0.98 |
| Solvent and other product use | 0.50 | 0.50 | 0.56 | 0.52 | 0.49 | 0.41 | 0.38 | 0.37 | 0.36 | 0.34 |
| Road transport | 0.41 | 0.49 | 0.52 | 0.51 | 0.46 | 0.43 | 0.41 | 0.41 | 0.41 | 0.33 |
| Other mobile sources and machinery | 0.03 | 0.03 | 0.03 | 0.03 | 0.03 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 |
| Waste treatment and disposal | 0.61 | 0.60 | 0.49 | 0.52 | 0.35 | 0.38 | 0.36 | 0.37 | 0.35 | 0.36 |
| Agriculture | 0.13 | 0.13 | 0.12 | 0.13 | 0.12 | 0.13 | 0.12 | 0.12 | 0.12 | 0.12 |
| Total | 10.98 | 10.47 | 10.04 | 8.48 | 5.14 | 4.30 | 4.29 | 4.27 | 4.14 | 3.83 |

Emissions show a global reduction of 62% between 1990 and 2020, from 11.0 Mg to 3.8 Mg, mainly driven by the reduction of emissions in the non ferrous metal industry, with the installation of the relevant abatement technologies and the drop of production. Among the most significant variations, emissions from *combustion in industry* and from *non industrial combustion plants* represent 30% and 12% of the total respectively, showing a decrease of 79% and 69% respectively. Emissions from *production processes* decrease by about 51% and represent 26% of the total. Emissions from *waste treatment and disposal* (i.e. waste incineration), accounting for 9% of the total, register a reduction of about 41% while emissions from *road transport*, accounting for 9% of the total levels, decreased by 19% and emissions from *stubble burning in agriculture* account for 3% of the total and decrease of about 9%.

2.3.3 Mercury (Hg)

The national atmospheric emissions of mercury show a reduction trend in the period 1990-2020. Figure 2.11 and Table 2.11 illustrate the emission trend from 1990 to 2020. Figure 2.11 also illustrates the share of Hg emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.

Emission trend shows a global reduction of about 62% from 1990 to 2020, varying from 15.3 Mg to 5.8 Mg. The general trend is driven by reduction of emissions in lead and zinc production industry as well as in cement production industry, with the installation of the relevant abatement technologies. The main variations concern: emissions from *combustion in industry - processes with contact*, accounting for 26% and decreasing by 58%; emissions from *production process - processes in iron and steel industries and collieries*, representing 45% of the total and increasing by 4%; emissions from *non industrial combustion plants* which represent 8% of the total and decrease by 23%. Emissions deriving from *combustion in energy and transformation industries*, accounting for 8%, show a 58% reduction. Emissions from *production process - processes in inorganic chemical industries*, not contributing to the total in 2020, show a reduction equal to 100% totally due to the technological changes for the production of chlorine. Emissions from *road transport* account for 3% and decrease of 23%. Emissions from *waste treatment and disposal* contributing to the total for 2% and *agriculture*, contributing to the total only for less than 1%, show a large reduction, equal respectively to 70% and 9%. Emissions from *geothermal production* account for 6% of the national total and shows a reduction of 90% with respect to 1990 due to the introduction of control and abatement system at the production plants.




Figure 2.11 Hg emission trend, percentage share by sector and variation 1990-2020

Table 2.11 Hg emission trend from 1990 to 2020 (Mg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|--------------|--------------|--------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | Mg | | | | | | | | | |
| Combustion in energy and transformation industries | 1.10 | 1.14 | 1.06 | 1.07 | 0.86 | 0.69 | 0.64 | 0.58 | 0.52 | 0.47 |
| Non-industrial combustion plants | 0.61 | 0.71 | 1.04 | 1.97 | 0.52 | 0.49 | 0.49 | 0.49 | 0.48 | 0.47 |
| Combustion - industry | 4.17 | 3.95 | 3.43 | 3.41 | 2.55 | 2.06 | 1.78 | 1.74 | 1.74 | 1.64 |
| Production processes | 5.47 | 4.36 | 3.59 | 3.36 | 2.87 | 2.70 | 3.00 | 3.07 | 2.91 | 2.60 |
| Geothermal production | 3.40 | 3.62 | 4.96 | 2.15 | 1.25 | 0.98 | 0.85 | 0.72 | 0.43 | 0.34 |
| Road transport | 0.19 | 0.23 | 0.24 | 0.24 | 0.21 | 0.19 | 0.17 | 0.18 | 0.18 | 0.15 |
| Waste treatment and disposal | 0.35 | 0.34 | 0.25 | 0.32 | 0.23 | 0.11 | 0.11 | 0.11 | 0.10 | 0.10 |
| Agriculture | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 |
| Total | 15.30 | 14.36 | 14.59 | 12.53 | 8.51 | 7.25 | 7.06 | 6.90 | 6.38 | 5.78 |

2.4 PERSISTENT ORGANIC POLLUTANTS (POPs)

In this section, the most significant peculiarities of polycyclic aromatic hydrocarbons and dioxins, occurred between 1990 and 2020, will be presented.

2.4.1 Polycyclic aromatic hydrocarbons (PAH)

The national atmospheric emissions of polycyclic aromatic hydrocarbons decreased from 90 Mg to 60 Mg between 1990 and 2020. Figure 2.12 and Table 2.12 illustrate the emission trend from 1990 to 2020. Figure 2.12 also illustrates the share of PAH emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.12 PAH emission trend, percentage share by sector and variation 1990-2020

Table 2.12 PAH emission trend from 1990 to 2020 (Mg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|------|------|------|------|------|------|------|------|------|------|
| | Mg | | | | | | | | | |
| Combustion in energy and transformation industries | 0.9 | 1.0 | 0.7 | 0.5 | 0.4 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 |
| Non-industrial combustion plants | 31.9 | 35.2 | 35.7 | 38.9 | 67.3 | 55.9 | 59.2 | 52.2 | 51.2 | 48.6 |
| Combustion - industry | 4.5 | 4.6 | 2.2 | 2.4 | 0.4 | 0.5 | 0.5 | 0.5 | 0.5 | 0.4 |
| Production processes | 48.4 | 47.4 | 16.8 | 17.7 | 14.1 | 9.3 | 9.6 | 9.4 | 9.0 | 7.1 |
| Solvent and other product use | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Road transport | 1.9 | 1.9 | 2.1 | 2.6 | 2.6 | 2.5 | 2.5 | 2.6 | 2.6 | 2.1 |
| Other mobile sources and machinery | 0.3 | 0.3 | 0.3 | 0.4 | 0.3 | 0.2 | 0.2 | 0.3 | 0.3 | 0.3 |
| Waste treatment and disposal | 1.3 | 1.3 | 1.2 | 1.2 | 1.0 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 |
| Agriculture | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 | 0.4 |

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | <i>Mg</i> | | | | | | | | | |
| Total | 89.7 | 92.2 | 59.5 | 64.0 | 86.5 | 70.5 | 74.0 | 66.8 | 65.5 | 60.4 |

Between 1990 and 2020, total emissions show a decrease of about 33%. Among the most significant changes, *non industrial combustion plants*, prevalently *residential plants*, account for 80% of the total in 2020 and show a strong increase (about 52%) due to the increase in wood consumption for heating.

Emissions from *production processes*, mainly *processes in iron and steel industries*, account for 12% of the total and show a decrease of 85% due to the adoption of best abatement technologies for the coke production; emissions from *waste treatment and disposal*, mainly open burning of agricultural wastes except stubble burning, account for 2% of the total and show a decrease of 16%. Emissions from *road transport*, accounting for 3% in 2020, show an increase of about 7%. The share of other subsectors is lower than 1%.

2.4.2 Dioxins

The national atmospheric emissions of dioxins show a decreasing trend between 1990 and 2020, with values varying from 529 g I Teq to 280 g I Teq. Figure 2.13 and Table 2.13 illustrate the emission trend from 1990 to 2020. Figure 2.13 also illustrates the share of dioxin emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.13 Dioxin emission trend, percentage share by sector and variation 1990-2020

Table 2.13 Dioxin emission trend from 1990 to 2020 (g I Teq)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|----------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | <i>g I Teq</i> | | | | | | | | | |
| Combustion in energy and transformation industries | 24.9 | 27.9 | 20.9 | 14.0 | 8.2 | 6.2 | 5.3 | 4.7 | 4.2 | 4.0 |
| Non-industrial combustion plants | 173.8 | 164.6 | 151.0 | 87.0 | 134.6 | 112.4 | 118.4 | 103.7 | 101.6 | 96.4 |
| Combustion - industry | 116.7 | 120.9 | 112.0 | 118.7 | 65.3 | 58.2 | 59.8 | 55.9 | 57.0 | 47.1 |
| Production processes | 67.2 | 71.7 | 70.7 | 78.6 | 76.2 | 76.8 | 86.0 | 88.9 | 84.5 | 75.7 |
| Solvent and other product use | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| Road transport | 10.7 | 13.6 | 18.2 | 21.4 | 20.2 | 13.6 | 11.5 | 10.0 | 9.2 | 7.1 |
| Other mobile sources and machinery | 1.3 | 1.4 | 1.3 | 1.5 | 1.2 | 0.9 | 1.0 | 1.1 | 1.0 | 1.0 |
| Waste treatment and disposal | 134.3 | 110.9 | 60.3 | 40.2 | 35.9 | 41.3 | 48.3 | 47.0 | 49.0 | 48.8 |
| Agriculture | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Total | 529.0 | 511.2 | 434.5 | 361.5 | 341.6 | 309.6 | 330.4 | 311.4 | 306.7 | 280.2 |

The general trend shows a decrease from 1990 to 2020 equal to 47%, with a noticeable decline between 1995 and 2004 and between 2008 and 2011 because of the implementation of abatement system in the largest Italian integrated iron and steel plant (steel production > 80% with respect to national production from integrated plants):

- Double filtering system ESP (Electrostatic Precipitator) + MEEP (Moving Electrode Electrostatic Precipitator);
- Reduction of the chlorine amount in the charge;
- Injections of urea (able to form stable compounds with metals that catalyse the formation of dioxins).

The most considerable reductions, between 1990 and 2020, are observed in *waste treatment and disposal*, *combustion in energy and transformation industries* and *combustion in industry*, (-64%, -84% and -60%, respectively). Specifically, the reduction is principally due to the cut of emissions from the combustion of municipal waste both with energy recovery, reported under the non-industrial sector, and without recovery, reported under the waste sector due to the introduction of regulations establishing more stringent limits of dioxin emissions from stacks. The waste sector includes *accidental fires of vehicles and buildings* which account in 2020 for 15% of total national emissions.

In 2020, the subsectors which have contributed most to total emissions are *non-industrial combustion plants*, *production processes* and *combustion in industry* accounting for 34%, 27% and 17% of the total respectively. In particular emissions from *production processes* show an increase of 13% in the period 1990-2020 due to the increase of the iron and steel production in electric arc furnaces.

2.4.3 Hexachlorobenzene (HCB)

The national atmospheric emissions of hexachlorobenzene show a decreasing trend in the period 1990-2020, varying from 142 kg to 13 kg due to the decrease of the use of pesticide in agriculture. Figure 2.14 and Table 2.14 illustrate the emission trend from 1990 to 2020. Figure 2.14 also illustrates the share of HCB emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.14 HCB emission trend, percentage share by sector and variation 1990-2020

Table 2.14 HCB emission trend from 1990 to 2020 (Mg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | <i>Mg</i> | | | | | | | | | |
| Combustion in energy and transformation industries | 0.000 | 0.000 | 0.000 | 0.000 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 |
| Non-industrial combustion plants | 0.002 | 0.003 | 0.006 | 0.006 | 0.004 | 0.003 | 0.004 | 0.003 | 0.003 | 0.003 |
| Combustion - industry | 0.008 | 0.008 | 0.009 | 0.009 | 0.007 | 0.007 | 0.007 | 0.006 | 0.007 | 0.005 |
| Road transport | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| Other mobile sources and machinery | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 |
| Waste treatment and disposal | 0.013 | 0.014 | 0.010 | 0.008 | 0.000 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 |
| Agriculture | 0.119 | 0.084 | 0.007 | 0.002 | 0.002 | 0.003 | 0.003 | 0.002 | 0.002 | 0.002 |
| Total | 0.142 | 0.110 | 0.033 | 0.027 | 0.016 | 0.016 | 0.016 | 0.015 | 0.014 | 0.013 |

The use of pesticide in *agriculture* category is the main driver for the decreasing trend of the HCB national emissions, emissions from this category show 98% decrease between 1990 and 2020. The second sector contributing to the general trend is *waste treatment and disposal*, in particular waste incineration - sludge incineration. Specifically, the considerable increase of the amount of sludge burnt at a specific incinerator is the reason of the peaks observed in 2001-2003 (incineration with energy recovery). The other relevant sectors are *combustion in industry* and *non industrial combustion plants* accounting for 23% and 42% respectively. Emissions from *combustion in energy and transformation industry* and emissions from *non industrial combustion plants* show an increase of 216% and 56% respectively between 1990 and 2020. In the same years for emissions from *waste treatment and disposal* a decrease of 94% must be noted while emissions from combustion in industry show a decrease of 30%.

2.4.4 Polychlorinated biphenyl (PCB)

The national atmospheric emissions of polychlorinated biphenyl show a decreasing trend in the period 1990-2020, about -33%, from 154 kg to 103 kg. Figure 2.15 and Table 2.15 illustrate the emission trend from 1990 to 2020. Figure 2.15 also illustrates the share of PCB emissions by category in 1990 and 2020 as well as the total and sectoral variation from 1990 to 2020.




Figure 2.15 PCB emission trend, percentage share by sector and variation 1990-2020

Table 2.15 PCB emission trend from 1990 to 2020 (Mg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | <i>Mg</i> | | | | | | | | | |
| Combustion in energy and transformation industries | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| Non-industrial combustion plants | 0.015 | 0.018 | 0.025 | 0.036 | 0.022 | 0.019 | 0.020 | 0.019 | 0.018 | 0.018 |
| Combustion - industry | 0.042 | 0.043 | 0.033 | 0.035 | 0.017 | 0.014 | 0.014 | 0.013 | 0.013 | 0.010 |
| Production processes | 0.092 | 0.100 | 0.096 | 0.106 | 0.093 | 0.079 | 0.087 | 0.088 | 0.083 | 0.073 |
| Road transport | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| Other mobile sources and machinery | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 | 0.000 |
| Waste treatment and disposal | 0.005 | 0.005 | 0.002 | 0.002 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 |
| Total | 0.154 | 0.166 | 0.157 | 0.179 | 0.133 | 0.114 | 0.122 | 0.121 | 0.116 | 0.103 |

Among the most significant variations, emissions from *combustion in industry* and from *production processes* represent 10% and 71% of the total respectively, showing the former a decrease of -75% and the latter of 20%. The noticeable decline between 2008 and 2009 was due to the implementation of abatement systems in the largest Italian steel plant. The other relevant sector is *non industrial combustion plants* accounting for 17% and relevantly increasing (21%) between 1990 and 2020. The share of other sectors is lower than 1%.

3 ENERGY (NFR SECTOR 1)

3.1 OVERVIEW OF THE SECTOR

For the pollutants and sources discussed in this section, emissions result from the combustion of fuel. All the pollutants reported under the UNECE/CLRTAP are estimated. Stationary and mobile categories are covered for:

- Electricity production (power plants and Industrial producers);
- Refineries (Combustion);
- Iron and steel industries (Combustion)
- Chemical and petrochemical industries (Combustion);
- Construction industries (roof tiles, bricks);
- Other industries (metal works factories, food, textiles, others);
- Road Transport;
- Coastal Shipping;
- Railways;
- Aircraft;
- Domestic heating;
- Commercial heating;
- Public Service;
- Fishing and Agriculture.

Fugitive emissions are also reported under the energy sector as well as emissions from geothermal production.

The national emission inventory is prepared using energy consumption information available from national statistics and an estimate of the actual use of the fuels. The latter information is available at sectoral level in a different number of publications and different details, such as fuel consumption, distance travelled or some other statistical data related to emissions. For most of the combustion source categories, emissions are estimated from fuel consumption data reported in the National Energy Balance (BEN) as supplied by the Ministry for the Economic Development (MSE, several years (a)) and reported to the international energy organization, and from emission factors appropriate to the type of combustion and the pollutant.

The estimate from fuel consumption emission factors refers to stationary combustion in boilers and heaters. The other categories are estimated by more complex methods discussed in the relevant sections. The fuel consumption of “Other industries” is estimated so that the total fuel consumption of these sources is consistent with the national energy balance.

Electricity generation by companies primarily for their own use is auto-generation, and the relevant emissions should be reported under the industry concerned. However, national energy statistics report emissions from electricity generation as a separate category. The Italian inventory makes an overall calculation and then attempts to report as far as possible according to the guidelines:

- auto-generators are reported in the relevant industrial sectors of section “1.A.2 Manufacturing Industries and Construction”;
- refineries auto-generation is included in section 1.A.1b;
- iron and steel auto-generation is included in section 1.A.1c
- incinerators auto-generation of energy and heat is included in section 1.A.4a.

These reports are based on estimates of fuel used for steam generation connected with electricity production supplied by the National Independent System Operator (TERNA, several years).

Emissions from the energy production plants in integrated iron and steel plants and emissions from coke ovens are included in 1.A.1c category. Emissions from waste incineration facilities with energy recovery are reported under category 1.A.4a i (Combustion activity, commercial/institutional sector), whereas emissions from other types of waste incineration facilities are reported under category 5.C (Waste incineration). In particular, for 2020, almost 99% of the total amount of waste incinerated is treated in plants with energy recovery system. The energy recovered by these plants is mainly used for district heating of commercial buildings or used to satisfy the internal energy demand of the plants. Different emission factors for municipal, industrial and oils, hospital waste, and sewage sludge are applied, as reported in the waste chapter. Waste amount is then converted in energy content applying the relevant factor as resulting from data provided by TERNA, which in 2020 is equal to 11.4 GJ/t of waste.

Landfill gas is generally recovered and used for heating and power in commercial facilities, the resulting emissions are reported under 1.A.4.a. Biogas recovered from the anaerobic digester of animal waste is used for utilities in the agriculture sector and relative emissions are reported under 1.A.4.c.

Under 1.A.2 g vii industrial off road machinery are reported; the methodology used to estimate emissions from a range of portable or mobile equipment powered by reciprocating diesel engines is summarized. Industrial off-road include construction equipment such as bulldozers, loaders, graders, scrapers, rollers and excavators and other industrial machines as portable generators, compressors and cement mixers. Estimates are calculated taking in account especially the population of the different classes, annual usage, average power rating, load factor and technology distribution (EURO) according to the Guidebook (EMEP/EEA, 2019). COPERT II has been used for years 1994 and 1995 to estimate emissions and average emission factors for vehicles and diesel fuel consumption. Population data have been estimated on the basis on a survey of machinery sales. Machinery lifetime was estimated on the European averages reported in EMEP/CORINAIR, 2007, the annual usage data were taken either from industry or published data by EEA. The emission factors used came from EMEP/EEA and COPERT. The load factors were taken from COPERT. It was possible to calculate fuel consumptions for each class based on fuel consumption factors given in EMEP/CORINAIR, 2007. Comparison with known fuel consumption for certain groups of classes suggested that the population method overestimated fuel consumption by factors of 1.2-1.5 for industrial vehicles. Time series is reconstructed in relation to the diesel fuel use in industry reported in the national energy balance as gasoil final consumption. Emission factors for NO_x, CO, NMVOC and PM have been updated taking in account the reduction factors established in the European Directive 97/68/EC, the timing of application of the new limits and the tax of penetration of the new industrial vehicles in the total fleet. Emission reduction factor reported in the European Directive 2004/26/EC Directive have been applied and introduced in the emission estimates.

The emission factors used are based on national sources, or else on values specified in the EMEP/EEA guidebook and/or IPCC guidelines which are appropriate for Italy. Emission factors used for energy and manufacturing industries and non-industrial combustion, specifically categories 1A1, 1A2, 1A4, and their references are available on the ISPRA website at <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/> as well as emission factors for road transport (1A3b) are available at <http://www.sinanet.isprambiente.it/it/sia-ispra/fetransp/>.

In 2020 the energy sector accounts for more than 50% of total emissions for all the estimated pollutants, except for NMVOC, which accounts for 36%, PCB for 28% and ammonia for 2%. In particular, emissions from the energy sector are 94% of CO, 93% of SO_x and BC, 89% of NO_x, 86% of PM_{2.5} and PAH national total emissions.

In 2020, the following categories are key categories for different pollutants: *public electricity and heat production* (1A1a), *stationary combustion in iron and steel industries* (1A2a), *stationary combustion in non-ferrous metal industries* (1A2b), *stationary combustion in non-metallic mineral industries* (1A2f), *stationary combustion in manufacturing industries and construction: other* (1A2gviii), *road transport categories* (1A3b), *national navigation* (1A3d ii), *stationary combustion plants in commercial/institutional* (1A4a i) and *residential* (1A4b i), *off-road vehicles in agriculture, forestry and fishing* (1A4c ii), *fugitive emissions from refining and storage* (1B2a iv), and *other fugitive emissions from energy production* (1B2d).

The same categories are key categories for 1990 and for the trend analysis. In addition, for 1990, petroleum refining (1A1b) for SO_x, PM_{2.5}, *stationary combustion in chemical industries* (1A2c) for SO_x and PM₁₀, *mobile combustion in manufacturing industries and construction* (1A2g vii) for BC, *stationary combustion in other industries* (1A2g viii) for SO_x, and *fugitive emissions from distribution of oil products* (1B2a v) for NMVOC emissions are also key categories.

The following sections present an outline of the main key categories in the energy sector. Table 3.1 highlights the key categories identified in the sector.

The energy sector is the main source of emissions in Italy with a share of more than 80% for different pollutants under the UNECE convention; specifically, for the main pollutants, in 2020 the sector accounts for:

- 94% in national total CO emissions;
- 93% in national total BC emissions;
- 93% in national total SO_x emissions;
- 89% in national total NO_x emissions;
- 86% in national total PM_{2.5} emissions;
- 86% in national total PAH emissions.

Moreover, the sector is also an important source for heavy metals; specifically in 2020, energy sector is responsible for 55% of total Cd emissions, 54% for Hg and 59% for lead emissions.

There are no differences as compared to the sectoral share in 1990, except for lead whose contribution in 1990 was 98% of total emissions, 30% higher than in 2020 and for PAH whose contribution in 1990 was 44%, 40% lower than in 2020.

One of the most important source of emissions in the sector and key category, in 2020, is represented by *road transport* (1A3b), at least for the main pollutants: NO_x (37.4%), BC (24.2%), CO (15.7%), NMVOC (10.4%), Cd (8.6%) and particulate matter (PM10 9.3%, PM2.5 8.0%). There has been a strong reduction in lead emissions from 1990 to 2020 in *road transport* due to replacement of lead gasoline. An in depth analysis of the road transport category and its emission trends is reported in paragraph 3.5.

Manufacturing industries and construction (1A2) is a main source of heavy metals and POPs, accounting for about 43% of lead total emissions, 30% for cadmium, 28% for mercury, 42% for HCB, and 17% for dioxin. The sector is key category also for PM10 and PM2.5 (4%) as well as SO_x, NO_x and CO, about 39%, 9% and 5% of total emissions. The main sectors are iron and steel sector, which is key for SO_x, CO, Cd and Hg, the non-ferrous metal sector, key for Hg, HCB and Dioxin, and non-metallic mineral sector that is key category for SO_x, NO_x, PM10, PM2.5, Pb, Cd and Hg.

Public electricity and heat production (1A1a) is a key source of SO_x emissions in 2020 with a share of 6.1%, Hg (6.4%) and NO_x emissions (4.2%).

National *navigation* (1A3d ii) is key category for SO_x (9.8%), NO_x (15.0%), PM10 (3.9%), PM2.5 (4.8%), CO (2.8%) and BC (6.9%). The weight of this category on the total emissions has increased for SO_x and NO_x during the period because of a sectoral delay in the introduction of relevant normative to reduce air emissions.

A sector increasing its level of emissions is the *non-industrial combustion* (1A4): NO_x and NMVOC, emissions of this category account in 2020 for 20.0% and 19.4% of national total, respectively; SO_x emission account for 12.0%; CO emissions account for 66.5%; Cd emissions account for 12.2%; PM10 and PM2.5 emissions account for 55.3% and 67.9% respectively while BC emissions account for 56.9%; dioxin is 34.7%, PAH is 80.7% , PCB is 17.4% and HCB is 26.7% of national totals. These emissions are prevalently due to biomass combustion, in winter, and they are also becoming critical for air quality issues and for HCB due to the increase of combustion of waste with energy recovery reported under the sector. An in depth analysis of this category is reported in the paragraph 3.12.

Fugitive emissions in refinery from fossil fuel distribution and storage (1B2a iv) is key category in 2020 for SO_x emissions (13.7%). Total SO_x fugitive emissions from distribution of fossil fuels account for 18.4% of the total. *Fugitive emissions from geothermal energy production* is key category for Hg for 5.9% of national total emissions.

Table 3.1 Key categories in the energy sector in 2020

| | 1A1 a | 1A1 b | 1A1 c | 1A2 | 1A2 g vii | 1A3 a i | 1A3 a ii | 1A3 b i | 1A3 b ii | 1A3 b iii | 1A3 b iv | 1A3 b v | 1A3 b vi | 1A3 b vii | 1A3 c | 1A3 d ii | 1A3 e i | 1A4 a i | 1A4 b i | 1A4 b ii | 1A4 c | 1A5 b | 1B1 a | 1B1 b | 1B2 |
|-------------------------|----------|----------|----------|------|--------------|------------|-------------|------------|-------------|--------------|-------------|------------|-------------|--------------|----------|-------------|------------|------------|------------|-------------|----------|----------|----------|----------|------|
| SO_x | 6.1 | 4.9 | 2.3 | 38.5 | 0.0 | 0.2 | 0.1 | 0.2 | 0.1 | 0.1 | 0.0 | | | | 0.0 | 9.8 | 0.0 | 5.3 | 6.6 | 0.0 | 0.1 | 0.2 | | | 18.4 |
| NO_x | 4.2 | 1.3 | 0.4 | 7.9 | 0.6 | 0.3 | 0.2 | 19.1 | 6.2 | 11.7 | 0.4 | | | | 0.3 | 15.0 | 0.1 | 5.5 | 6.8 | 0.0 | 7.7 | 0.6 | | | 0.9 |
| NH₃ | 0.0 | 0.0 | 0.0 | 0.3 | 0.0 | | | 1.1 | 0.0 | 0.1 | 0.0 | | | | 0.0 | 0.0 | | 0.0 | 0.3 | 0.0 | 0.0 | 0.0 | | | 0.3 |
| NM VOC | 0.3 | 0.1 | 0.0 | 0.7 | 0.1 | 0.0 | 0.0 | 1.9 | 0.1 | 0.2 | 3.0 | 5.2 | | | 0.0 | 1.5 | 0.0 | 3.2 | 15.1 | 0.1 | 0.9 | 0.1 | | 0.1 | 3.8 |
| CO | 1.0 | 0.2 | 0.9 | 4.7 | 0.3 | 0.1 | 0.1 | 9.5 | 0.3 | 1.1 | 4.7 | | | | 0.0 | 2.8 | 0.0 | 1.5 | 62.3 | 0.1 | 2.6 | 1.3 | | | 0.0 |
| PM₁₀ | 0.2 | 0.1 | 0.0 | 4.1 | 0.1 | 0.0 | 0.0 | 1.6 | 0.4 | 0.6 | 0.3 | | 4.2 | 2.3 | 0.0 | 3.9 | 0.0 | 0.8 | 53.1 | 0.0 | 1.4 | 0.3 | 0.1 | 0.0 | 0.0 |
| PM_{2.5} | 0.2 | 0.1 | 0.0 | 4.2 | 0.1 | 0.0 | 0.0 | 2.0 | 0.5 | 0.7 | 0.4 | | 2.9 | 1.5 | 0.0 | 4.8 | 0.0 | 1.0 | 65.2 | 0.0 | 1.7 | 0.4 | 0.0 | 0.0 | 0.0 |
| BC | 0.1 | 0.0 | 0.1 | 1.6 | 0.6 | 0.0 | 0.0 | 13.3 | 3.2 | 4.0 | 0.6 | | 2.6 | 0.5 | 0.2 | 6.9 | 0.0 | 0.8 | 49.7 | 0.0 | 6.3 | 1.7 | 0.1 | 0.1 | 0.0 |
| Pb | 0.7 | 0.2 | 0.0 | 43.5 | | 0.2 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | | 5.5 | | | 0.1 | | 3.2 | 5.5 | | 0.0 | 0.1 | | 0.2 | 0.0 |
| Cd | 1.2 | 0.4 | 0.0 | 30.4 | 0.0 | 0.0 | 0.0 | 5.6 | 0.6 | 0.5 | 0.8 | | 1.1 | | 0.0 | 0.4 | | 2.2 | 9.5 | 0.0 | 0.5 | 0.0 | | 1.9 | 0.0 |
| Hg | 6.4 | 1.4 | 0.2 | 28.3 | | | | 1.7 | 0.3 | 0.5 | 0.1 | | | | | | | 3.8 | 4.0 | | 0.3 | | 0.7 | 5.9 | |
| PAH | 0.7 | 0.0 | 0.0 | 0.7 | 0.0 | 0.0 | 0.0 | 2.3 | 0.4 | 0.6 | 0.0 | | 0.0 | | 0.0 | 0.1 | | 1.0 | 79.0 | 0.0 | 0.7 | 0.0 | | 0.0 | |
| Dioxin | 0.7 | 0.8 | | 16.8 | | | | 1.9 | 0.2 | 0.3 | 0.1 | | | | | 0.1 | | 1.0 | 33.2 | | 0.5 | | | | |
| HCB | 5.4 | | | 42.3 | | | | 0.0 | 0.0 | 0.0 | 0.0 | | | | | 0.8 | | 11.1 | 11.5 | | 4.2 | | | | |
| PCB | 0.4 | | | 10.1 | | | | 0.0 | 0.0 | 0.0 | 0.0 | | | | | 0.0 | | 2.6 | 14.5 | | 0.3 | | | | |

Note: key categories are shaded in blue

3.2 ENERGY INDUSTRIES

This chapter deals with emissions from categories 1A1a Public electricity and heat production, 1A1b Petroleum refining and 1A1c Manufacture of solid fuels and other energy industries.

3.2.1 Methodological issues

Methodologies used for estimating emissions from this sector are based on and conform to the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007; EMEP/EEA, 2013; EMEP/EEA, 2016), the IPCC Guidelines (IPCC, 1997; IPCC, 2006) and the Good Practice Guidance (IPCC, 2000). A detailed description on the methods and national specific circumstances as well as reference material of the energy sector is documented in the national inventory report of the Italian greenhouse gas inventory (ISPRA, 2022[a]). The National Energy Balance, published by the Ministry of Economic Development, is the main source of information to estimate emissions from the energy sector as it reports fuel consumption for different sectors at national level. Additional information for electricity production is provided by the major national electricity producers and by the major national industry corporation. Other data are communicated by different category associations.

The use of PRTR data has been intensified in recent years, as also described in the following paragraphs. In this regard it should be noted that in response to the review process a survey has been conducted to verify if emission data submitted by operators are calculated subtracting the confidence interval. The issue has been discussed also with the colleagues from the Ministry of Environment (IMELS) in charge of the implementation at national level for the IED legislation. In principle it is to be noted that the validated average values (with the confidence interval subtracted from the measured data) are the data used to verify the compliance of the operators to prescriptions included in the permits issued to the same operators and not for the calculation of the total annual emissions submitted in the framework of the relevant European Union Directives and Regulations. In addition, the implementation at national level of the IED requires Italian operators with emissions reported on the basis of Continuous Monitoring System (CMS) data to refer to confidence intervals which are not those included in the IED: in fact, the confidence intervals must result from quality assurance procedure and the implementation of UNI EN 14181:2005 and QAL2 procedure. This national circumstance implies that the validated average values used by the Italian operators are more realistic compared to those calculated using the procedure laid out in the IED. Moreover, the use of CMS at the stacks is largely implemented at facilities with installations exceeding 50 MWth input. In order to assess consistency along the timeseries, data reported by the largest Italian operator in the Energy production (about 25% of energy production) show that no issues concerning consistency can be raised. For consistency issue we believe that official data, as air emission values, communicated by the operators in the EU official frameworks, as the LCP Directive, PRTR registry and IPPC Directive should be considered as they were reported and without any further adjustment (apart from QA/QC procedures).

Notation key NO for activity is used indicating that a fuel is not consumed at all while NA is reported in the column where is requested to specify a different indicator than fuel consumption.

3.2.1.1 Public Electricity and Heat Production

For 1A1 categories, a Tier 3 is used and SO_x, NO_x and PM₁₀ emissions are estimated on the basis of emission and consumption data provided by the relevant plants in the framework of LCP and ETS European Directives and EPRTR Regulation. The average implied emission factors at fuel level result from the analysis of the information provided and available at plant level, including technologies for energy production and emissions abatement. These IEF at fuel level have been used to estimate emissions for those plants where some pollutants have not been declared and to verify emissions declared. PM_{2.5} is estimated applying the ratio between PM_{2.5} and PM₁₀ reported in the Tier 2 tables of the EMEP/EEA 2019 Guidebook at fuel level.

With regard to heavy metals country specific emission factors for each fuel have been used to estimate emissions as provided by the main national operator in relation with the technologies in 2001 while for PCB, emission factors for coal, oil products and wood biomass from the EMEP/EEA Guidebook 2019 have been used following the recommendation of the review process (EEA, 2019). Emission factors for the PAH, Dioxin and HCB for Italy are from a study of TNO at European level (Berdowski et al, 1997). A comparison with data

from PRTR has been made but in the case of HMs and POPs the information provided in that framework is generally poor and regard only few plants probably because of the ceiling which is high. For most of the pollutants the available information is not representative of the total. Dioxin, PCB and HCB emissions are generally missing. For the next submission we plan to improve the quality of HMs and POPs estimate on the basis of the figures reported in the PRTR updating the EFs where they are representative of the national total or at least checking the order of magnitude of the values already used when the figures from PRTR are not sufficient to be considered representative of the total. In particular for power plants and coal fuelled main plants some HMs under the PRTR are available.

In the following Table 3.2 the main EFs for Public Electricity and Heat production sector are reported. The data shown in the blue cells are country specific emission factors as they are derived from specific measurement campaigns and direct communications by the operators. PM10 EFs too are country specific while PM2.5 EFs have been derived from PM10 EFs and the ratio PM2.5/PM10 per fuel as reported in the EMEP/EEA Guidebook 2019. Finally, Dioxins and PAH EFs come from a study of TNO at European level (Berdowski et al, 1997).

Table 3.2 *Main emission factors for Public Electricity and Heat Production - 2020*

| | SO _x | NO _x | NMVOC | CO | Cd | Hg | Pb | PM2.5 | Dioxins | PAH |
|-----------------------|-----------------|-----------------|-------|------|-------|-----|-----|-------|---------|-----------|
| | g/GJ | | | | mg/GJ | | | g/GJ | | |
| Steam coal | 31.7 | 40.2 | 1.5 | 12.0 | 0.1 | 0.9 | 4.6 | 0.36 | 0.0038 | 0.0000009 |
| Lignite | 382.2 | 40.2 | 1.5 | 12.0 | 0.1 | 0.9 | 4.6 | 0.33 | 0.0096 | 0.0000024 |
| Natural gas | 0.1 | 14.4 | 2.5 | 20.0 | 0.0 | 0.3 | 0.6 | 0.01 | - | - |
| Gasoil | 14.7 | 61.9 | 1.5 | 12.0 | 0.1 | 0.8 | 2.5 | 0.05 | 0.0234 | 0.0008903 |
| Low sulphur fuel oil | 38.0 | 28.1 | 3.0 | 15.0 | 0.1 | 0.8 | 2.5 | 0.77 | 0.0244 | 0.0008903 |
| High sulphur fuel oil | 38.0 | 28.1 | 3.0 | 15.0 | 0.1 | 0.8 | 2.5 | 0.77 | 0.0244 | 0.0008903 |
| Other fuel oils | 13.2 | 28.1 | 3.0 | 15.0 | 0.1 | 0.8 | 2.5 | 0.77 | 0.0244 | 0.0008903 |
| Biomass | 2.7 | 59.2 | 1.5 | 12.0 | | | | 2.10 | 0.0096 | 0.0036558 |

3.2.1.2 Refineries

As stated above, for 1A1 categories, a Tier 3 is used and SO_x, NO_x and PM10 emissions are estimated on the basis of emission and consumption data provided by the relevant plants in the framework of LCP and ETS European Directives and EPRTR Regulation. The average implied emission factors at fuel level result from the analysis of the information provided and available at plant level, including technologies for energy production and emissions abatement. These IEF at fuel level have been used to estimate emissions for those plants where some pollutants have not been declared and to verify emissions declared. PM2.5 is estimated applying the ratio between PM2.5 and PM10 reported in the Tier 2 tables of the EMEP/EEA 2019 Guidebook at fuel level. In particular for 1A1b category, the implied emission factor refers both to the production of energy and heat and to the other combustion activities in refineries.

With regard to heavy metals country specific emission factors for each fuel have been used to estimate emissions as provided by the main national operator in relation with the technologies in 2001 while for PCB, emission factors for coal, oil products and wood biomass from the EMEP/EEA Guidebook 2019 have been

used following the recommendation of the review process (EEA, 2019). A comparison with data from PRTR has been made but in the case of HMs and POPs the information provided in that framework is generally poor and regard only few plants probably because of the ceiling which is high. For most of the pollutants the available information is not representative of the total. Dioxin, PCB and HCB emissions are generally missing. For the next submissions we plan to improve the quality of HMs and POPs estimate on the basis of the figures reported in the PRTR updating the EFs where they are representative of the national total or at least checking the order of magnitude of the values already used when the figures from PRTR are not sufficient to be considered representative of the total. For refineries a quite complete reporting is available for selenium and zinc. In the first case the average EFs is very close to the default used while for zinc the resulting EFs from EPRTR is one order bigger than the default used up to now.

In the following Table 3.3 the main EFs for the Refinery sector are reported. The data shown in the blue cells are country specific emission factors as they are derived from specific measurement campaigns and direct communications by the operators. PM10 EFs too are country specific while PM2.5 EFs have been derived from PM10 EFs and the ratio PM2.5/PM10 per fuel as reported in the EMEP/EEA Guidebook 2019. Finally, Dioxins and PAH EFs come from a study of TNO at European level (Berdowski et al, 1997).

Table 3.3 *Main emission factors for Refineries - 2020*

| | SO _x | NO _x | NM VOC | CO | Cd | Hg | Pb | PM2.5 | Dioxins | PAH |
|---------------------------------|-----------------|-----------------|--------|------|-------|------|------|-------|---------|--------|
| | g/GJ | | | | mg/GJ | | | g/GJ | | |
| Combustion plants in refineries | | | | | | | | | | |
| Refinery gas | 5.8 | 28 | 2.5 | 13 | - | - | - | 0.30 | - | - |
| Kerosene | 42 | 28 | 3 | 60 | 0.14 | 0.77 | 2.52 | 0.15 | 0.0230 | 0.0004 |
| Petcoke | 42 | 28 | 1.5 | 9 | 0.14 | 0.77 | 2.52 | 0.40 | 0.0288 | 0.0004 |
| Tar + synthesis gasses | 42 | 28 | 3 | 8 | 0.14 | 0.77 | 2.52 | 0.69 | 0.0244 | 0.0004 |
| High sulphur fuel oil | 42 | 28 | 3 | 10 | 0.14 | 0.77 | 2.52 | 0.69 | 0.0244 | 0.0004 |
| Natural gas | 0.2 | 9 | 2.5 | 13 | 0.03 | 0.31 | 0.60 | 0.01 | - | - |
| LPG | 0.2 | 28 | 2 | 10 | - | - | - | 0.01 | 0.0217 | - |
| Refinery furnaces | | | | | | | | | | |
| LPG | 0.2 | 42 | 2 | 10 | - | - | - | 0.01 | 0.0217 | - |
| Refinerygas | 5.8 | 42 | 2.5 | 13.0 | - | - | - | 0.30 | - | - |
| Kerosene | 42 | 42 | 3 | 60 | 0.14 | 0.77 | 2.52 | 0.15 | 0.0230 | 0.0004 |
| Gasoline | 23 | 42 | 3 | 60 | 0.14 | 0.77 | 2.52 | 0.08 | 0.0230 | 0.0004 |
| Gasoil | 47 | 42 | 3 | 10 | 0.14 | 0.77 | 2.52 | 0.08 | 0.0234 | 0.0004 |
| Highsulphur ueloil | 42 | 42 | 3 | 10 | 0.14 | 0.77 | 2.52 | 0.69 | 0.0244 | 0.0004 |
| Lowsulphur eloil | 42 | 42 | 3 | 10 | 0.14 | 0.77 | 2.52 | 0.69 | 0.0244 | 0.0004 |
| Petcoke | 42 | 42 | 1.5 | 9 | 0.14 | 0.77 | 2.52 | 0.40 | 0.0288 | 0.0004 |
| Noenergyfuel | 42 | 42 | 3 | 10 | 0.14 | 0.77 | 2.52 | 0.69 | 0.0244 | 0.0004 |

3.2.1.3 Manufacture of Solid Fuels and Other Energy Industries

As already mentioned, for 1A1 categories, a Tier 3 is used and SO_x, NO_x and PM₁₀ emissions are estimated on the basis of emission and consumption data provided by the relevant plants in the framework of LCP and ETS European Directives and EPRTTR Regulation. The average implied emission factors at fuel level result from the analysis of the information provided and available at plant level, including technologies for energy production and emissions abatement. These IEF at fuel level have been used to estimate emissions for those plants where some pollutants have not been declared and to verify emissions declared. PM 2.5 is estimated applying the ratio between PM_{2.5} and PM₁₀ reported in the Tier 2 tables of the EMEP/EEA 2019 Guidebook at fuel level.

In particular, for coke production according to the review (EEA, 2019) PAH emission factor have been disaggregated into those deriving from the combustion process and the fugitive ones and estimated with the emission factors in the Guidebook (EMEP/EEA, 2019).

In the following Table 3.4 the main EFs for the Manufacture of Solid Fuels and Other Energy Industries are reported. The data shown in the blue cells are country specific emission factors as they are derived from specific measurement campaigns and direct communications by the operators. PM₁₀ emission factors too are country specific while PM_{2.5} have been derived from PM₁₀ and the ratio PM_{2.5}/PM₁₀ per fuel as reported in the EMEP/EEA Guidebook 2019. Finally, Dioxins and PAH emission factors come from a study of TNO at European level (Berdowski et al, 1997).

Table 3.4 Main emission factors for Manufacture of Solid Fuels and Other Energy Industries - 2020

| | SO _x | NO _x | NMVOC | CO | Cd | Hg | Pb | PM _{2.5} | Dioxins | PAH |
|-------------------------|-----------------|-----------------|-------|-------|-------|------|------|-------------------|---------|----------|
| | g/GJ | | | | mg/GJ | | | g/GJ | | |
| Combustion in coke oven | | | | | | | | | | |
| Steam coal | 32 | 40 | 1.5 | 12 | 0.15 | 0.85 | 4.63 | 3.89 | 0.004 | 0.000001 |
| Natural gas | 0.1 | 9 | 2.5 | 20 | 0.03 | 0.31 | 0.60 | 0.14 | - | - |
| Oxygen furnaces gas | 78 | 68 | 2.5 | 13 | 0.03 | 0.31 | 0.60 | 0.14 | - | - |
| Coke oven gas | 78 | 68 | 2.5 | 13 | 0.03 | 0.31 | 0.60 | 0.14 | - | - |
| Blast furnace gas | 78 | 68 | 2.5 | 13 | 0.03 | 0.31 | 0.60 | 0.14 | - | - |
| High sulphur fuel oil | 80 | 30 | 3 | 15 | 0.14 | 0.77 | 2.52 | 0.14 | 0.024 | 0.00089 |
| Coking coal | 369 | 492 | 2 | 12 | 0.03 | 0.31 | 0.60 | 18 | | |
| Coke oven furnaces g/Mg | | | | | | | | | | |
| Coke production | 396 | 520 | 145 | 11157 | | | | 20 | | 0.08 |

3.2.2 Time series and key categories

The trend of emissions from energy industries has been influenced by the implementation of the legislative framework and by the evolution and replacement of fuels and these effects can be noticed in particular from the analysis of the NO_x and SO_x trends.

The adoption of new regulations, in the late 80s, has led to a shift in energy consumption from oil with high sulphur content to oil with lower sulphur content and to natural gas.

In recent years, the conversion to the use of natural gas to replace fuel oil has intensified, thanks to incentives granted for the improvement of energy efficiency. Furthermore, a significant reduction in the use of coal fuels for energy production has been recorded in the last years. These measures, together with those of promoting renewable energy and energy saving, have led to a further reduction of emissions in the sector. In addition, in the last years, more stringent emission limits to the new plants have been established during the authorisation process with the aim to prevent air quality issues at local level.

3.2.2.1 Public Electricity and Heat production

This paragraph refers to the main electricity producers that produce electricity for the national grid. From 1998 onwards, the expansion of the industrial cogeneration of electricity and the split of the national monopoly have transformed many industrial producers into “independent producers”, regularly supplying the national grid. These producers account in 2020 for 92.8% of all electricity produced with combustion processes in Italy (TERNA, several years).

In 2020, Public Electricity and Heat production is a key category for SO_x, NO_x and Hg.

Public electricity and heat production (1A1a) is a key source of SO_x emissions in 2020 with a share of 6.1%, Hg (6.4) and NO_x emissions (4.2%). A strong reduction of SO_x, NO_x and PM10 emissions is observed for this category along the time series (as well as for 1A2 sector). The introduction of two regulatory instruments: the DPR 203/88 (Decree of President of the Republic of 24th May 1988), laying down rules concerning the authorisation of plants, and the Ministerial Decree of 12th July 1990, which introduced plant level limits to emissions of PM10, NO_x and SO_x for new plants and required old plants to conform to the limit by 1997, explained the emission reduction in the nineties. The shift from fuel oil to natural gas combined with the increase of energy efficiency of the plants and the introduction of PM10 abatement technologies have been implemented to comply with the emission limit values. From 2000 lower limits to emissions at the stacks have been introduced, in the framework of environmental integrated authorisations, for the authorisation of new plants and the implementation of the old ones, especially for those facilities located in areas with air quality critical values. For this reason the plants have increased the use of natural gas heat and power combined technology. In 2020 in Italy there are still 8 coal plants, and 2 fuel oil plant out of around 190 power plants included in this source category. With exception of few biomass plants and some gasoil stationary engines in the small islands, the other plants are natural gas combined cycle thermoelectric power plant.

3.2.2.2 Refineries

This subsector covers the energy emissions from the national refineries (14 plants in 2020), including the energy used to generate electricity for internal use and exported to the national grid by power plants that directly use off-gases or other residues of the refineries. These power plants are generally owned by other companies but are located inside the refinery premises or just sideway. In 2020 the power plants included in this source category have generated 6.7% of all electricity produced with combustion processes in Italy.

Petroleum refining (1A1b) is not a key category in 2020. In addition, for 1990, petroleum refining (1A1b) is a key category for SO_x and PM2.5. Emissions are estimated on the basis of emission and consumption data provided by refineries in the framework of LCP, ETS European Directives and EPRTR Regulation and refer both to the production of energy and heat and to the other combustion activities in the plants. Emission trends are driven by the same legislation quoted for 1A1a category, where specific rules and ceiling were set up for refineries.

3.2.2.3 Manufacture of Solid Fuels and Other Energy Industries

In this section, emissions from power plants, which use coal gases, are also reported. In particular, we refer to the electricity generated in the iron and steel plant sites (using coal gases and other fuels). In 2020 the power plants included in this source category have generated about 1.6% of all electricity produced with combustion processes in Italy.

Manufacture of Solid Fuels and Other Energy Industries (1A1c) is not a key category in 2020. Emissions show a decreasing trend linked to a reduction in activity data (only 2 plants remaining in 2020) and to the implementation of abatement technologies and the plants revamping. About abatement technologies, the largest integrated plant in Italy (and in Europe), in the last ten years, has carried out several interventions on the coking plant in the framework of the IPPC permit, applying BAT in several cases.

3.2.3 QA/QC and verification

A complete description of methodological and activity data improvements is documented every year in a QA/QC plan (ISPRA, 2022[b]).

The analysis of data collected from point sources allowed to distribute emissions at local level, for 2019 and previous years, as submitted under the CLTRAP. To illustrate an example, NO_x emissions from point sources are reported in Figure 3.1 for the year 2019. Point sources include public electricity and heat production plants and stationary combustion plants in manufacturing industries and construction.

The figure highlights that the most critical industrial areas are distributed in few regions.




Figure 3.1 NO_x emissions from point sources in 2019 (t)

Similarly, in Figure 3.2 NO_x emissions from point sources in 2015 are reported.




Figure 3.2 *NO_x emissions from point sources in 2015 (t)*

In Figure 3.3, NO_x emissions communicated by 229 facilities (power plants, refineries, cement plants and iron and steel integrated plants), in the framework of the national E-PRTR register and LCP Directive, have been processed and geographically located. The territorial distribution shows similar results to those reported in the previous figure highlighting the industrial areas still in activity in 2010.




Figure 3.3 *NO_x emissions from point sources in 2010 (t)*

Every four years emissions are disaggregated at regional and provincial levels and figures are compared to the results obtained by regional bottom-up inventories. Emissions disaggregated at local level are also used as input for air quality modelling. NO_x emissions from *road transport* have been disaggregated at NUTS3 level; the disaggregation related to the year 2019 is reported in Figure 3.4 whereas methodologies are described in the relevant publication (ISPRA, 2009).




Figure 3.4 *NO_x emissions from road transport in 2019 (t)*

3.2.4 Recalculations

In the 2022 submission minor recalculations have been performed for energy industries, in particular emissions from refineries have been updated from 2017.

3.2.5 Planned improvements

Specific improvements are detailed in the 2022 QA/QC plan (ISPRA, 2022[b]).

For the *energy* sector, a major progress regards the management of the information system where data collected in the framework of different obligations, Large Combustion Plant, E-PRTR and Emissions Trading, are gathered together thus highlighting the main discrepancies in information and detecting potential errors. Moreover, the complete use of the energy data provided by the Ministry of Economic Development to the Joint Questionnaire IEA/OECD/EUROSTAT is planned in substitution of the national energy balances used till now; liquid, gaseous and solid fuel are now aligned for the whole time series and we plan for the next submission to update as possible renewable fuels and biomass.

With respect to PM10 and heavy metals emissions from *Public Electricity and Heat Production* category (1A1a) while PM10 emissions are updated every year on the basis of data submitted by the plants in the framework of the EPRTR registry, Large Combustion Plants Directive and Environmental Reports, heavy metals emission factors time series have been reconstructed from 1990 to 2001 on the basis of a study conducted by ENEL (major company in Italy) which reports heavy metals emissions measurements by fuel and technology (with or without PM10 abatement technologies) of relevant national plants. From 2001 these emission factors have not been updated. Heavy metals emission data in the EPRTR registry refer only to few not representative plants and are not sufficient to calculate average emission factors. Further work is planned to update/change emission factors for those pollutants where figures reported in the EPRTR lead to average values significantly different from those actually used.

3.3 MANUFACTURING INDUSTRIES AND CONSTRUCTION

This chapter deals with emissions from Stationary combustion in manufacturing industries and construction, in particular from categories 1A2a Iron and steel, 1A2b Non-ferrous metals, 1A2c Chemicals, 1A2d Pulp, Paper and print, 1A2e Food processing, beverages and tobacco, 1A2f Non-metallic minerals, 1A2g Other.

3.3.1 Methodological issues

For 1A2 categories, estimates for chemical, food processing, and other sectors (as textile, mechanics, extraction) are based on fuel consumption where EMEP/CORINAIR 2007 emission factors at fuel level have been used except for SO_x, NO_x and PM10 which are estimated on the basis of emission and consumption data provided by the relevant plants in the framework of LCP and ETS European Directives and EPRTR Regulation. PM 2.5 is estimated applying the ratio between PM2.5 and PM10 reported in the Tier 2 tables of the EMEP/EEA 2019 Guidebook at fuel level. Emissions of NH₃ have been also included when available at plant level. With regard to heavy metals, country specific emission factors for each fuel have been used to estimate emissions as provided by the main national operator in relation with the technologies, while for PCB emission factors for coal, oil products and wood biomass from the EMEP/EEA Guidebook 2019 have been used following the recommendation of the review process (EEA,2019). Emission factors for the PAH, Dioxin and HCB for Italy are from a study of TNO at European level (Berdowski et al, 1997). For the iron and steel, non-ferrous metal, pulp and paper and non-metallic minerals sectors emission estimates are based on production data at SNAP category level. SO_x, NO_x and PM10 emission factors time series are estimated based on the communication from operators in the framework of LCP Directive and EPRTR Regulation and industrial association at SNAP activity code level. For NMVOC, default EFs of EMEP/CORINAIR 2007 Guidebook are prevalently used except for glass and lead production where country specific emission factors are used; emission factors provided in the EMEP/EEA 2019 Guidebook are not appropriate because of they are calculated for small combustion boilers while emissions in this category refer prevalently to boilers >20 MWt for auto-production of energy and heat in the industrial sectors.

The Institute, specifically the same unit responsible for the inventory, also collects data in the context of the European Emissions Trading Scheme, the National Pollutant Release and Transfer Register (Italian PRTR) and the Large Combustion Plants (LCP) Directives. All these data are managed and used to compile the inventory. Figures are cross checked to develop country-specific emission factors and input activity data; whenever data cannot be straight used for the inventory compilation, they are considered as verification. EPER/EPRTR data are yearly available from 2002 while ETS data from 2005 and LCP data from 1990 all on yearly basis. In the EPRTR registry total emissions divided by category are reported by plants if they exceed the relevant ceiling for each pollutant. LCP data refer only to SO_x, NO_x and PM emissions that are collected in stacks over 50 MWth and could result in figures lower than those reported in the EPRTR. In the ETS only CO₂ and fuel consumption data are reported. QA/QC checks at plants level are directed to check the submissions of data in the different context and evaluate the differences if any. For example, if emissions submitted by a plant under LCP are higher than those submitted under the EPRTR, ISPRA asks the operator of the reporting plant for an explanation and the verification of data submitted. In addition, on the basis of fuel consumption supplied under the ETS and average emission factor by fuel the energy experts estimate emissions at plant level and compare them with those submitted in the EPRTR and LCP. Also in this case ISPRA ask for clarifications to the reporting plant if necessary.

3.3.1.1 Iron and steel

More in detail 1A2a includes combustion activities from the iron and steel sector as blast furnace coppers, sinter plants and reheating furnaces. In 1990 there were four integrated iron and steel plants in Italy. In 2020, there are only two of the above-mentioned plants, one of which lacks BOF; oxygen steel production represents about 16.5% of the total production and the arc furnace steel the remaining 83.5% (FEDERACCIAI, several years). Currently, long products represent about 48% of steel production in Italy, flat products about 41%, and pipe the remaining 12%. Most of the flat production derives from only one integrated iron and steel plant while, in steel plants equipped with electric ovens almost all located in the northern regions, long products are predominantly produced (e.g carbon steel, stainless steels) and seamless pipes (only one plant) (FEDERACCIAI, several years). Dioxins, PCB, HCB, PAH, Cd and Pb emissions are estimated on the basis of country specific emission factors at activity level, especially referring to sinter plants production, as provided by the main national operators. In particular, HCB emissions come from sinter plant productions and the emission factor is from the 2006 EMEP/CORINAIR Guidebook and it is coherent with data provided by the main national operator, at least for past years. The update of HMs and POPs emission factors with last available data started in the 2022 submission. On the basis of data collected in the framework of the IPPC permits, emission factors from point sources (sinter plants and blast furnaces) have been updated for As, Ni, Pb, Se, Cu, Hg, Zn, in the case of blast furnace it was also possible to estimate the PM10 emission factor. Again, on the basis of the IPPC data, the emission factors of NO_x, SO_x and PM10 from reheating furnaces have been estimated. Emission factors derived from this survey on the IPPC permits have been applied since 2016.

As regards Cd emissions, these refer to blast furnaces, sinter and reheating activities; emission factors are those reported by the main Italian plant and emissions have been revised since 2003 based on the last review process and of E-PRTR data. For Hg and the other HMs emission factors are from the IPPC Bref sectoral report (JRC, 2013) and/or EMEP/EEA Guidebook 2006.

In the following Table 3.5 the main emission factors for the sector are reported. The data shown in the light blue cells are country specific based on emission data at plant level communicated in the framework of EPRTR, LCP, ETS or specific studies. Data reported in the dark blue cells represent country specific and fuel based emission factors. PM10 emission factors too, are country specific while PM2.5 factors have been derived from PM10 and the ratio PM2.5/PM10 per fuel as reported in the EMEP/EEA Guidebook 2019.

Table 3.5 Main emission factors 1A2 Iron and steel - 2020

| | SO _x | NO _x | NM VOC | CO | Cd | Hg | Pb | PM2.5 | PAH | Dioxins |
|-----------------------------------|-----------------|-----------------|--------|--------|--------------|-------|-------|--------|-------|---------------|
| | Kg/Mg product | | | | g/Mg product | | | | | mg/Mg product |
| Blast furnace coppers | | | | | | | | | | |
| areal sources | 0.300 | 0.100 | 0.0003 | 0.189 | 0.002 | 0.026 | 0.100 | 9.938 | - | - |
| point sources | 0.300 | 0.057 | 0.0003 | 0.189 | 0.002 | 0.001 | 0.007 | 0.754 | - | - |
| Sinter and pelletizing plants | | | | | | | | | | |
| areal sources | 0.515 | 0.859 | 0.090 | 16.105 | 0.023 | 0.020 | 3.700 | 54.593 | 0.025 | 0.001 |
| point sources | 0.860 | 0.684 | 0.090 | 13.215 | 0.004 | 0.004 | 0.323 | 29.131 | 0.025 | 0.0003 |
| Reheating furnaces steel and iron | | | | | | | | | | |
| areal sources | 0.036 | 0.103 | 0.010 | 0.050 | 0.017 | 0.016 | 0.117 | 5.444 | - | - |
| point sources | 0.040 | 0.116 | 0.005 | 0.272 | 0.017 | 0.016 | 0.117 | 0.329 | - | - |

3.3.1.2 Non-ferrous metals

1A2b, non-ferrous metal sector, includes emissions from grey iron foundries, lead, zinc, copper and secondary aluminium production. In particular, emission from the production of lead and zinc at the moment

are entirely reported in the energy sector because up to now there was no information to distinguish between energy and process emissions and, above all, these processes are considered combustion processes with contact, consequently, emissions are dependent on the combustion process. In particular, in Italy no production of primary copper has ever occurred while, as regards lead and zinc, there is a sole integrated plant for the primary productions, and this makes it difficult to ensure a good breakdown. Consequently, the issue related to the allocation of emissions is not only about combustion and process but also about the different productions of different metals in the same factory. To resolve this issue, an in-depth investigation has been started with the aim to better specify the technology used on the basis of E-PRTR and IPPC permits. The first result of this investigation has been the update of certain EFs since 2014 (ISPRA, 2021). HCB emission factors available in the Guidebook refer to the consumption of coal and other solid fuels and wood biomass while in Italy only natural gas and small amount of LPG and fuel oil are used so that HCB and PCB emissions from secondary aluminum production have been estimated and reported based on the information available in a national study. Dioxin emissions from this category is driven by emissions from secondary aluminum production where country specific emission factors are used from a research project of 2002 based on measurements at production plant level; such emissions are due prevalently to the role played by recycled material and there is no evidence of changes in the quality of the aluminum scraps as well as in the pretreatment process. Average emission factor is equal to 69 micrograms per Mg, and it is in the range of values of the Guidebook reported in the IPPU relevant sector but representing total emissions from this category. Dioxins emissions reported in 1A2b occur also for secondary production of lead, zinc and copper but in total their emissions are 10% of the total of 1A2b, one order of magnitude lower than those from aluminum production. An investigation is ongoing with the aim to report these emissions separately in the energy and IPPU sectors. Because emissions are up to now reported in the 1.A.2.b category, notation key IE has been provided for 2.C.3 category.

For Hg emission factors are from EMEP/CORINAIR 2007 Guidebook. Moreover, up to 2013, for primary and secondary lead production, emission factors for SO_x, NO_x, NMVOC, CO, Pb, PM₁₀ are country specific, from a sectoral technical survey (ENEA, 2000) and from the communication of the operators, as well as for PAH e dioxins (ENEA-AIB-MATTM, 2002). For the other pollutants emission factors are from EMEP/CORINAIR 2007 but they have been shared and checked with the main operator. For primary zinc production, up to 2013, SO_x, CO, Pb, PM, Zn and Cd emission factors are country specific as provided by the only operator while for the other pollutants are from the EMEP/CORINAIR Guidebook 2007 taking in account the weight of the different production processes, electrolytic and Imperial Smelting Furnace. Thanks to the investigation above mentioned emission factors for NO_x, SO_x, PM₁₀, Pb, Cd e Zn from zinc and lead production have been updated on the basis of data at plant level. For secondary aluminium production PAH and dioxins country specific emission factors have been used (ENEA-AIB-MATTM, 2002).

In the following Table 3.6 the main emission factors for the sector are reported. The data shown in the light blue cells are country specific based on emission data at plant level communicated in the framework of EPRTR, LCP, ETS or specific studies. PM₁₀ emission factors too are country specific while PM_{2.5} emission factors have been derived from PM₁₀ emission factors and the ratio PM_{2.5}/PM₁₀ per fuel as reported in the EMEP/EEA Guidebook 2019. The remaining emission factors derive from the EMEP/EEA Guidebook.

Table 3.6 *Main emission factors 1A2 Non-Ferrous metals - 2020*

| | SO _x | NO _x | NMVOC | CO | Cd | Hg | Pb | PM _{2.5} | PAH | Dioxins |
|---------------------------|-----------------|-----------------|-------|-------|--------------|-------|-------|-------------------|---------------|---------|
| | Kg/Mg product | | | | g/Mg product | | | | mg/Mg product | |
| Grey iron foundries | 0.125 | 0.160 | 0.090 | 9.500 | 0.140 | | 7.200 | 423.529 | | |
| Primary lead production | 0.758 | 0.163 | 0.045 | 0.085 | 0.159 | 3.000 | 0.898 | 26.337 | | |
| Primary zinc production | 0.088 | | | 0.598 | 0.140 | 6.120 | 0.635 | 18.989 | | |
| Primary copper production | | | | | | | | | | |

| | | | | | | | | | | |
|----------------------------------|-------|-------|-------|-------|-------|-------|--------|---------|-------|-------|
| Secondary lead production | 2.452 | 0.631 | 0.006 | 0.013 | | | 7.971 | 20.769 | 0.006 | 0.005 |
| Secondary zinc production | | | 1.000 | | 0.140 | 0.020 | 0.635 | | | 0.065 |
| Secondary copper production | | 0.200 | 2.000 | | 5.000 | | 50.000 | 967.989 | | 0.020 |
| Secondary aluminium production | 1.300 | 0.400 | 1.250 | | | | | 258.008 | 0.189 | 0.068 |
| Other | | | | | | | | | | |
| copper manufactures | | 0.200 | 2.000 | | | | | | | |
| zinc-copper & brass manufactures | | 0.200 | 2.000 | | | | | | | |

3.3.1.3 Chemicals

Category 1A2c refer to the stationary combustion emissions from the chemical and petrochemical sectors. Emissions from this category are calculated on the basis of fuel consumption and the relevant emission factors and they are due to the auto production of energy and heat for the production processes. Natural gas and petrochemical gases are prevalently used in gas turbine plants. It is not level and trend key category for any pollutant. Emission from the production processes are reported in the relevant IPPU sub sector. SO_x, NO_x and PM10 emission factors are country specific and derive from the communications of the main operators in the framework of the EPRTR and LCP framework while fuel consumption at plant level are provided in the ETS together with CO₂ emissions.

3.3.1.4 Pulp, paper and print

Category 1A2d refer to the stationary combustion emissions from the pulp and paper sector. Emissions from this category are calculated on the basis of fuel consumption and the relevant emission factors and they are due to the auto production of energy and heat for the production processes. Natural gas is prevalently used in gas turbine plants. It is not level and trend key category for any pollutant. Emission from the production processes are reported in the relevant IPPU sub sector. SO_x, NO_x and PM10 emission factors are country specific and derive from the communications of the main operators in the framework of the EPRTR and LCP framework while fuel consumption at plant level are provided in the ETS together with CO₂ emissions.

3.3.1.5 Food processing, beverages and tobacco

Category 1A2e refer to the stationary combustion emissions from the food and drink sector. Emissions from this category are calculated on the basis of fuel consumption and the relevant emission factors and they are due to the auto production of energy and heat for the production processes. Natural gas, and biogas are prevalently used in gas turbine plants. It is not level and trend key category for any pollutant. Emission from the production processes are reported in the relevant IPPU sub sector. SO_x, NO_x and PM10 emission factors are country specific and derive from the communications of the main operators in the framework of the EPRTR and LCP framework while fuel consumption at plant level are provided in the ETS together with CO₂ emissions.

3.3.1.6 Non-metallic minerals

Category 1A2f, stationary combustion in non-metallic mineral industry, refers to a multitude of production activities such as cement, lime, glass, brick and tiles, ceramics, and asphalt production which means a multitude of different emission factors. For cement production, PM emissions from kilns are reported in this category where emissions from mills are reported in IPPU (emission factor from USEPA 1991 emission factor handbook) while for lime production PM emission factors referring to the complete process are used (from USEPA 1996 emission factor handbook) and emissions are distributed between energy and IPPU. For Hg, emission factors are country specific (especially cement production which is the emission driver of this category); for Dioxin, HCB, PCB and Cd emission factors are from the relevant Bref reports or EMEP/EEA 2007 Guidebook; for Pb, emission factors are country specific for ceramic production and from the Bref report or EMEP 2006 Guidebook for glass, cement and lime productions.

In the following Table 3.7 the main emission factors for the sector are reported. The data shown in the light blue cells are country specific based on emission data at plant level communicated in the framework of EPRTR, LCP, ETS or specific studies. PM10 emission factors too are country specific while PM2.5 emission factors have been derived from PM10 emission factors and the ratio PM2.5/PM10 per fuel as reported in the EMEP/EEA Guidebook 2019. The remaining emission factors derive from the EMEP/EEA Guidebook.

Table 3.7 Main emission factors 1A2 Non-Metallic minerals - 2020

| | SO _x | NO _x | NM VOC | CO | Cd | Hg | Pb | PM _{2.5} | PAH | Dioxins |
|-------------------------------|-----------------|-----------------|--------|-------|--------------|-------|-------|-------------------|--------|---------------|
| | Kg/Mg product | | | | g/Mg product | | | | | mg/Mg product |
| Plaster furnaces | 0.115 | 0.088 | 0.009 | 0.013 | | | | 0.990 | | |
| Cement | 0.309 | 0.723 | 0.023 | 0.932 | 0.008 | 0.030 | 0.006 | 2.183 | | 0.0001 |
| Lime | | | | | | | | | | |
| areal sources | 0.098 | 1.197 | 0.004 | 0.013 | | 0.001 | | 1.200 | | |
| point sources | 0.004 | 0.023 | 0.004 | 0.013 | | 0.001 | | 0.600 | | |
| Asphalt concrete plants | 0.002 | 0.011 | 0.004 | 0.002 | | | | 6.500 | 0.0004 | |
| Flat glass | 1.010 | 2.810 | 0.044 | 0.100 | | | | 350.000 | | |
| Container glass | 1.156 | 1.618 | 0.047 | 0.100 | 0.090 | | | 448.800 | | |
| Glass wool (except binding) | 2.500 | 1.480 | 0.048 | 0.081 | | | | 123.390 | | |
| Other glass | | 2.000 | 0.100 | 0.260 | | | | 17.778 | | |
| Mineral wool (except binding) | 1.500 | | | 3.200 | | | | 493.559 | | |
| Bricks and tiles | 0.785 | 0.250 | 0.007 | 0.040 | | | | 49.841 | | |
| Fine ceramic materials | 0.569 | 0.174 | 0.014 | 0.130 | | | | 193.898 | | |

3.3.1.7 Other

Category 1A2g refer to the stationary combustion emissions from different industrial sectors as machinery, mining and quarrying, construction, textile and others. Emissions from this category are calculated on the basis of fuel consumption and the relevant emission factors and they are due to the auto production of energy and heat for the production processes. Natural gas is prevalently used in gas turbine plants. Fugitive emissions from the oil and gas extraction activities are reported in the relevant energy sub sector. SO_x, NO_x and PM10 emission factors are country specific and derive from the communications of the main operators in the framework of the EPRTR and LCP framework while fuel consumption at plant level are provided in the ETS together with CO₂ emissions. In general it is not level and trend key category for any pollutant. In 2020 this category is key category for SO_x because of a new oil and gas extraction plant which had in the same year different accidental events.

3.3.2 Time series and key categories

Manufacturing industries and construction (1A2) is a main source of heavy metals and POPs, accounting for about 43% of lead total emissions, 30% for cadmium, 28% for mercury, 42% for HCB, and 17% for dioxin. The sector is key category also for PM10 and PM2.5 (4%) as well as SO_x, NO_x and CO, about 39%, 8% and 5% of total emissions. In 2020 the main sectors are *Stationary combustion in iron and steel industries* (1A2a), which is key for SO_x, CO, Cd, and Hg, *Stationary combustion in non-ferrous metal industries* (1A2b), key for Hg, Dioxin and HCB, and *stationary combustion in non-metallic mineral industries* (1A2f) that is key category for SO_x, NO_x, PM10, PM2.5, Pb, Cd and Hg.

As already reported for 1A1 category, a strong reduction of SO_x, NO_x and PM10 emissions is observed for this category along the time series. The introduction of two regulatory instruments: the DPR 203/88 (Decree of President of the Republic of 24th May 1988), laying down rules concerning the authorisation of plants, and the Ministerial Decree of 12th July 1990, which introduced plant level limits to emissions of PM10, NO_x and SO_x for new plants and required old plants to conform to the limit by 1997, explained the emission reduction in the nineties. The shift from fuel oil to natural gas combined with the increase of energy efficiency of the plants and the introduction of PM10 abatement technologies have been implemented to comply with the emission limit values. From 2000 lower limits to emissions at the stacks have been introduced, in the framework of environmental integrated authorisations, for the authorisation of new plants and the implementation of the old ones, especially for those facilities located in areas with air quality critical values. For this reason, the plants have increased the use of natural gas heat and power combined technology.

3.3.2.1 Iron and steel

The category 1A2a in 2020 is a key category for SO_x, CO, Cd, and Hg. The trend of emissions is linked above all to the production levels and then to the abatement technologies.

In fact, for the majority of pollutants, in 2009 a strong reduction of emissions is observed due to the effects of the economic recession that in 2010 and 2011 has partially recovered. In 2012 a further drop occurred for the economic crisis and for environmental constrains of the main iron and steel integrated plants that should reduce its productions. In 2015 a drop is still observed consistently with the production activities reduction of the main iron and steel integrated plants. There were four integrated steel plants in 1990 that from 2005 are reduced to two, with another plant that still has a limited production of pig iron. Nevertheless, the steel production in integrated plants has not changed significantly in the 1990-2008 period due to an expansion in capacity of the two operating plants. From 2015 only one integrated plant remains in operation. The maximum production was around 11 Mt/y in 1995 and in 2005-2008, with lower values in other years and the lowest of 3.4 Mt in 2020.

3.3.2.2 Non-ferrous metals

The category 1A2b in 2020 is a key category for Hg, dioxins and HCB.

In Italy, the production of primary aluminium stopped in 2013 (and was 232 Gg in 1990) while secondary aluminium accounts for 350 Gg in 1990 and 698 Gg in 2020. These productions, however, use electricity as the primary energy source so the emissions due to the direct use of fossil fuels are limited. The sub sector

comprises also the production of other non-ferrous metals, both primary and secondary copper, lead, zinc and others; but also those productions have a limited share of emissions.

3.3.2.3 Chemicals

In Italy there are petrochemical plants integrated with a nearby refinery and stand-alone plants that get the inputs from the market. Main products are Ethylene, Propylene, Styrene. In particular, ethylene and propylene are produced in petrochemical industry by steam cracking. Ethylene is used to manufacture ethylene oxide, styrene monomer and polyethylene. Propylene is used to manufacture polypropylene but also acetone and phenol. Styrene, also known as vinyl benzene, is produced on industrial scale by catalytic dehydrogenation of ethyl benzene. Styrene is used in the rubber and plastic industry to manufacture through polymerisation processes such products as polystyrene, ABS, SBR rubber, SBR latex. Except for ethylene oxide, whose production has stopped in 2002, the other productions of the above-mentioned chemicals still occur in Italy. Activity data are stable from 1990 to 2012, with limited yearly variations along the timeseries and a reduction in the last years. Chemical industry includes non-organic chemicals as chlorine/soda, sulphuric acid, nitric acid, ammonia. A limited production of fertilizers is also present in Italy. From 1990 to 2020 the sum of productions of this source category has greatly reduced: in 2020 it was about a half of the production in 1990.

3.3.2.4 Pulp, paper and print

Emissions from the manufacturing of paper are included in this source category. In Italy the manufacture of virgin paper pulp is rather limited, with a production feeding less than 5% of the paper produced in 2020. Most of the pulp was imported in 1990, while in 2020 half of the pulp used is produced locally from recycled paper. The paper production is expanding and activity data (total paper produced) were 6.2 Mt in 1990 and 8.9 Mt in 2020. The printing industry represents a minor part of the source category emissions.

3.3.2.5 Food processing, beverages and tobacco

In Italy the food production industry is expanding. A comprehensive activity data for this sector is not available; more in detail while energy data are those reported in the national energy balance for this sector, information at subsector and technological level is not available and only few plants are part of the ETS; energy fuel consumption was estimated to be 62 PJ in 1990 and 113 PJ in 2020, about half of energy consumptions derives from biomass. Value added at constant prices has increased of 0.6% per years from 1990 to 2003 and almost constant from 2004.

3.3.2.6 Non-metallic minerals

The category 1A2f in 2020 is a key category for SO_x, NO_x, PM₁₀, PM_{2.5}, Pb, Cd and Hg. This sector, which refers to construction materials, is quite significant in terms of emissions due to the energy intensity of the processes involved. Construction materials subsector includes the production of cement, lime, bricks, tiles and glass. It comprises thousands of small and medium size enterprises, with only few large operators, mainly connected to cement production. Some of the production is also exported.

3.3.2.7 Other

This sector comprises emissions from many different industrial subsectors, some of which are quite significant in Italy in terms of both value added and export capacity. In particular, engineering sectors (vehicles and machines manufacturing) is the main industrial sub sector in terms of value added and revenues from export and textiles was the second subsector up to year 2000. The remaining “other industries” include furniture and other various “made in Italy” products that produce not negligible amounts of emissions. In 2020 this category is key category for SO_x because of a new oil and gas extraction plant which had in the same year different accidental events.

3.3.3 QA/QC and verification

As already reported, QA/QC checks at plants level are directed to check the submissions of data in the different context and evaluate the differences if any. For example, if emissions submitted by a plant under LCP are higher than those submitted under the EPRTR ISPRA asks the operator of the reporting plant for an explanation and the verification of data submitted. In addition, on the basis of fuel consumption supplied under the ETS and average emission factor by fuel energy experts estimate emissions at plant level and compare them with

those submitted in the EPRTR and LCP. Also, in this case ISPRA asks for clarifications to the reporting plant if necessary. As for 1A1 point sources, the analysis of data collected from point sources allowed to distribute emissions at local level and to check and verify data at local level.

3.3.4 Recalculations

For 1.A.2 category several recalculations occurred in this submission, most of them as a follow up to the review process.

In particular for 1.A.2.a, iron and steel activities, SO_x, NO_x, PM and HMs emission factors since 2016, as well as HCB emission factors since 2010, have been updated on the basis of plant level emissions data.

For 1.A.2.b, secondary aluminum production, HCB and PCB emission have been estimated for the whole time series on the basis of country specific emission factors.

For 1.A.2.f, according to the EMEP/EEA Guidebook, SO_x emissions from cement production before reported in the IPPU sector have been moved and reported under the energy sector. Moreover further improvement and recalculations have been occurred for SO_x, NO_x, and NH₃ emissions from glass production from 2007 and for NO_x, from lime production from 1990, because of the use of E-PRTR data. Lime and other glass activity data for 2019 have been updated.

3.3.5 Planned improvements

Specific improvements are detailed in the 2022 QA/QC plan (ISPRA, 2022[b]).

For the *energy* sector, a major progress regards the management of the information system where data collected in the framework of different obligations, Large Combustion Plant, E-PRTR and Emissions Trading, are gathered together thus highlighting the main discrepancies in information and detecting potential errors. Moreover, the complete use of the energy data provided by the Ministry of Economic Development to the Joint Questionnaire IEA/OECD/EUROSTAT is planned in substitution of the national energy balances used till now; liquid, gaseous and solid fuel are now aligned for the whole time series and we plan for the next submission to update as possible renewable fuels and biomass.

3.4 AVIATION (NFR SUBSECTOR 1.A.3.A)

3.4.1 Overview

Emissions from categories 1.A.3.a.i International Aviation and 1.A.3.a.ii Domestic Aviation are estimated, including figures both for landing and take-off cycles (LTO) and for the cruise phase of the flight (the latter reported as memo items and not included in the national totals).

3.4.2 Methodological issues

According to the IPCC Guidelines and Good Practice Guidance (IPCC, 1997; IPCC, 2006; IPCC, 2000) and the EMEP/EEA Guidebook 2016 (EMEP/EEA, 2019), a national technique has been developed and applied to estimate emissions.

The current method estimates emissions from the following assumptions and information.

Activity data comprise both fuel consumptions and aircraft movements, which are available in different level of aggregation and derive from different sources as specified here below:

- Total inland deliveries of aviation gasoline and jet fuel are provided in the national energy balance (MSE, several years (a)). This figure is the best approximation of aviation fuel consumption, for international and domestic use, but it is reported as a total and not split between domestic and international.
- Data on annual arrivals and departures of domestic and international landing and take-off cycles at Italian airports are reported by different sources: National Institute of Statistics in the statistics yearbooks (ISTAT, several years), Ministry of Transport in the national transport statistics yearbooks (MIMS, several years), the Italian civil aviation in the national aviation statistics yearbooks (ENAC/MIMS, several years), EUROCONTROL flights data time series 2002–2018 (EUROCONTROL, several years).

An overall assessment and comparison with EUROCONTROL emission estimates was carried out over the years and that lead to an update of the methodology used by Italy for this category. Data on the number of flights, fuel consumption and emission factors were provided by EUROCONTROL in the framework of a specific project funded by the European Commission, and quality checked by the European Environmental Agency and its relevant Topic Centre (ETC/ACM), aimed at improving the reporting and the quality of emission estimates from the aviation sector of each EU Member State under both the UNFCCC and LRTAP conventions. The Advanced Emissions Model (AEM) was applied by EUROCONTROL to derive these figures, according to a Tier 3 methodology (EMEP/EEA, 2019).

EUROCONTROL fuel and emissions time series cover the period 2005-2020, while the number of flights is available since 2002. In this submission, recalculation of the time series affected PAH from 1990 to 2017.

For the time series from 1990 to 1999, figures for emission and consumption factors are derived by the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007), both for LTO cycles and cruise phases, taking into account national specificities.

These specificities derived from the results of a national study which, taking into account detailed information on the Italian air fleet and the origin-destination flights for the year 1999, calculated national values for both domestic and international flights (Romano et al., 1999; ANPA, 2001; Trozzi et al., 2002 (a)) on the basis of the default emission and consumption factors reported in the EMEP/CORINAIR guidebook. National average emissions and consumption factors were therefore estimated for LTO cycles and cruise both for domestic and international flights from 1990 to 1999. Specifically, for the year referred to in the survey, the method estimates emissions from the number of aircraft movements broken down by aircraft and engine type (derived from ICAO database if not specified) at each of the principal Italian airports; information about whether the flight is international or domestic and the related distance travelled has also been considered. A Tier 3 method has been applied for 1999. In fact, figures on the number of flights, destination, aircraft fleet and engines have been provided by the local airport authorities, national airlines and EUROCONTROL, covering about 80% of the national official statistics on aircraft movements for the relevant years. Data on 'Times in mode' have also been supplied by the four principal airports and estimates for the other minor airports have been carried out on the basis of previous sectoral studies at local level. Consumption and emission factors are those derived from the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007). Based on sample information, estimates have been carried out at national level from 1990 to 1999 considering the official statistics of the aviation sector (ENAC/MIMS, several years) and applying the average consumption and emission factors.

From 2005, fuel consumption and emission factors were derived from the database made available to EU Member States by EUROCONTROL, as previously described. These data were used for updating fuel

consumption factors, and emission factors of all pollutants. For the period between 1999 and 2005, interpolation has been applied to calculate these parameters. Estimates were carried out applying the consumption and emission factors to the national official aviation statistics (ENAC/MIMS, several years) and EUROCONTROL data on movements from 2002 (EUROCONTROL, several years).

In general, to carry out national estimates of greenhouse gases and other pollutants for LTO cycles, both domestic and international, consumptions and emissions are calculated for the complete time series using the average consumption and emission factors multiplied by the total number of flights. The same method is used to estimate emissions for domestic cruise; on the other hand, for international cruise, consumptions are derived by difference from the total fuel consumption reported in the national energy balance and the estimated values as described above and emissions are therefore calculated.

The fuel split between national and international fuel use in aviation is then supplied to the Ministry of the Economic Development to be included in the official international submission of energy statistics to the IEA in the framework of the Joint Questionnaire OECD/EUROSTAT/IEA compilation together with other energy data.

Data on domestic and international aircraft movements from 1990 to 2020 are shown in Table 3.8 where domestic flights are those entirely within Italy.

Since 2002, EUROCONTROL flights data have been considered, accounting for departures from and arrivals to all airports in Italy, regarding flights flying under instrument flight rules (IFR), including civil helicopters flights and excluding flights flagged as military, when the above flights can be identified.

Total fuel consumptions, both domestic and international, are reported by LTO and cruise in Table 3.9.

Table 3.8 Aircraft Movement Data (LTO cycles)

| Number of flights | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|-----------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| Domestic flights | 172,148 | 185,220 | 319,748 | 350,140 | 354,520 | 280,645 | 277,872 | 281,498 | 284,627 | 288,470 | 151,156 |
| International flights | 147,875 | 198,848 | 303,608 | 381,206 | 406,990 | 425,410 | 446,817 | 462,896 | 484,764 | 502,764 | 172,835 |

Source: ISTAT, several years; ENAC/MIMS, several years; Eurocontrol, several years.

Table 3.9 Aviation jet fuel consumptions for domestic and international flights (Gg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|----------------------|-----------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| | Gg | | | | | | | | | | |
| Domestic LTO | 111 | 120 | 208 | 233 | 227 | 168 | 166 | 169 | 179 | 180 | 89 |
| International LTO | 130 | 175 | 258 | 269 | 296 | 328 | 344 | 355 | 382 | 399 | 130 |
| Domestic cruise | 357 | 384 | 654 | 666 | 704 | 526 | 527 | 544 | 563 | 580 | 291 |
| International cruise | 1,246 | 1,688 | 2,297 | 2,456 | 2,534 | 2,745 | 2,962 | 3,230 | 3,461 | 3,584 | 1,087 |

Source: ISPRA elaborations

Emissions from military aircrafts are also estimated and reported under category 1.A.5 Other. The methodology to estimate military aviation emissions is simpler than the one described for civil aviation since LTO data are not available in this case. As for activity data, total consumption for military aviation is published in the petrochemical bulletin (MSE, several years (b)) by fuel. Emission factors are those provided in the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007). Therefore, emissions are calculated by multiplying military fuel consumption data for the EMEP/CORINAIR default emission factors.

3.4.3 Time series and key categories

Emission time series of NO_x, NMVOC, SO_x, TSP, CO, Pb are reported in Table 3.10, Table 3.11, Table 3.12, Table 3.13, Table 3.14 and Table 3.15, respectively.

An upward trend in emission levels for civil aviation is observed from 1990 to 2019, which is explained by the increasing number of LTO cycles. Nevertheless, the propagation of more modern aircrafts in the fleet slows down the trend in the most recent years. There has also been a decrease in the number of domestic flights from 2000, although a new increasing trend in the last couple of years has been registered. Year 2020 is to be considered separately, because the aviation sector was severely hit by the pandemic measures: domestic flights in 2020 were about the 52% of the 2019, while international flights during 2020 were the 34% of the flights of the previous year. Aviation is not a key category.

Table 3.10 *Time series of NO_x (Gg)*

| Source categories for NFR Subsector 1.A.3.a, 1.A.5.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|--------------|--------------|-------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Gg | | | | | | | | | | | |
| 1 A 3 a ii (i) Domestic aviation LTO (civil) | 1.36 | 1.47 | 2.50 | 2.55 | 2.71 | 2.12 | 2.11 | 2.14 | 2.30 | 2.32 | 1.15 |
| 1 A 3 a i (i) International aviation LTO (civil) | 1.60 | 2.16 | 3.20 | 3.47 | 3.99 | 4.55 | 4.84 | 5.02 | 5.35 | 5.60 | 1.85 |
| 1 A 3 a Civil Aviation (LTO) | 2.97 | 3.62 | 5.70 | 6.02 | 6.70 | 6.68 | 6.96 | 7.16 | 7.65 | 7.92 | 3.01 |
| 1A3 a ii (ii) Domestic aviation cruise (civil) | 5.23 | 5.63 | 9.43 | 8.71 | 10.16 | 8.09 | 8.10 | 8.21 | 8.51 | 8.78 | 4.25 |
| 1A3a i (ii) International aviation cruise (civil) | 18.85 | 26.83 | 38.99 | 36.55 | 41.02 | 47.05 | 50.25 | 52.04 | 55.71 | 59.65 | 18.56 |
| 1 A 5 b Other, Mobile (including military, land based and recreational boats) | 11.16 | 11.99 | 7.24 | 13.50 | 6.11 | 3.29 | 3.28 | 2.36 | 2.05 | 2.73 | 3.48 |

Table 3.11 *Time series of NMVOC (Gg)*

| Source categories for NFR Subsector 1.A.3.a, 1.A.5.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Gg | | | | | | | | | | | |
| 1 A 3 a ii (i) Domestic aviation LTO (civil) | 0.12 | 0.13 | 0.23 | 0.25 | 0.33 | 0.27 | 0.26 | 0.26 | 0.28 | 0.30 | 0.14 |
| 1 A 3 a i (i) International aviation LTO (civil) | 0.15 | 0.20 | 0.31 | 0.39 | 0.45 | 0.48 | 0.49 | 0.48 | 0.51 | 0.52 | 0.16 |
| 1 A 3 a Civil Aviation (LTO) | 0.27 | 0.33 | 0.54 | 0.64 | 0.78 | 0.75 | 0.75 | 0.75 | 0.79 | 0.82 | 0.31 |
| 1A3 a ii (ii) Domestic aviation cruise (civil) | 0.10 | 0.10 | 0.18 | 0.20 | 0.37 | 0.34 | 0.34 | 0.34 | 0.36 | 0.38 | 0.19 |
| 1A3a i (ii) International aviation cruise (civil) | 0.25 | 0.36 | 0.55 | 0.69 | 0.81 | 0.89 | 0.92 | 0.92 | 0.93 | 0.94 | 0.29 |
| 1 A 5 b Other, Mobile (including military, land based and recreational boats) | 3.00 | 3.13 | 1.90 | 3.00 | 1.05 | 0.66 | 0.70 | 0.48 | 0.44 | 0.58 | 0.77 |

Table 3.12 *Time series of SO_x (Gg)*

| Source categories for NFR Subsector 1.A.3.a, 1.A.5.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Gg | | | | | | | | | | | |
| 1 A 3 a ii (i) Domestic aviation LTO (civil) | 0.11 | 0.12 | 0.21 | 0.23 | 0.23 | 0.17 | 0.17 | 0.17 | 0.18 | 0.18 | 0.09 |
| 1 A 3 a i (i) International aviation LTO (civil) | 0.13 | 0.17 | 0.26 | 0.27 | 0.30 | 0.33 | 0.34 | 0.35 | 0.38 | 0.40 | 0.13 |
| 1 A 3 a Civil Aviation (LTO) | 0.24 | 0.29 | 0.47 | 0.50 | 0.52 | 0.50 | 0.51 | 0.52 | 0.56 | 0.58 | 0.22 |
| 1A3 a ii (ii) Domestic aviation cruise (civil) | 0.36 | 0.38 | 0.65 | 0.67 | 0.70 | 0.53 | 0.53 | 0.54 | 0.56 | 0.58 | 0.29 |
| 1A3a i (ii) International aviation cruise (civil) | 1.25 | 1.78 | 2.60 | 2.45 | 2.65 | 2.95 | 3.11 | 3.24 | 3.48 | 3.69 | 1.16 |
| 1 A 5 b Other, Mobile (including military, land based and recreational boats) | 1.19 | 0.81 | 0.21 | 0.17 | 0.13 | 0.12 | 0.15 | 0.08 | 0.10 | 0.13 | 0.19 |

Table 3.13 *Time series of TSP (Gg)*

| Source categories for NFR Subsector 1.A.3.a, 1.A.5.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Gg | | | | | | | | | | | |
| 1 A 3 a ii (i) Domestic aviation LTO (civil) | 0.01 | 0.01 | 0.02 | 0.03 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.01 |
| 1 A 3 a i (i) International aviation LTO (civil) | 0.02 | 0.03 | 0.04 | 0.05 | 0.03 | 0.04 | 0.04 | 0.04 | 0.04 | 0.04 | 0.01 |
| 1 A 3 a Civil Aviation (LTO) | 0.03 | 0.04 | 0.06 | 0.07 | 0.06 | 0.05 | 0.05 | 0.05 | 0.06 | 0.06 | 0.02 |
| 1A3 a ii (ii) Domestic aviation cruise (civil) | 0.07 | 0.08 | 0.13 | 0.10 | 0.10 | 0.07 | 0.07 | 0.07 | 0.08 | 0.08 | 0.04 |
| 1A3a i (ii) International aviation cruise (civil) | 0.36 | 0.52 | 0.75 | 0.71 | 0.83 | 0.91 | 0.94 | 0.94 | 0.95 | 0.96 | 0.30 |
| 1 A 5 b Other, Mobile (including military, land based and recreational boats) | 1.30 | 1.57 | 0.91 | 1.63 | 0.83 | 0.48 | 0.50 | 0.34 | 0.32 | 0.43 | 0.56 |

Table 3.14 *Time series of CO (Gg)*

| Source categories for NFR Subsector 1.A.3.a, 1.A.5.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| | Gg | | | | | | | | | | |
| 1 A 3 a ii (i) Domestic aviation LTO (civil) | 1.23 | 1.33 | 2.26 | 2.33 | 2.28 | 1.69 | 1.65 | 1.71 | 1.85 | 1.88 | 0.94 |
| 1 A 3 a i (i) International aviation LTO (civil) | 1.73 | 2.32 | 3.33 | 2.86 | 3.00 | 3.25 | 3.31 | 3.35 | 3.73 | 3.89 | 1.25 |
| 1 A 3 a Civil Aviation (LTO) | 2.96 | 3.64 | 5.59 | 5.19 | 5.28 | 4.94 | 4.95 | 5.06 | 5.57 | 5.77 | 2.19 |
| 1A3 a ii (ii) Domestic aviation cruise (civil) | 1.31 | 1.41 | 2.43 | 2.66 | 3.07 | 2.35 | 2.32 | 2.42 | 2.51 | 2.60 | 1.47 |
| 1A3a i (ii) International aviation cruise (civil) | 2.03 | 2.89 | 4.42 | 5.55 | 5.74 | 6.07 | 6.29 | 6.45 | 6.98 | 7.23 | 2.49 |
| 1 A 5 b Other, Mobile (including military, land based and recreational boats) | 65.12 | 79.02 | 45.49 | 54.48 | 17.33 | 16.49 | 19.73 | 11.93 | 13.23 | 17.42 | 24.93 |

Table 3.15 *Time series of Pb (Mg)*

| Source categories for NFR Subsector 1.A.3.a, 1.A.5.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|--------------|-------------|-------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | Gg | | | | | | | | | | |
| 1 A 3 a ii (i) Domestic aviation LTO (civil) | 0.19 | 0.20 | 0.35 | 0.38 | 0.38 | 0.30 | 0.30 | 0.30 | 0.31 | 0.31 | 0.16 |
| 1 A 3 a i (i) International aviation LTO (civil) | 0.21 | 0.28 | 0.43 | 0.54 | 0.57 | 0.60 | 0.63 | 0.65 | 0.68 | 0.71 | 0.24 |
| 0.24 1 A 3 a Civil Aviation (LTO) | 0.39 | 0.48 | 0.77 | 0.91 | 0.96 | 0.90 | 0.93 | 0.96 | 0.99 | 1.02 | 0.41 |
| 1A3 a ii (ii) Domestic aviation cruise (civil) | 0.57 | 0.62 | 1.06 | 1.16 | 1.18 | 0.93 | 0.92 | 0.94 | 0.95 | 0.96 | 0.50 |
| 1A3a i (ii) International aviation cruise (civil) | 2.01 | 2.86 | 4.36 | 5.48 | 5.85 | 6.11 | 6.42 | 6.65 | 6.96 | 7.22 | 2.48 |
| 1 A 5 b Other, Mobile (including military, land based and recreational boats) | 16.34 | 4.22 | 1.16 | 0.001 | NA | 0.12 | 0.02 | 0.02 | 0.09 | 0.18 | 0.18 |

3.4.4 QA/QC and Uncertainty

Data used for estimating emissions from the aviation sector derive from different sources: local airport authorities, national airlines operators, EUROCONTROL and official statistics by different Ministries and national authorities.

Different QA/QC and verification activities are carried out for this category. As regards past years, the results of the national studies and methodologies, applied at national and airport level, were shared with national experts in the framework of an *ad hoc* working group on air emissions instituted by the National Aviation Authority (ENAC). The group, chaired by ISPRA, included participants from ENAC, Ministry of Environment, Land and Sea, Ministry of Transport, national airlines and local airport authorities. The results reflected differences between airports, aircrafts used and times in mode spent for each operation.

Currently, verification and comparison activities regard activity data and emission factors. In particular, number of flights have been compared considering different sources: ENAC, ASSAEROPORTI, ISTAT, EUROCONTROL and verification activities have been performed on the basis of the updated EUROCONTROL data on fuel consumption and emission factors resulting in an update and improving of the national inventory. Furthermore, there is an ongoing collaboration and data exchange with regional environmental agencies on this issue.

3.4.5 Recalculations

In this submission recalculation of the time series affected PAH from 1990 to 2017 due to the correction of an error in the calculation of the emissions.

3.4.6 Planned improvements

Improvements for next submissions are planned on the basis of the outcome of the ongoing quality assurance and quality control activities, in particular with regard to the results of investigation about data and information deriving from different sources, in particular further assessment of EUROCONTROL data, and comparison with information provided by the national institute of statistics, ISTAT, on the number of flights.

3.5 ROAD TRANSPORT (NFR SUBSECTOR 1.A.3.B)

3.5.1 Overview

The road transport sector contributes to the total national emissions in 2020 as follows: nitrogen oxides emissions for 37.4% of the total, emissions of carbon monoxide for 15.7%, non-methane volatile organic compounds for 10.4%, PM10 and PM2.5, for 9.3% and 8.0%, respectively, of the total.

The estimation refers to the following vehicle categories:

- 1.A.3.b.i Passenger cars
- 1.A.3.b.ii Light-duty trucks
- 1.A.3.b.iii Heavy-duty vehicles including buses
- 1.A.3.b.iv Mopeds and motorcycles
- 1.A.3.b.v Gasoline evaporation
- 1.A.3.b.vi Road transport: Automobile tyre and brake wear
- 1.A.3.b.vii Road transport: Automobile road abrasion

3.5.2 Methodological issues

A national methodology has been developed and applied to estimate emissions according to the IPCC Guidelines and Good Practice Guidance (IPCC, 1997; IPCC, 2000; IPCC, 2006) and the EMEP/EEA Guidebook (EMEP/EEA, 2019).

In general, the annual update of the model is based on the availability of new measurements and studies regarding road transport emissions (for further information see: <https://www.emisia.com/utilities/copert/>).

The model COPERT 5 (updated version 5.5.1, September 2021) has been used and applied for the whole time series in 2022 submission. COPERT 5 introduced over the years upgrades both from software and methodological point of view (<https://www.emisia.com/utilities/copert/versions/>).

Several new methodological features had been introduced respect to the previous model COPERT 4.

As regards fuel, updates concerned: fuel energy instead of fuel mass calculations; distinction between primary and end (blends) fuels, automated energy balance.

Regarding vehicle types, updated vehicle category naming, new vehicle types and emission control technology level, have been introduced.

As regards emission factors, one function type and the possibility to distinguish between peak/off-peak urban, have been implemented.

Main methodological innovations introduced since version 5.4.36, used in last submission, relate: introduction of Solid Particle Number (SPN23) emission factors for all vehicle categories; updated PM exhaust emission factor; the calculation of N₂O and NH₃ emissions for PC LPG & CNG and Urban Buses CNG has been corrected; NFR export has been corrected; the calculation of HM emissions has been corrected; fuel consumption from fossil part of biofuels is reported in the CRF export (sheet: "Other fossil fuels"); correction of NO_x emission factors for PC Diesel PHEV (Euro 6d-temp and 6d); hot emission factors for Quad & ATVs and Micro-cars have been corrected; furthermore various software upgrades have been implemented.

The model, on the basis of the inputs inserted, gives output results separately for vehicles category and urban (peak/off-peak urban), rural, highway areas, concerning emission estimates of CO, VOC, NMVOC, CH₄, NO_x, N₂O, NH₃, PM2.5, PM10, PM exhaust (the emission factors of particulate matter from combustion refer to particles smaller than 2.5 µm, that implicitly assumes that the fraction of particulate matter with diameter between 2.5 µm and 10 µm is negligible), CO₂, SO₂, heavy metals, NO_x speciation in NO and NO₂, the speciation in elemental and organic carbon of PM, the speciation of NMVOC.

Resulting national emission factors at detailed level are available on the following public web address: <http://www.sinanet.isprambiente.it/it/sia-ispra/fetransp>.

Data on fuel consumption of gasoline, diesel, liquefied petroleum gas (LPG), natural gas (CNG) and biofuels are those reported in the EUROSTAT energy balance; Italian road electricity consumption, introduced since 2021 submission, derives from Eurostat database (<https://ec.europa.eu/eurostat/data/database>). Time series of consumptions, by fuel and vehicle categories, are detailed in the NFR.

Lubricants consumption due to 2 stroke engines is estimated and reported in 1A3b. All the other national lubricants consumption, including 4 stroke engines, and relevant emissions are reported in 2D3 category.

3.5.2.1 Exhaust emissions

Exhaust emissions from vehicles subsectors are split between cold and hot emissions; estimates are calculated either on the basis of a combination of total fuel consumption and fuel properties data or on the basis of a combination of drive related emission factors and road traffic data.

The calculation of emissions is based on emission factors calculated for the vehicle models most widely and systematically used, distinguishing between the type of vehicle, fuel, engine size or weight class, standard legislation. The legislative standards introduced become more stringent over the years, ensuring that new vehicles emit much less than the older ones as regards the regulated pollutants.

With reference to four groups of pollutants, the method of calculation of exhaust emissions is different. The methodology implemented is derived from the EMEP/EEA Emission Inventory Guidebook 2019 (EMEP/EEA, 2019).

As regards the first two groups, methods are used leading to high standard detailed emissions data.

The first group includes: CO, NO_x, VOC, CH₄, NMVOC, N₂O, NH₃ and PM. For these pollutants, specific emission factors are applied relating to different engine conditions and urban, rural and highway driving shares.

The second group includes: CO₂, SO₂, Pb, Cd, Cr, Cu, Ni, Se, Zn. The emissions of these pollutants are estimated on the basis of fuel consumption.

For the third group of pollutants, including PAHs and PCDDs and PCDFs, detailed data are not available and then a simplified methodology is applied.

Finally the fourth group includes pollutants (alkanes, alkenes, alkynes, aldehydes, ketones, cycloalkanes and aromatic compounds) obtained as a fraction of the total emissions of NMVOC, assuming that the fraction of residual NMVOC are PAHs.

Because of the availability in Italy of an extensive and accurate database, a detailed methodology is implemented in the model COPERT 5. Total emissions are calculated as the sum of hot emissions, deriving from the engine when it reaches a hot temperature, and cold emissions produced during the heating process. The different methodological approach is justified by the performance of vehicles in the two different phases.

The production of emissions is also closely linked to the driving mode, differentiating for activity data and emission factors, with reference to urban (where it is assumed that almost all cold emissions are produced), rural and highway shares. Several factors contribute to the production of hot emissions such as mileage, speed, type of road, vehicle age, engine capacity and weight. Cold emissions are mainly attributed to urban share, and are attributed only to passenger cars and light duty vehicles. Varying according to the weather conditions and driving behaviour, are related to the specific country.

Emissions of NMVOC, NO_x, CO and PM are calculated on the basis of emission factors expressed in grams per kilometre and road traffic statistics estimated by ISPRA on account of data released from Ministry of Transport, ACI and ANCM (several years). The emission factors are based on experimental measurements of emissions from in-service vehicles of different types driven under test cycles with different average speeds calculated from the emission functions and speed-coefficients provided by COPERT 5 (EMISIA SA, 2021). This source provides emission functions and coefficients relating emission factors (in g/km) to average speed for each vehicle type and Euro emission standard derived by fitting experimental measurements to polynomial functions. These functions were then used to calculate emission factor values for each vehicle type and Euro emission standard at each of the average speeds of road and area types.

As regards the speciation of PM into elemental (EC, assumed to be equal to black carbon for road transport) and organic carbon (OC), considering the organic material (OM) as the mass of organic carbon corrected for the hydrogen content of the compounds collected, since the estimates are based on the assumption that low-sulphur fuels are used, when advanced after treatments are used, EC and OM do not add up to 100%, assuming that the remaining fraction consists of ash, nitrates, sulphates, water and ammonium salts (EMEP/EEA 2019).

Emissions of fuel dependent pollutants have been estimated applying a different approach.

Data on consumption of various fuels are derived from official statistics aggregated at national level and then estimated in the detail of vehicle categories, emission regulation and road type in Italy. The resulting error of approximation deriving from the comparison between the calculated value and the statistical value of the total fuel consumption, is corrected by applying a normalisation procedure to the breakdown of fuel consumption by each vehicle type calculated on the basis of the fuel consumption factors added up, with reference to the BEN figures for total fuel consumption in Italy (adjusted for off-road consumption).

The 1990-2020 inventory uses fuel consumption factors expressed as MJ per kilometre for each vehicle type and average speed calculated from the emission functions and speed-coefficients provided by the model COPERT 5, version 5.5.1. Emissions of sulphur dioxide and heavy metals are calculated applying specific

factors to consumption of gasoline, diesel, liquefied petroleum gas (LPG) and natural gas (CNG), taken from the BEN (MiSE, MiTE, several years (a)), updated since 2017 according to EUROSTAT methodology.

Emissions of SO₂ are based on the sulphur content of the fuel. Values for SO₂ vary annually as the sulphur-content of fuels change and are calculated every year for gasoline and gas oil and officially communicated to the European Commission in the framework of the European Directives on fuel quality; these figures are also published by the refineries industrial association (UP, several years).

Fuel specifications for gasoline, diesel fuel and LPG, derive from *ad hoc* studies about the properties of transportation fuels sold in Italy and whose results are representative and applicable with reference to three different time phases: 1990 – 1999; 2000 – 2012; 2013 – 2019; since 2020 (Innovhub – Fuel Experimental Station surveys, several years).

As regards natural gas, the national market is characterized by the commercialisation of gases with different chemical composition in variable quantities from one year to the other. Each year the quantities of natural gas imported or produced in Italy are published on the web by the MSE <http://dgerm.sviluppoeconomico.gov.it/dgerm/bilanciogas.asp>.

In Italy, biodiesel, biogasoline and biogas are used in road transportation and the respective emissions have been estimated in the Inventory.

As regards biodiesel and biogasoline, almost all the commercial gasoline is practically still substantially an E0 (in 2020 the share of biogasoline is 0.3%, respect to the total road gasoline consumption), while the distributed diesel reaches up to 5-7% by volume of biodiesel in diesel fuel (in 2020 the share of biodiesel is 6.7%, respect to the total road diesel consumption). That is because Italian producers/refineries have decided since the beginning of the introduction of the obligations on biofuels to focus on biodiesel rather than on ethanol to comply with the European/Italian obligations to introduce bio-fuels on the market.

Biogasoline represents to date a minimum percentage out of the total gasoline including biogasoline consumption. According to the Renewable energy Directive (2009/28/EC) the amount of biogasoline reported in the Energy balance is equal to the renewable part of the fuel, calculated as the 37% of the total volume placed on the market.

Biodiesel has been tested since 1994 to 1996 before entering in production since 1998.

Moreover biogas road consumption has been taken into account in the Inventory, representing about 11% of total road natural gas consumption in 2020. It is reported only for the last reporting year 2020 in the IEA - Eurostat – UNECE Energy Questionnaire (as element not covered by Regulation EC No 1099/2008 on energy statistics, therefore it is not mandatory for Countries to transmit it to Eurostat).

Emissions of heavy metals are estimated on the basis of data regarding the fuel and lubricant content and the engine wear; as reported in the EMEP/EEA Emission Inventory Guidebook 2019, these apparent fuel metal content factors originate from the work of Winther and Slentø, 2010, and have been reviewed by the TFEIP expert panel in transport and because of the scarce available information, the uncertainty in the estimate of these values is still considered quite high. In COPERT model heavy metals emission factors have been then updated focusing on the distinction between exhaust and non exhaust share.

Non exhaust emissions of PAHs have also been estimated on the basis of brake and tyres debris-bound values resulting from the EMEP/EEA guidebook 2019.

3.5.2.2 Evaporative emissions

As regards NMVOC, the share of evaporative emissions is provided. These emissions are calculated only for gasoline vehicles: passenger cars, light duty vehicles, mopeds and motorcycles. Depending on temperature and vapour pressure of fuel, evaporative emissions have shown a growth over the years, nevertheless recently the contribution has been reduced by the introduction of control systems such as the canister. The estimation procedure is differentiated according to the processes of diurnal emission, running losses and hot soak emissions (EMEP/EEA, 2019).

3.5.2.3 Emissions from automobile tyre and brake wear

Not exhaust PM emissions from road vehicle tyre and brake wear are estimated. The focus is on the primary particles, deriving directly from tyre and brake wear. The material produced by the effects of wear and attrition between surfaces is subject to evaporation at high temperatures developed by the contact.

Emissions are influenced by, as regards tyres, composition and pressure of tyres, structure and characteristics of vehicles, the peculiarities of the road and, as regards brakes, by the composition of the materials of the components, the position, the configuration systems, and the mechanisms of actuation (EMEP/EEA, 2019).

3.5.2.4 Emissions from automobile road abrasion

Particulate non-exhaust emissions deriving from road surface wear have been introduced in COPERT model, according to the Guidebook methodology (EMEP/EEA, 2019).

Emissions depend on the type of asphalt-based and concrete-based road surfaces, taking into account that composition can vary widely, both from country to country and within countries. The type of tyres used also affect emissions, for instance the wear of the road surface, and the resulting PM concentrations due to resuspension, are considerably high when studded tyres are extensively used during the winter.

The wear of the road surface increases with moisture level, also increasing after salting of the road, since the surface remains wet for longer periods. Other influencing factors are vehicle speed, tyre pressure and air temperature. As a consequence of the decrease of temperature, tyres become less elastic, causing the increase of the road surface wear rates (EMEP/EEA, 2019).

3.5.3 Activity data

The road traffic data used are vehicle-kilometre estimates for the different vehicle types and different road classifications in the national road network. These data have to be further broken down by composition of each vehicle fleet in terms of the fraction of different fuels types powered vehicles on the road and in terms of the fraction of vehicles on the road set by the different emission regulations which applied when the vehicle was first registered. These are related to the age profile of the vehicle fleet.

Basic data derive from different sources. Detailed data on the national fleet composition are found in the yearly report from ACI (ACI, several years), used from 1990 to 2006, except for mopeds for which estimates have been elaborated, for the whole time series, on the basis of National Association of Cycle-Motorcycle Accessories data on mopeds fleet composition and mileages (ANCMA, several years).

The Ministry of Transport (MIMS) provides specific fleet composition data for all vehicle categories from 2007 onwards, starting from 2013 submission. The Ministry of Transport in the national transport yearbook (MIMS, several years) reports mileages time series. Furthermore since 2015 MIMS supplies information relating the distribution of old gasoline cars over the detailed vehicles categories (PRE ECE; ECE 15/00-01; ECE 15/02; ECE 15/03; ECE 15/04; information obtained from the registration year; data used for the updating of the time series since 2007). MIMS data are used relating to: the passenger cars (petrol hybrid and diesel hybrid passenger cars are introduced from 2007 onwards, the detailed “Gasoline < 0.8 l” passenger cars subsector is introduced since 2012 and “Diesel<1.4 l” subsector since 2007 onwards, in addition to the gasoline, diesel, LPG, CNG traditional ones); the diesel and gasoline light commercial vehicles; the breakdown of the heavy duty trucks, buses and coaches fleet according to the different weight classes and fuels (diesel almost exclusively for HDT, a negligible share consists of gasoline vehicles; diesel for coaches; diesel, diesel hybrid and CNG for urban buses); the motorcycles fleet in the detail of subsector and legislation standard of both 2-stroke and 4-stroke categories. Fleet values for mopeds are updated according to the revisions of data published by ANCMA; fleet values for diesel buses are updated according to the updating of the data on urban public buses, published on CNIT.

The National Institute of Statistics carries out annually a survey on heavy goods vehicles, including annual mileages (ISTAT, several years).

The National Association of concessionaries of motorways and tunnels produces monthly statistics on highway mileages by light and heavy vehicles (AISCAT, several years).

The National General Confederation of Transport and Logistics (CONFETRA, several years) and the national Central Committee of road transporters (Giordano, 2007) supplied useful information and statistics about heavy goods vehicles fleet composition and mileages.

Fuel consumption data derive basically from the National Energy Balance (MiSE, MiTE, several years (a)); supplementary information is taken from the Oil Bulletin (MiSE, MiTE, several years (b)). As regards biofuels, the consumption has increased in view of the targets to be respected by Italy and set in the framework of the European directive 20-20-20. The trend of biodiesel is explained by the fact that this biofuel has been tested since 1994 to 1996 before entering in production since 1998. The consumption of bioethanol is introduced since 2008, according to data resulting on the BEN. Biogas is introduced in 2020 being reported only for the last reporting year 2020 in the IEA - Eurostat – UNECE Energy Questionnaire.

Emissions are calculated from vehicles of the following types:

- Gasoline passenger cars;
- Diesel passenger cars;
- LPG passenger cars;
- CNG passenger cars;

- Petrol Hybrid passenger cars;
- Diesel Hybrid passenger cars;
- Gasoline Light Commercial Vehicles (Gross Vehicle Weight (GVW) <= 3.5 tonnes);
- Diesel Light Commercial Vehicles (Gross Vehicle Weight (GVW) <= 3.5 tonnes);
- Rigid-axle Heavy Duty Trucks (GVW > 3.5 tonnes);
- Articulated Heavy Duty Trucks (GVW > 3.5 tonnes);
- Diesel Buses and coaches;
- Diesel Hybrid Buses;
- CNG Buses;
- Mopeds and motorcycles.

In Table 3.16 the historical series of annual consumption data (Mg) for the different fuel types is reported.

Table 3.16 Annual fuel consumption data (Mg)

| Fuel | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|-------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|
| Gasoline Leaded | 12,280,212 | 10,112,250 | 4,542,113 | - | - | - | - | - | - | - | - |
| Gasoline Unleaded | 639,115 | 7,060,391 | 12,175,814 | 13,482,132 | 9,806,890 | 7,809,940 | 7,297,739 | 7,089,221 | 7,286,305 | 7,353,002 | 5,797,395 |
| Diesel | 15,278,022 | 14,445,441 | 17,059,010 | 22,327,864 | 21,557,266 | 21,128,587 | 21,228,198 | 20,101,587 | 20,901,623 | 20,985,104 | 17,103,910 |
| LPG | 1,342,000 | 1,478,000 | 1,422,000 | 1,029,000 | 1,214,000 | 1,654,000 | 1,598,000 | 1,667,372 | 1,614,000 | 1,653,017 | 1,309,461 |
| CNG | 183,770 | 216,804 | 292,214 | 342,756 | 610,502 | 787,148 | 784,406 | 740,695 | 746,783 | 832,667 | 671,528 |
| of which biogas | | | | | | | | | | | 71,349 |
| Biodiesel | - | 44,491 | 64,723 | 200,000 | 1,468,086 | 1,292,079 | 1,141,334 | 1,164,023 | 1,377,205 | 1,409,548 | 1,408,889 |
| Biogasoline | - | - | - | - | 142,106 | 30,420 | 37,808 | 38,455 | 38,238 | 35,401 | 22,841 |

Source: ISPRA elaborations on BEN, BP, UP data

The final reports on the physic-chemical characterization of fossil fuels used in Italy, carried out by the Fuel Experimental Station, that is an Italian Institute operating in the framework of the Department of Industry, are used with the aim to improve fuel quality specifications (surveys conducted in 2000, in 2012 – 2013 and in 2020). Fuel information has also been updated for the entire time series on the basis of the annual reports published by ISPRA about the fuel quality in Italy.

Monitoring of the carbon content of the fuels used in Italy is an ongoing activity at ISPRA (Italian Institute for Environmental Protection and Research). The purpose is to analyse regularly the chemical composition of the used fuels or relevant commercial statistics to estimate the carbon content/emission factor (EF) of the fuels. With reference to the whole inventory, for each primary fuel, a specific procedure has been established.

As regards road transport, Italy fuel specifications values for gasoline, diesel fuel and LPG, derive from Fuel Experimental Station analysis about the properties of transportation fuels sold in Italy and whose results are representative and applicable with reference to four different time phases: 1990 – 1999; 2000 – 2012; 2013 – 2019; since 2020 (Innovhub – Fuel Experimental Station surveys, several years).

As regards natural gas, the national market is characterized by the commercialisation of gases with different chemical composition in variable quantities from one year to the other. The methodology used to estimate the average EF for natural gas per year is based on the available consumption data, referring to the lower heat value (each year the quantities of natural gas imported or produced in Italy are published on the web by the MSE <http://dgerm.sviluppoeconomico.gov.it/dgerm/bilanciogas.asp>).

A normalisation procedure is applied to ensure that the breakdown of fuel consumption by each vehicle type calculated on the basis of the fuel consumption factors then added up matches the BEN figures for total fuel consumption in Italy (adjusted for off-road consumption).

The automatic energy balance process, introduced by COPERT 5, has been applied. The simulation is started up having the target to equalize calculated and statistical consumptions, separately for fuel, at national level, with the aim to obtain final estimates the most accurate as possible. Once all data and input parameters have been inserted and all options have been set reflecting the peculiar situation of the Country, emissions and consumptions are calculated by the model in the detail of the vehicle category legislation standard; then the aggregated consumption values so calculated are compared with the input statistical national aggregated values

(deriving basically from the National Energy Balance, as described above), with the aim to minimize the deviation.

In the following Tables 3.17, 3.18, 3.19 and 3.20 detailed data on the relevant vehicle mileages in the circulating fleet are reported, subdivided according to the main emission regulations (ISPRA elaborations on ACI, ANCMA and MIMS data).

Table 3.17 *Passenger Cars technological evolution: circulating fleet calculated as stock data multiplied by actual mileage (%)*

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|-------|------|------|-------|--------|-------|-------|-------|-------|-------|-------|
| PRE ECE, pre-1973 | 0.04 | 0.03 | 0.01 | 0.01 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 |
| ECE 15/00-01, 1973-1978 | 0.10 | 0.04 | 0.01 | 0.005 | 0.003 | 0.003 | 0.003 | 0.002 | 0.002 | 0.003 | 0.002 |
| ECE 15/02-03, 1978-1984 | 0.30 | 0.15 | 0.03 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| ECE 15/04, 1985-1992 | 0.55 | 0.55 | 0.28 | 0.10 | 0.04 | 0.03 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 |
| PC Euro 1 - 91/441/EEC, from 1/1/93 | 0.001 | 0.24 | 0.27 | 0.17 | 0.05 | 0.02 | 0.02 | 0.01 | 0.01 | 0.01 | 0.01 |
| PC Euro 2 - 94/12/EEC, from 1/1/97 | - | - | 0.39 | 0.32 | 0.21 | 0.12 | 0.10 | 0.09 | 0.08 | 0.07 | 0.06 |
| PC Euro 3 - 98/69/EC Stage2000, from 1/1/2001 | - | - | - | 0.31 | 0.20 | 0.13 | 0.12 | 0.11 | 0.10 | 0.09 | 0.08 |
| PC Euro 4 - 98/69/EC Stage2005, from 1/1/2006 | - | - | - | 0.09 | 0.44 | 0.37 | 0.35 | 0.33 | 0.31 | 0.28 | 0.27 |
| PC Euro 5 - EC 715/2007, from 1/1/2011 | - | - | - | - | 0.04 | 0.25 | 0.24 | 0.23 | 0.21 | 0.18 | 0.17 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 a/b/c | - | - | - | - | - | 0.06 | 0.13 | 0.19 | 0.26 | 0.25 | 0.24 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 d-temp (2019 - 2020) | - | - | - | - | - | - | - | - | - | 0.09 | 0.14 |
| Total | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| a. Gasoline cars technological evolution | | | | | | | | | | | |
| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
| Conventional, pre-1993 | 1.00 | 0.92 | 0.36 | 0.06 | 0.01 | 0.004 | 0.003 | 0.003 | 0.003 | 0.003 | 0.003 |
| PC Euro 1 - 91/441/EEC, from 1/1/93 | - | 0.08 | 0.10 | 0.03 | 0.01 | 0.002 | 0.002 | 0.001 | 0.001 | 0.001 | 0.001 |
| PC Euro 2 - 94/12/EEC, from 1/1/97 | - | - | 0.54 | 0.22 | 0.05 | 0.02 | 0.02 | 0.01 | 0.01 | 0.01 | 0.01 |
| PC Euro 3 - 98/69/EC Stage2000, from 1/1/2001 | - | - | - | 0.57 | 0.31 | 0.15 | 0.14 | 0.11 | 0.09 | 0.08 | 0.08 |
| PC Euro 4 - 98/69/EC Stage2005, from 1/1/2006 | - | - | - | 0.12 | 0.55 | 0.41 | 0.38 | 0.37 | 0.30 | 0.28 | 0.26 |
| PC Euro 5 - EC 715/2007, from 1/1/2011 | - | - | - | - | 0.07 | 0.36 | 0.32 | 0.30 | 0.30 | 0.28 | 0.27 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 a/b/c | - | - | - | - | 0.0001 | 0.05 | 0.14 | 0.21 | 0.29 | 0.28 | 0.27 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 d-temp (2019 - 2020) | - | - | - | - | - | - | - | - | - | 0.06 | 0.10 |
| Total | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| b. Diesel cars technological evolution | | | | | | | | | | | |
| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
| Conventional, pre-1993 | 1.00 | 0.90 | 0.71 | 0.47 | 0.04 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|------|------|------|------|------|------|------|------|------|-------|-------|
| PC Euro 1 - 91/441/EEC, from 1/1/93 | - | 0.10 | 0.20 | 0.26 | 0.02 | 0.01 | 0.01 | 0.01 | 0.01 | 0.004 | 0.004 |
| PC Euro 2 - 94/12/EEC, from 1/1/97 | - | - | 0.09 | 0.19 | 0.08 | 0.03 | 0.03 | 0.02 | 0.02 | 0.02 | 0.01 |
| PC Euro 3 - 98/69/EC Stage2000, from 1/1/2001 | - | - | - | 0.06 | 0.08 | 0.05 | 0.04 | 0.04 | 0.03 | 0.03 | 0.02 |
| PC Euro 4 - 98/69/EC Stage2005, from 1/1/2006 | - | - | - | 0.01 | 0.75 | 0.46 | 0.42 | 0.38 | 0.35 | 0.31 | 0.29 |
| PC Euro 5 - EC 715/2007, from 1/1/2011 | - | - | - | - | 0.03 | 0.36 | 0.34 | 0.32 | 0.30 | 0.28 | 0.26 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 a/b/c | - | - | - | - | - | 0.08 | 0.15 | 0.09 | 0.11 | 0.13 | 0.13 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 d-temp (2017-2019) | - | - | - | - | - | - | - | 0.14 | 0.18 | 0.23 | 0.22 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 d (2020 and later) | - | - | - | - | - | - | - | - | - | - | 0.05 |
| Total | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| c. Lpg cars technological evolution | | | | | | | | | | | |
| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
| PC Conventional - Euro 4 | 1.00 | 1.00 | 1.00 | 1.00 | 0.91 | 0.58 | 0.54 | 0.51 | 0.47 | 0.44 | 0.42 |
| PC Euro 5 - EC 715/2007, from 1/1/2011 | - | - | - | - | 0.09 | 0.32 | 0.31 | 0.31 | 0.30 | 0.29 | 0.29 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 a/b/c | - | - | - | - | - | 0.10 | 0.15 | 0.10 | 0.12 | 0.14 | 0.14 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 d-temp (2017-2019) | - | - | - | - | - | - | - | 0.08 | 0.10 | 0.13 | 0.12 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 d (2020 and later) | - | - | - | - | - | - | - | - | - | - | 0.04 |
| Total | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| d. CNG cars technological evolution | | | | | | | | | | | |
| | 2007 | 2008 | 2009 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | |
| PC Euro 4 - 98/69/EC Stage2005, from 1/1/2006 | | 1.00 | 1.00 | 0.65 | 0.54 | 0.22 | 0.07 | 0.04 | 0.03 | 0.02 | 0.01 |
| PC Euro 5 - EC 715/2007, from 1/1/2011 | | - | - | 0.35 | 0.46 | 0.61 | 0.42 | 0.27 | 0.18 | 0.11 | 0.07 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 a/b/c | | - | - | - | - | 0.16 | 0.51 | 0.13 | 0.15 | 0.15 | 0.09 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 d-temp (2017-2019) | | - | - | - | - | - | - | 0.56 | 0.64 | 0.72 | 0.44 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 d (2020 and later) | | - | - | - | - | - | - | - | - | - | 0.39 |
| Total | | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| e. Hybrid Gasoline cars technological evolution (from 2007 onwards) | | | | | | | | | | | |
| | 2007 | 2008 | 2009 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | |

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|------|------|------|------|------|------|------|------|------|------|------|
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 a/b/c | | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 0.69 | 0.42 | 0.11 | 0.04 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 d-temp (2017-2019) | | - | - | - | - | - | - | 0.31 | 0.58 | 0.89 | 0.39 |
| PC Euro 6 (Since EC 715/2007, from 9/1/2015) - Euro 6 d (2020 and later) | | - | - | - | - | - | - | - | - | - | 0.57 |
| Total | | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| e. Hybrid Diesel cars technological evolution (from 2007 onwards) | | | | | | | | | | | |

Source: ISPRA elaborations on MIMS and ACI data

Table 3.18 *Light Duty Vehicles technological evolution: circulating fleet calculated as stock data multiplied by actual mileage (%)*

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|------|------|------|------|------|------|------|------|------|------|------|
| Conventional, pre 10/1/94 | 1.00 | 0.93 | 0.63 | 0.35 | 0.08 | 0.06 | 0.06 | 0.06 | 0.05 | 0.03 | 0.03 |
| LCV Euro 1 - 93/59/EEC, from 10/1/94 | - | 0.07 | 0.22 | 0.17 | 0.10 | 0.04 | 0.04 | 0.04 | 0.03 | 0.02 | 0.02 |
| LCV Euro 2 - 96/69/EEC, from 10/1/98 | - | - | 0.15 | 0.15 | 0.30 | 0.15 | 0.15 | 0.14 | 0.12 | 0.06 | 0.06 |
| LCV Euro 3 - 98/69/EC Stage2000, from 1/1/2002 | - | - | - | 0.31 | 0.26 | 0.19 | 0.19 | 0.18 | 0.15 | 0.11 | 0.10 |
| LCV Euro 4 - 98/69/EC Stage2005, from 1/1/2007 | - | - | - | 0.01 | 0.25 | 0.32 | 0.32 | 0.32 | 0.28 | 0.24 | 0.22 |
| LCV Euro 5 - 2008 Standards 715/2007/EC, from 1/1/2012 | - | - | - | - | 0.00 | 0.22 | 0.20 | 0.17 | 0.19 | 0.17 | 0.16 |
| LCV Euro 6 (Since 2007/715/EC, from 9/1/2016) - Euro 6 a/b/c | - | - | - | - | - | 0.02 | 0.05 | 0.09 | 0.06 | 0.11 | 0.10 |
| LCV Euro 6 (Since 2007/715/EC, from 9/1/2016) - Euro 6 d-temp (2018 - 2020) | - | - | - | - | - | - | - | - | 0.11 | 0.25 | 0.32 |
| Total | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| a. Gasoline Light Commercial Vehicles technological evolution | | | | | | | | | | | |
| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
| Conventional, pre 10/1/94 | 1.00 | 0.92 | 0.55 | 0.23 | 0.07 | 0.02 | 0.02 | 0.02 | 0.01 | 0.01 | 0.01 |
| LCV Euro 1 - 93/59/EEC, from 10/1/94 | - | 0.08 | 0.21 | 0.11 | 0.05 | 0.02 | 0.02 | 0.01 | 0.01 | 0.01 | 0.01 |
| LCV Euro 2 - 96/69/EEC, from 10/1/98 | - | - | 0.23 | 0.20 | 0.18 | 0.07 | 0.06 | 0.04 | 0.03 | 0.02 | 0.02 |
| LCV Euro 3 - 98/69/EC Stage2000, from 1/1/2002 | - | - | - | 0.45 | 0.34 | 0.21 | 0.18 | 0.12 | 0.09 | 0.07 | 0.06 |
| LCV Euro 4 - 98/69/EC Stage2005, from 1/1/2007 | - | - | - | 0.01 | 0.34 | 0.33 | 0.29 | 0.27 | 0.23 | 0.18 | 0.17 |
| LCV Euro 5 - 2008 Standards 715/2007/EC, from 1/1/2012 | - | - | - | - | 0.01 | 0.35 | 0.37 | 0.35 | 0.34 | 0.33 | 0.30 |
| LCV Euro 6 (Since 2007/715/EC, from 9/1/2016) - Euro 6 a/b/c | - | - | - | - | 0.00 | 0.01 | 0.07 | 0.19 | 0.12 | 0.17 | 0.15 |
| LCV Euro 6 (Since 2007/715/EC, from 9/1/2016) - Euro 6 d-temp (2018 - 2020) | - | - | - | - | - | - | - | - | 0.16 | 0.22 | 0.29 |

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|------|------|------|------|------|------|------|------|------|------|------|
| Total | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| b. Diesel Light Commercial Vehicles technological evolution | | | | | | | | | | | |

Source: ISPRA elaborations on MIMS and ACI data

Table 3.19 Heavy Duty Trucks and Buses technological evolution: circulating fleet calculated as stock data multiplied by actual mileage (%)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|
| Conventional, pre 10/1/93 | 1.00 | 0.90 | 0.67 | 0.39 | 0.20 | 0.02 | 0.02 | 0.02 | 0.02 | 0.01 | 0.01 |
| HDT Euro I - 91/542/EEC Stage I, from 10/1/93 | - | 0.10 | 0.10 | 0.06 | 0.04 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| HDT Euro II - 91/542/EEC Stage II, from 10/1/96 | - | - | 0.22 | 0.27 | 0.15 | 0.08 | 0.07 | 0.06 | 0.05 | 0.05 | 0.04 |
| HDT Euro III - 2000 Standards, 99/96/EC, from 10/1/2001 | - | - | - | 0.27 | 0.36 | 0.34 | 0.31 | 0.28 | 0.25 | 0.22 | 0.20 |
| HDT Euro IV - 2005 Standards, 99/96/EC, from 10/1/2006 | - | - | - | - | 0.07 | 0.10 | 0.09 | 0.08 | 0.08 | 0.07 | 0.07 |
| HDT Euro V - 2008 Standards, 99/96/EC, from 10/1/2009 | - | - | - | - | 0.18 | 0.38 | 0.37 | 0.36 | 0.34 | 0.33 | 0.32 |
| HDT Euro VI (Since 2009/595/EC, from 12/31/2013) - Euro VI A/B/C | - | - | - | - | - | 0.07 | 0.13 | 0.19 | 0.25 | 0.25 | 0.23 |
| HDT Euro VI (Since 2009/595/EC, from 12/31/2013) - Euro VI D/E | - | - | - | - | - | - | - | - | - | 0.06 | 0.12 |
| Total | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| a. Heavy Duty Trucks technological evolution | | | | | | | | | | | |
| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
| Conventional, pre 10/1/93 | 1.00 | 0.93 | 0.65 | 0.34 | 0.13 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.004 |
| Buses Euro I - 91/542/EEC Stage I, from 10/1/93 | - | 0.07 | 0.07 | 0.08 | 0.04 | 0.01 | 0.01 | 0.01 | 0.00 | 0.01 | 0.005 |
| Buses Euro II - 91/542/EEC Stage II, from 10/1/96 | - | - | 0.28 | 0.32 | 0.27 | 0.14 | 0.13 | 0.11 | 0.08 | 0.08 | 0.07 |
| Buses Euro III - 2000 Standards, 99/96/EC, from 10/1/2001 | - | - | - | 0.26 | 0.34 | 0.38 | 0.35 | 0.32 | 0.28 | 0.25 | 0.24 |
| Buses Euro IV - 2005 Standards, 99/96/EC, from 10/1/2006 | - | - | - | - | 0.12 | 0.13 | 0.12 | 0.11 | 0.11 | 0.11 | 0.10 |
| Buses Euro V - 2008 Standards, 99/96/EC, from 10/1/2009 | - | - | - | - | 0.11 | 0.28 | 0.28 | 0.29 | 0.27 | 0.27 | 0.26 |
| Buses Euro VI (Since 2009/595/EC, from 12/31/2013) - Euro VI A/B/C | - | - | - | - | - | 0.05 | 0.10 | 0.16 | 0.24 | 0.22 | 0.21 |
| Buses Euro VI (Since 2009/595/EC, from 12/31/2013) - Euro VI D/E | - | - | - | - | - | - | - | - | - | 0.06 | 0.11 |
| Total | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| b. Diesel Buses technological evolution | | | | | | | | | | | |
| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
| Urban CNG Buses Conventional, pre 10/1/93; Urban CNG Buses Euro I - 91/542/EEC Stage I, from 10/1/93 | 1.00 | 1.00 | 0.11 | 0.01 | 0.003 | 0.003 | 0.003 | 0.002 | 0.002 | 0.001 | 0.001 |
| Urban CNG Buses Euro II - 91/542/EEC Stage II, from 10/1/96 | - | - | 0.89 | 0.20 | 0.10 | 0.05 | 0.04 | 0.04 | 0.03 | 0.03 | 0.02 |
| Urban CNG Buses Euro III - 2000 Standards, 99/96/EC, from 10/1/2001; Urban CNG Buses Euro IV - 2005 Standards, 99/96/EC, from 10/1/2006 | - | - | - | 0.79 | 0.09 | 0.07 | 0.07 | 0.07 | 0.06 | 0.06 | 0.05 |

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|------|------|------|------|------|------|------|------|------|------|------|
| Euro V - 2008 Standards, 99/96/EC, from 10/1/2009; EEV (Enhanced environmentally friendly vehicle; ref. 2001/27/EC and 1999/96/EC line C, optional limit emission values); Urban CNG Buses Euro VI – EC 595/2009, from 12/31/2013 | - | - | - | - | 0.81 | 0.88 | 0.88 | 0.89 | 0.90 | 0.91 | 0.93 |
| Total | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| c. CNG Buses technological evolution | | | | | | | | | | | |
| | 2007 | 2008 | 2009 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 | |
| Buses Euro VI (Since 2009/595/EC, from 12/31/2013) - Euro VI A/B/C | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 0.21 | 0.16 | |
| Buses Euro VI (Since 2009/595/EC, from 12/31/2013) - Euro VI D/E | - | - | - | - | - | - | - | - | 0.79 | 0.84 | |
| Total | | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |
| d. Diesel Hybrid Buses technological evolution (from 2007 onwards) | | | | | | | | | | | |

Source: ISPRA elaborations on MIMS and ACI data

Table 3.20 *Mopeds and motorcycles technological evolution: circulating fleet calculated as stock data multiplied by actual mileage (%)*

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---------------------------------------|------|------|------|------|------|------|---------------|---------------|---------------|--------|-------|
| Mopeds and motorcycles - Conventional | 1.00 | 1.00 | 0.86 | 0.43 | 0.18 | 0.11 | 0.10 | 0.11 | 0.11 | 0.11 | 0.10 |
| Mopeds and motorcycles - Euro 1 | - | - | 0.14 | 0.30 | 0.20 | 0.14 | 0.13 | 0.13 | 0.11 | 0.11 | 0.10 |
| Mopeds and motorcycles - Euro 2 | - | - | - | 0.22 | 0.35 | 0.34 | 0.33 | 0.30 | 0.30 | 0.23 | 0.22 |
| Mopeds and motorcycles - Euro 3 | - | - | - | 0.04 | 0.27 | 0.42 | 0.43 | 0.41 | 0.38 | 0.41 | 0.39 |
| Mopeds and motorcycles - Euro 4 | - | - | - | - | - | - | 0.01 | 0.05 | 0.10 | 0.14 | 0.18 |
| Mopeds and motorcycles - Euro 5 | - | - | - | - | - | - | 0.0000 002 | 0.0000 002 | 0.0000 005 | 0.0002 | 0.007 |
| Total | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 | 1.00 |

Source: ISPRA elaborations on ANCMA, ACI and MIMS data

Average emission factors are calculated for average speeds by three driving modes (urban, rural and motorway) combined with the vehicle kilometres travelled and vehicle categories.

ISPRA estimates total annual vehicle kilometres for the road network in Italy by vehicle type, see Table 3.21, based on data from various sources:

- Ministry of Transport (MIMS, several years) for rural roads and on other motorways; the latter estimates are based on traffic counts from the rotating census and core census surveys of ANAS (management authority for national road and motorway network);
- highway industrial association for fee-motorway (AISCAT, several years);
- local authorities for built-up areas (urban).

Table 3.21 *Evolution of fleet consistency and mileage*

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|------|------|------|------|------|------|------|------|------|------|------|
| All passenger vehicles (including moto), total mileage (10 ⁹ veh-km/y) | 350 | 412 | 455 | 453 | 450 | 444 | 445 | 447 | 442 | 445 | 354 |
| Car fleet (10 ⁶) | 27 | 30 | 32 | 34 | 36 | 37 | 38 | 38 | 39 | 40 | 40 |
| Moto, total mileage (10 ⁹ veh-km/y) | 30 | 41 | 42 | 42 | 36 | 33 | 31 | 28 | 27 | 28 | 24 |

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|------|------|------|------|------|------|------|------|------|------|------|
| Moto fleet (10 ⁶) | 7 | 7 | 9 | 9 | 9 | 10 | 10 | 10 | 10 | 10 | 10 |
| Goods transport, total mileage (10 ⁹ veh-km/y) | 69 | 76 | 83 | 104 | 83 | 62 | 60 | 51 | 65 | 66 | 59 |
| Truck fleet (10 ⁶), including LDV | 2 | 3 | 3 | 4 | 5 | 5 | 5 | 5 | 5 | 5 | 5 |

Source: ISPRA elaborations

Notes: The passenger vehicles include passenger cars, buses and moto; the moto fleet includes mopeds and motorcycles; in the goods transport light commercial vehicles and heavy duty trucks are included.

3.5.4 Time series and key categories

The analysis of time series on transport data shows a trend that is the result of the general growth in mobility demand and consumptions until 2007, followed by a decrease basically due to the economic crisis on one side, and of the introduction of advanced technologies limiting emissions in modern vehicles in recent years, on the other side; then the growth in the years 2018-2019 is followed by a sharp decrease in 2020 due to the pandemic crisis.

More in details, passenger cars and light duty vehicles emissions trends are driven by a gradual decrease over the years of gasoline fuel consumption balanced by an increase of diesel fuel which is the main driver for NO_x and PM emissions. At pollutant level emission trends are driven not only by fuel but also by changes in technologies which are reflected in the COPERT model by the annual vehicle fleet. Due to the penetration of new vehicles with more stringent pollutant limits, some pollutant emissions decreased faster than other. An important role has been played also by the distribution between diesel and gasoline fuel consumptions. In the last years an increase of diesel fuelled vehicles and a decrease of gasoline ones have been registered and diesel fuel new technologies resulted in a slower decrease of NO_x emission than expected.

Regarding heavy duty vehicles emissions trends are explained by the variations estimated in mileages time series data correlated to the variations registered in fuel consumptions; annual variation are explained by the general trend of national economic growth and in particular commercial and industrial activities.

Emissions trends regarding mopeds and motorcycles are explained by the variations estimated in mileages time series data correlated to the variations registered in gasoline consumptions. The annual penetration of new technologies explains annual emission trends.

The general decrease between 2019 and 2020 is explained by the pandemic crisis.

In Table 3.22 the list of key categories by pollutant identified for road transport in 2020, 1990 and at trend assessment is reported.

Table 3.22 List of key categories for pollutant in the road transport in 2020, 1990 and in the trend

| | Key categories in 2020 | | | Key categories in 1990 | | | Key categories in trend | | | |
|-------------------|------------------------|---------|---------|------------------------|--------|---------|-------------------------|--------|---------|--------|
| | 1A3bi | 1A3bii | 1A3biii | 1A3bi | 1A3bii | 1A3biii | 1A3bi | 1A3bii | 1A3biii | |
| NO _x | 1A3bi | 1A3bii | 1A3biii | 1A3bi | 1A3bii | 1A3biii | 1A3bi | 1A3bii | 1A3biii | |
| NMVOG | 1A3bi | 1A3biv | 1A3bv | 1A3bi | 1A3biv | 1A3bv | 1A3bi | 1A3biv | | |
| NH ₃ | | | | | | | 1A3bi | | | |
| CO | 1A3bi | 1A3biv | | 1A3bi | 1A3biv | | 1A3bi | | | |
| PM ₁₀ | 1A3bi | 1A3bvi | 1A3bvii | 1A3bi | 1A3bii | 1A3biii | 1A3bvi | 1A3bii | 1A3biii | 1A3bvi |
| PM _{2.5} | 1A3bvi | | | 1A3bi | 1A3bii | 1A3biii | 1A3bvi | 1A3bii | 1A3biii | |
| BC | 1A3bi | 1A3biii | | 1A3bi | 1A3bii | 1A3biii | | 1A3bii | 1A3biii | |
| Pb | 1A3bvi | | | 1A3bi | 1A3biv | | | 1A3bi | | |
| Cd | 1A3bi | | | | | | | 1A3bi | | |

Source: ISPRA elaborations

In 2020 key categories are identified for the following pollutants: nitrogen oxides, non methane volatile organic compounds, carbon monoxide, particulate matter with diameter less than 10 µm, particulate matter with diameter less than 2.5 µm, black carbon, lead and cadmium.

Nitrogen oxides emissions show a decrease since 1990 of 78.6%. Emissions are mainly due to diesel vehicles. The decrease observed since 1990 in emissions relates to all categories except for diesel passenger cars, hybrid categories and CNG buses.

In 2020, emissions of nitrogen oxides (Table 3.23) from passenger cars, light-duty vehicles and heavy-duty trucks including buses are key categories. The same categories are identified as key categories in 1990 and in trend.

Table 3.23 *Time series of nitrogen oxides emissions in road transport (Gg)*

| Source categories for NFR Subsector 1.A.3.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|--------------|----------------|--------------|--------------|---------------|--------------|--------------|--------------|---------------|--------------|--------------|
| <i>Gg</i> | | | | | | | | | | | |
| 1.A.3.b.i Passenger cars | 590.9 | 628.5 | 388.6 | 240.2 | 176.9 | 167.4 | 162.7 | 158.7 | 152.6 | 144.4 | 108.8 |
| 1.A.3.b.ii Light-duty vehicles | 60.6 | 69.5 | 71.2 | 75.15 | 53.3 | 41.1 | 40.7 | 34.8 | 41.5 | 41.5 | 35.2 |
| 1.A.3.b.iii Heavy-duty vehicles including buses | 340.3 | 336.1 | 311.4 | 306.7 | 186.9 | 1149 | 106.2 | 86.4 | 88.8 | 81.9 | 66.8 |
| 1.A.3.b.iv Mopeds and motorcycles | 4.29 | 5.55 | 6.07 | 6.85 | 5.08 | 4.30 | 3.93 | 3.49 | 3.20 | 3.08 | 2.53 |
| Total emissions | 996.1 | 1,039.7 | 777.3 | 628.9 | 422.15 | 327.6 | 313.5 | 283.3 | 286.15 | 270.9 | 213.3 |

Source: ISPRA elaborations

As regards non methane volatile organic compounds, emissions from passenger cars, mopeds and motorcycles and gasoline evaporation are key categories in 2020 and 1990; emissions from passenger cars and mopeds and motorcycles are key categories in trend.

Despite the decline of about -88.0% since 1990 of emissions of non methane volatile organic compounds from this category, road transport (Table 3.24) is the fourth source at national level after the use of solvents, the not industrial combustion and agriculture; this trend is due to the combined effects of technological improvements that limit VOCs from tail pipe and evaporative emissions (for cars) and the expansion of two-wheelers fleet. In Italy there is in fact a remarkable fleet of motorbikes and mopeds (about 10 million vehicles in 2020) that uses gasoline and it is increased of about 52% since 1990 (this fleet not completely complies with strict VOC emissions controls).

Table 3.24 *Time series of non methane volatile organic compounds emissions in road transport (Gg)*

| Source categories for NFR Subsector 1.A.3.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--------------|
| <i>Gg</i> | | | | | | | | | | | |
| 1.A.3.b.i Passenger cars | 430.45 | 467.18 | 261.44 | 122.45 | 48.33 | 28.29 | 24.93 | 23.88 | 22.07 | 23.33 | 16.58 |
| 1.A.3.b.ii Light-duty vehicles | 15.56 | 18.47 | 13.91 | 11.52 | 5.77 | 2.09 | 1.78 | 1.25 | 1.30 | 0.92 | 0.72 |
| 1.A.3.b.iii Heavy-duty vehicles including buses | 26.35 | 25.20 | 20.52 | 17.34 | 8.32 | 3.51 | 3.23 | 2.61 | 2.67 | 2.46 | 2.01 |
| 1.A.3.b.iv Mopeds and motorcycles | 175.51 | 248.77 | 225.42 | 152.44 | 76.97 | 47.05 | 42.81 | 38.97 | 35.67 | 32.27 | 26.69 |
| 1.A.3.b.v Gasoline evaporation | 119.22 | 119.10 | 87.87 | 55.55 | 45.28 | 46.05 | 44.90 | 46.81 | 46.52 | 44.14 | 45.73 |
| Total emissions | 767.08 | 878.71 | 609.16 | 359.30 | 184.67 | 126.99 | 117.65 | 113.52 | 108.23 | 103.12 | 91.74 |

Source: ISPRA elaborations

Carbon monoxide emissions from passenger cars and mopeds and motorcycles are key categories in 2020 and 1990; passenger cars are also key category in trend. The time series of CO emissions is reported in Table 3.25.

Table 3.25 *Time series of carbon monoxide emissions in road transport (Gg)*

| Source categories for NFR Subsector 1.A.3.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|-----------------|----------------|----------------|----------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <i>Gg</i> | | | | | | | | | | | |
| 1.A.3.b.i Passenger cars | 4,124.60 | 4,151.8 | 2,137.7 | 1,078.9 | 478.11 | 290.83 | 260.19 | 251.34 | 238.80 | 249.91 | 178.15 |
| 1.A.3.b.ii Light-duty vehicles | 175.06 | 207.29 | 140.17 | 97.63 | 45.28 | 17.98 | 15.44 | 12.52 | 12.67 | 8.23 | 6.54 |
| 1.A.3.b.iii Heavy-duty vehicles including buses | 81.44 | 79.76 | 70.24 | 70.21 | 46.69 | 33.33 | 31.09 | 25.90 | 26.96 | 25.30 | 20.89 |
| 1.A.3.b.iv Mopeds and motorcycles | 493.42 | 667.33 | 625.65 | 434.49 | 206.23 | 137.21 | 124.09 | 113.88 | 108.03 | 106.21 | 88.40 |
| Total emissions | 4,874.52 | 5,106.1 | 2,973.8 | 1,681.2 | 776.32 | 479.34 | 430.81 | 403.64 | 386.46 | 389.65 | 293.98 |

Source: ISPRA elaborations

A strong contribution to total emissions is given by gasoline vehicles (about 78.4% in 2020, although since 1990 a decrease of about -94.9% is observed); since 1990 to 2020 a general decrease, of about -94.0%, is observed.

Emissions of PM10 (Table 3.26) deriving from passenger cars, light-duty vehicles, heavy-duty vehicles including buses, road vehicle tyre and brake wear are key categories in 1990; emissions from passenger cars, road vehicle tyre and brake wear and emissions from road surface wear are key categories in 2020; emissions from passenger cars, light-duty vehicles, heavy-duty vehicles including buses and from road vehicle tyre and brake wear are key category in trend.

As regards PM2.5 (Table 3.27), emissions from passenger cars, light-duty vehicles, heavy-duty vehicles including buses are key categories in 1990 and in trend, road vehicle tyre and brake wear is also key category in 1990 and in 2020.

With regard to particulate matter, the relative weight of the non-exhaust component in total emissions becomes progressively more significant over the years.

Table 3.26 Time series of particulate matter with diameter less than 10 µm emissions in road transport (Gg)

| Source categories for NFR Subsector 1.A.3.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Gg | | | | | | | | | | | |
| 1.A.3.b.i Passenger cars | 18.88 | 13.64 | 11.84 | 9.55 | 8.16 | 5.62 | 5.16 | 4.79 | 3.80 | 3.71 | 2.65 |
| 1.A.3.b.ii Light-duty vehicles | 10.36 | 11.51 | 10.27 | 8.84 | 4.66 | 1.68 | 1.42 | 0.98 | 1.05 | 0.83 | 0.65 |
| 1.A.3.b.iii Heavy-duty vehicles including buses | 13.56 | 13.30 | 10.74 | 9.09 | 4.41 | 1.85 | 1.68 | 1.34 | 1.33 | 1.18 | 0.94 |
| 1.A.3.b.iv Mopeds and motorcycles | 3.70 | 5.26 | 4.86 | 3.16 | 1.47 | 0.88 | 0.80 | 0.74 | 0.69 | 0.66 | 0.54 |
| 1 A 3 b vi Road Transport: Automobile tyre and brake wear | 7.68 | 8.71 | 9.40 | 10.13 | 9.47 | 9.27 | 8.63 | 8.10 | 8.39 | 8.44 | 6.93 |
| 1.A.3.b.vii Road transport: Automobile road abrasion | 4.18 | 4.68 | 4.99 | 5.29 | 4.81 | 4.48 | 4.48 | 4.36 | 4.52 | 4.55 | 3.74 |
| Total emissions | 58.35 | 57.09 | 52.10 | 46.06 | 32.98 | 23.79 | 22.17 | 20.31 | 19.78 | 19.37 | 15.46 |

Source: ISPRA elaborations

Table 3.27 Time series of particulate matter with diameter less than 2.5 µm emissions in road transport (Gg)

| Source categories for NFR Subsector 1.A.3.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| Gg | | | | | | | | | | | |
| 1.A.3.b.i Passenger cars | 18.88 | 13.64 | 11.84 | 9.55 | 8.16 | 5.62 | 5.16 | 4.79 | 3.80 | 3.71 | 2.65 |
| 1.A.3.b.ii Light-duty vehicles | 10.36 | 11.51 | 10.27 | 8.84 | 4.66 | 1.68 | 1.42 | 0.98 | 1.05 | 0.83 | 0.65 |
| 1.A.3.b.iii Heavy-duty vehicles including buses | 13.56 | 13.30 | 10.74 | 9.09 | 4.41 | 1.85 | 1.68 | 1.34 | 1.33 | 1.18 | 0.94 |
| 1.A.3.b.iv Mopeds and motorcycles | 3.70 | 5.26 | 4.86 | 3.16 | 1.47 | 0.88 | 0.80 | 0.74 | 0.69 | 0.66 | 0.54 |
| 1 A 3 b vi Road Transport: Automobile tyre and brake wear | 4.17 | 4.74 | 5.13 | 5.52 | 5.17 | 5.03 | 4.73 | 4.46 | 4.63 | 4.66 | 3.83 |
| 1.A.3.b.vii Road transport: Automobile road abrasion | 2.26 | 2.52 | 2.69 | 2.86 | 2.60 | 2.42 | 2.42 | 2.35 | 2.44 | 2.46 | 2.02 |
| Total emissions | 52.92 | 50.97 | 45.54 | 39.01 | 26.46 | 17.49 | 16.21 | 14.67 | 13.94 | 13.50 | 10.64 |

Source: ISPRA elaborations

Emissions of particulate matter with diameter less than 10µm and less than 2.5µm show a decreasing trend since 1990 respectively of about -73.5% and -79.9%; despite the decrease, diesel vehicles (passenger cars, light duty vehicles and heavy duty trucks including buses) are mainly responsible for road transport emissions giving a strong contribution to total emissions, in 2020 about 74.7% and 77.0% out of the total for PM10 and PM2.5 respectively.

Emissions of black carbon are reported in Table 3.28. Emissions from passenger cars and heavy-duty trucks including buses are key categories in 2020; emissions from passenger cars, light-duty vehicles and heavy-duty trucks including buses are key categories in 1990 and in trend.

The emissions trend is generally decreasing (-84.1% since 1990). The main contribution to total emissions is given by diesel vehicles, in 2020 equal to 93.4% out of the total. Despite of the decrease, road transport is the second source of emissions (the main source is non industrial combustion) at national level in 2020 (24.2%).

Table 3.28 *Time series of black carbon emissions in road transport (Gg)*

| Source categories for NFR Subsector 1.A.3.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|--------------|--------------|--------------|--------------|--------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <i>Gg</i> | | | | | | | | | | | |
| 1.A.3.b.i Passenger cars | 10.33 | 7.51 | 7.37 | 7.23 | 6.73 | 4.59 | 4.21 | 3.90 | 3.04 | 2.96 | 2.10 |
| 1.A.3.b.ii Light-duty vehicles | 5.69 | 6.37 | 6.06 | 5.92 | 3.49 | 1.34 | 1.13 | 0.77 | 0.82 | 0.64 | 0.51 |
| 1.A.3.b.iii Heavy-duty vehicles including buses | 6.78 | 6.79 | 5.71 | 5.17 | 2.67 | 1.24 | 1.13 | 0.90 | 0.89 | 0.79 | 0.63 |
| 1.A.3.b.iv Mopeds and motorcycles | 0.66 | 0.95 | 0.87 | 0.55 | 0.25 | 0.15 | 0.14 | 0.12 | 0.12 | 0.11 | 0.09 |
| 1 A 3 b vi Road Transport: Automobile tyre and brake wear | 0.44 | 0.50 | 0.54 | 0.58 | 0.55 | 0.53 | 0.51 | 0.48 | 0.50 | 0.50 | 0.41 |
| 1.A.3.b.vii Road transport: Automobile road abrasion | 0.09 | 0.10 | 0.11 | 0.11 | 0.10 | 0.09 | 0.09 | 0.09 | 0.10 | 0.10 | 0.08 |
| Total emissions | 23.99 | 22.21 | 20.66 | 19.56 | 13.79 | 7.94 | 7.20 | 6.27 | 5.46 | 5.11 | 3.82 |

Source: ISPRA elaborations

Emissions of cadmium are reported in Table 3.29. Cadmium emissions from passenger cars are key categories in 2020 and in trend.

Emissions show a decrease since 1990 of about -18.9%, representing in 2020 the 8.6% of the national total. In 2020 most of the emissions derive from passenger cars (65.0%); non exhaust emissions from automobile tyre and brake wear are equal to 12.7% of the total.

Table 3.29 *Time series of Cadmium emissions in road transport (Gg)*

| Source categories for NFR Subsector 1.A.3.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| <i>Mg</i> | | | | | | | | | | | |
| 1.A.3.b.i Passenger cars | 0.21 | 0.24 | 0.27 | 0.26 | 0.26 | 0.27 | 0.27 | 0.27 | 0.27 | 0.27 | 0.21 |
| 1.A.3.b.ii Light-duty vehicles | 0.02 | 0.03 | 0.03 | 0.04 | 0.04 | 0.03 | 0.02 | 0.02 | 0.03 | 0.03 | 0.02 |
| 1.A.3.b.iii Heavy-duty vehicles including buses | 0.03 | 0.03 | 0.03 | 0.03 | 0.03 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 |
| 1.A.3.b.iv Mopeds and motorcycles | 0.09 | 0.13 | 0.13 | 0.11 | 0.08 | 0.05 | 0.05 | 0.04 | 0.04 | 0.03 | 0.03 |
| 1.A.3.b.vi Road transport: Automobile tyre and brake wear | 0.05 | 0.05 | 0.06 | 0.06 | 0.06 | 0.06 | 0.05 | 0.05 | 0.05 | 0.05 | 0.04 |
| Total emissions | 0.41 | 0.49 | 0.52 | 0.51 | 0.46 | 0.43 | 0.42 | 0.41 | 0.41 | 0.41 | 0.33 |

Source: ISPRA elaborations

Emissions of SO_x, NH₃ and Pb are shown in Table 3.30; SO_x are not key categories; Pb emissions from road vehicle tyre and brake wear are key category in 2020, Pb emissions from passenger cars and from mopeds and motorcycles are key categories in 1990, Pb emissions from passenger cars are key categories in trend; emissions of NH₃ from passenger cars are key categories in trend. In 2020 emissions of these pollutants deriving from road transport are less important compared to other sectors. Emissions of SO_x and Pb show strong decreases. Since 2002, due to limits on fuels properties imposed by legislation, Pb resulting emissions are almost completely non exhaust (road vehicle tyre and brake wear Pb emissions decrease of about 14.7% since 1990); total Pb emissions decrease of -99.8% since 1990, representing in 2020 about 5.5% of the national total. SO_x emissions decrease by -99.8%, representing 0.4% of the total in 2020. Emissions of NH₃, despite the strong increase since 1990, in 2020 account for just 1.2% out of the total.

Table 3.30 *Time series of sulphur oxides, ammonia and lead emissions in road transport*

| SO _x , NH ₃ , Pb Total Emissions for NFR Subsector 1.A.3.b | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|----------|----------|--------|-------|-------|-------|-------|-------|-------|-------|------|
| SO _x (Gg) | 129.29 | 71.60 | 11.93 | 2.21 | 0.43 | 0.38 | 0.41 | 0.41 | 0.41 | 0.39 | 0.32 |
| NH ₃ (Gg) | 0.76 | 5.30 | 20.53 | 15.69 | 9.91 | 6.35 | 5.94 | 5.64 | 5.82 | 5.70 | 4.47 |
| Pb (Mg) | 3,782.34 | 1,617.64 | 690.00 | 13.22 | 12.30 | 12.19 | 10.99 | 10.13 | 10.46 | 10.51 | 8.63 |

Source: ISPRA elaborations

3.5.5 QA/QC and Uncertainty

Data used for estimating emissions from the road transport sector, derive from different sources, including official statistics providers and industrial associations.

A specific procedure undertaken for improving the inventory in the sector regards the establishment of a national expert panel in road transport which involves, on a voluntary basis, different institutions, local agencies and industrial associations cooperating for improving activity data and emission factors accuracy. In this group emission estimates are presented annually and new methodologies are shared and discussed. Reports and data of the meetings can be found at the following address:

http://groupware.sinanet.isprambiente.it/expert_panel/library.

Besides, over time recalculations of time series estimates have been discussed with national experts in the framework of an *ad hoc* working group on air emissions inventories. The group is chaired by ISPRA and includes participants from the local authorities responsible for the preparation of local inventories, sectoral experts, the Ministry of Environment, Land and Sea, and air quality model experts. Recalculations are comparable with those resulting from application of the model at local level. Top-down and bottom-up approaches have been compared with the aim at identifying the major problems and future possible improvements in the methodology to be addressed (emission estimates at the link: <https://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>).

A Montecarlo analysis has been carried out by EMISIA on behalf of the Joint Research Centre (Kouridis et al., 2010) in the framework of the study “Uncertainty estimates and guidance for road transport emission calculations” for 2005 emissions. The study shows an uncertainty assessment, at Italian level, for road transport emissions on the basis of 2005 input parameters of the COPERT 4 model (v. 7.0).

3.5.6 Recalculation

The annual update of the emissions time series from road transport implies a periodic review process.

In 2022 submission the historical series has been revised mainly as a result of the upgrade of COPERT model version used (from version 5.4.36 in last submission to 5.5.1 in submission 2022), which resulted in various methodological updates, the main ones being: introduction of Solid Particle Number (SPN23) emission factors for all vehicle categories; updated PM exhaust emission factor; the calculation of N₂O and NH₃ emissions for PC LPG & CNG and Urban Buses CNG has been corrected; NFR export has been corrected; the calculation of HM emissions has been corrected; fuel consumption from fossil part of biofuels is reported in the CRF export (sheet: “Other fossil fuels”); correction of NO_x emission factors for PC Diesel PHEV (Euro 6d-temp and 6d); hot emission factors for Quad & ATVs and Micro-cars have been corrected. Furthermore various software upgrades have been implemented.

Italian road vehicles electricity consumption data derive from Eurostat database (<https://ec.europa.eu/eurostat/data/database>). COPERT input road electricity consumption data have been calibrated on the basis of the vehicles categories using electricity actually included in COPERT classification. The change of the version was also the chance for a general revision of input data and parameters.

The analysis has been updated on the basis of the results of the survey, carried out by Innovhub, on the fuels used in road transport in Italy in 2020. In particular the results obtained for biodiesel in 2020 have been used to update the entire historical series (previously, in the absence of country specific parameters, Eurostat Energy Balance values were used).

The results of the studies, carried out over the years, about the properties of transportation fuels sold in Italy, for consistency reasons respect to the procedure applied for the other Inventory sectors, are considered representative with reference to four different time phases: 1990 – 1999; 2000 – 2012; 2013 – 2019; since 2020 (Innovhub – Fuel Experimental Station surveys, several years).

Respect to last submission, biogas road consumption has been taken into account in the Inventory, reported only for the last reporting year 2020 in the IEA - Eurostat – UNECE Energy Questionnaire (about 11% of total road natural gas consumption).

Natural gas estimates have been revised since 2017 according to the revision of parameters applied for the estimation of emissions from this fuel for the whole Inventory.

Respect to last submission, the estimation of the allocation of the Euro 6 circulating fleet (among the categories Euro 6 a/b/c, Euro 6 d-temp and Euro 6 d) has been revised according the classification reported on the EMEP/EEA air pollutant emission inventory guidebook 2019 (Update Oct. 2021) due to the fact that between COPERT version 5.2.2 classification (previously kept as term of reference) and the current version 5.5.1 there was a change in the Euro 6 standards allocation to year registrations.

Mopeds circulating fleet estimation has been revised since 2012 in accordance with data provided by the Ministry of Transport for 2020, in order to address inconsistencies related to old and not classified vehicles in Ministry of Transport database.

Estimation of urban buses fleet data has been updated since 2019 in consequence of the updating of Ministry of transport statistics. The estimation of the buses fleet data has been revised since 2007 in relation to the urban/extra-urban split with reference to data and information about vehicles radiations per bus category, supplied from Ministry of Transport.

The estimation of non exhaust PAH has been revised, respect to last submission, differentiating the calculation for the two components tyre and brake wear (on the basis of PM TSP non exhaust emissions, now COPERT model gives in output the detail but previously non exhaust emissions were aggregated), according to 2019 EMEP\EEA Guidebook.

3.5.7 Planned improvements

Improvements for the next submission will be connected to the possible new availability of data and information regarding activity data, calculation factors and parameters, new developments of the methodology and the update of the software.

3.6 RAILWAYS (NFR SUBSECTOR 1.A.3.c)

The electricity used by the railways for electric traction is supplied from the public distribution system, so the emissions arising from its generation are reported under category 1.A.1.a Public Electricity.

Emissions from diesel trains are reported under the IPCC category 1.A.3.c Railways. Estimates are based on the gasoil consumption for railways reported in BEN (MSE, several years [a]), updated since 2018 according to EUROSTAT methodology (<http://dgsaie.mise.gov.it/dgerm/ben.asp>), and on the methodology Tier1, and emission factors from the EMEP/EEA Emission Inventory Guidebook 2019 (EMEP/EEA, 2019).

Fuel consumption data are collected by the Ministry of Economic Development, responsible of the energy balance, from the companies with diesel railways. The activity is present only in those areas without electrified railways, which are limited in the national territory. The trend reflects the decrease of the use of these railways. Because of low values, emissions from railways do not represent a key category. In Table 3.31, diesel consumptions (TJ) and nitrogen oxides, non-methane volatile organic compounds, sulphur oxides, ammonia, particulate and carbon monoxide emissions (Gg) are reported.

Emissions of Pb from 2002 are reported as 'NA', because of the introduction of unleaded liquid fuels in the market in 2002. In particular, heavy metals contents values derive from the analysis about the physical - chemical characterization of fossil fuels used in Italy (Innovhub, Fuel Experimental Station, several years).

Table 3.31 *Consumptions and emissions time series in railways*

| Consumptions and Emissions for NFR Subsector 1.A.3.c | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|----------|----------|----------|----------|----------|--------|--------|----------|----------|----------|----------|
| Diesel Consumption (TJ) | 8,370.25 | 8,199.43 | 5,850.63 | 4,142.42 | 2,690.44 | 939.52 | 640.58 | 1,409.28 | 1,879.04 | 1,836.33 | 1,851.66 |
| Emissions from diesel trains (Gg) | | | | | | | | | | | |
| NO _x | 10.27 | 10.06 | 7.18 | 5.08 | 3.24 | 1.00 | 0.67 | 1.43 | 1.85 | 1.75 | 1.72 |
| NM VOC | 0.91 | 0.89 | 0.64 | 0.45 | 0.29 | 0.09 | 0.06 | 0.13 | 0.18 | 0.17 | 0.17 |
| SO _x | 1.18 | 0.77 | 0.08 | 0.01 | 0.001 | 0.0003 | 0.0002 | 0.001 | 0.001 | 0.001 | 0.001 |
| NH ₃ | 0.001 | 0.001 | 0.001 | 0.001 | 0.0004 | 0.0002 | 0.0001 | 0.0002 | 0.0003 | 0.0003 | 0.0003 |
| PM _{2.5} | 0.27 | 0.26 | 0.19 | 0.13 | 0.08 | 0.03 | 0.02 | 0.04 | 0.05 | 0.05 | 0.04 |
| PM ₁₀ | 0.28 | 0.28 | 0.20 | 0.14 | 0.09 | 0.03 | 0.02 | 0.04 | 0.05 | 0.05 | 0.04 |
| TSP | 0.29 | 0.28 | 0.20 | 0.14 | 0.09 | 0.03 | 0.02 | 0.04 | 0.05 | 0.05 | 0.04 |
| BC | 0.18 | 0.18 | 0.13 | 0.09 | 0.06 | 0.02 | 0.01 | 0.03 | 0.03 | 0.03 | 0.03 |
| CO | 2.10 | 2.05 | 1.47 | 1.04 | 0.67 | 0.24 | 0.16 | 0.35 | 0.47 | 0.46 | 0.46 |

Source: ISPRA elaborations

In the review process has been observed the existence of at least one steam engine still operating in Italy. It is an historic train used only for few days per year and probably fuelled with biomass nowadays instead of coal. Nor biomass or coal are reported in the energy balance for railways activities. Anyway this possible source of emission could be considered insignificant.

In this submission recalculation affected PM2.5 emissions estimate from 1990 to 2009, updated according the methodology reported in the 2019 EMEP/EEA Guidebook, in consequence of the implementation of the recommendation raised during the 2021 NECD Review.
No specific improvements are planned for the next submission.

3.7 NAVIGATION (NFR SUBSECTOR 1.A.3.D)

3.7.1 Overview

This source category includes all emissions from fuels delivered to water-borne navigation. Emissions decreased from 1990 to 2020, because of the reduction in fuel consumed in harbour and navigation activities; the number of movements, showing an increase since 1990, reverses the trend in recent years. National navigation is a key category in 2020 with respect to emissions of SO_x, NO_x, PM₁₀, PM_{2.5}, BC and CO.

3.7.2 Methodological issues

Emissions of the Italian inventory from the navigation sector are carried out according to the IPCC Guidelines and Good Practice Guidance (IPCC, 1997; IPCC, 2000) and the EMEP/EEA Guidebook (EMEP/EEA, 2019). In particular, a national methodology has been developed following the EMEP/EEA Guidebook which provides details to estimate emissions from domestic navigation, specifying recreational craft, ocean-going ships by cruise and harbour activities; emissions from international navigation are also estimated and included as memo item but not included in national totals (EMEP/EEA, 2019). Inland, coastal and deep-sea fishing are estimated and reported under 1.A.4.c. International inland waterways do not occur in Italy.

The methodology developed to estimate emissions is based on the following assumptions and information.

Activity data comprise both fuel consumptions and ship movements, which are available in different level of aggregation and derive from different sources as specified here below:

- Total deliveries of fuel oil, gas oil and marine diesel oil to marine transport are given in national energy balance (MSE, several years (a)) but the split between domestic and international is not provided;
- Naval fuel consumption for inland waterways, ferries connecting mainland to islands and leisure boats, is also reported in the national energy balance as it is the fuel for shipping (MSE, several years (a));
- Data on annual arrivals and departures of domestic and international shipping calling at Italian harbours are reported by the National Institute of Statistics in the statistics yearbooks (ISTAT, several years (a)) and Ministry of Transport in the national transport statistics yearbooks (MIMS, several years).

As for emission and consumption factors, figures are derived by the EMEP/EEA guidebook (EMEP/EEA, 2019), both for recreational and harbour activities and national cruise, taking into account national specificities. These specificities derive from the results of a national study which, taking into account detailed information on the Italian marine fleet and the origin-destination movement matrix for the year 1997, calculated national values (ANPA, 2001; Trozzi et al., 2002 (b)) on the basis of the default emission and consumption factors reported in the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007).

National average emissions and consumption factors were therefore estimated for harbour and cruise activities both for domestic and international shipping from 1990 to 1999. In 2009 submission the study was updated for the years 2004, 2005 and 2006 in order to consider most recent trends in the maritime sector both in terms of modelling between domestic and international consumptions and improvements of operational activities in harbour (TECHNE, 2009). On the basis of the results, national average emissions and consumption factors were updated from 2000.

Specifically, for the years referred to in the surveys, the current method estimates emissions from the number of ships movements broken down by ship type at each of the principal Italian ports considering the information of whether the ship movement is international or domestic, the average tonnage and the relevant distance travelled.

For those years, in fact, figures on the number of arrivals, destination, and fleet composition have been provided by the local port authorities and by the National Institute of Statistics (ISTAT, 2009), covering about 90% of the official national statistics on ship movements for the relevant years. Consumption and emission factors are those derived from the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007) and refer to the Tier 3 ship movement methodology that takes into account origin-destination ship movements matrices as well as technical information on the ships, as engine size, gross tonnage of ships and operational times in harbours. On the basis of sample information, estimates have been carried out at national level for the relevant years considering the official statistics of the maritime sector.

In general, to carry out national estimates of greenhouse gases and other pollutants in the Italian inventory for harbour and domestic cruise activities, consumptions and emissions are calculated for the complete time

series using the average consumption and emission factors multiplied by the total number of movements. On the other hand, for international cruise, consumptions are derived by difference from the total fuel consumption reported in the national energy balance and the estimated values as described above and emissions are therefore calculated.

For maritime transportation only by Directive 1999/32/EC European Union started to examine environmental impact of navigation and in particular the sulphur content of fuels. This directive was amended by Directive 2005/33/EC that designated the Baltic Sea, the English Channel and the North Sea as sulphur emission control areas (SECA) limiting the content of sulphur in the fuel for these areas and introducing a limit of 0.1% of the sulphur content in the fuel used in EU harbours from 2010.

EU legislation combined with national normative resulted in the introduction of a limit of sulphur content in maritime gasoil equal to 0.2% (2% before) from 2002 and 0.1% from 2010 while for fuel oil some limits occur only from 2008 (maximum sulphur content of 1.5 % in harbour) and from 2010, 2% in domestic waters and 1% in harbour. For inland waterways, which include the navigation on the Po river and ferry-boats in the Venice lagoon, the same legislation is applied. Moreover, since January 2020, IMO introduced a limit of sulphur content in marine fuel oil equal to 0.5% (previously equal to 3.5%) in IMO jurisdiction areas (except in ECAS areas, which have a stricter sulphur content limit, equal to 0.1%).

The composition of the fleet of gasoline fuelled recreational craft distinguished in two strokes and four strokes engine distribution is provided by the industrial category association (UCINA, several years); the trend of the average emission factors takes into account the switch from two strokes to four strokes engines of the national fleet due to the introduction in the market of new models. In 2000, the composition of the fleet was 90% two stroke engine equipped and 10% four stroke while in the last year four strokes engines are about 56 % of the fleet.

The fuel split between national and international fuel use in maritime transportation is then supplied to the Ministry of Economic Development to be included in the official international submission of energy statistics to the IEA in the framework of the Joint Questionnaire OECD/EUROSTAT/IEA compilation together with other energy data. A discrepancy with the international bunkers reported to the IEA still remains, especially for the nineties, because the time series of the energy statistics to the IEA are not updated.

PCB, HCB and Dioxins emissions are estimated with Tier1 emission factors available in the 2019 EMEP/EEA Guidebook (EMEP/EEA, 2019).

3.7.3 Time series and key categories

In Table 3.32 the list of key categories by pollutant identified for navigation in 2020, 1990 and at trend assessment is reported. Navigation is, in 2020, key category for many pollutants: SO_x, NO_x, PM10, PM2.5, BC, CO; furthermore, it is a key driver of the SO_x and NO_x trend.

Table 3.32 List of key categories for pollutant in navigation in 2020, 1990 and in the trend

| | Key categories in 2020 | Key categories in 1990 | Key categories in trend |
|-----------------|------------------------|------------------------|-------------------------|
| SO _x | 1A3dii | 1A3dii | 1A3dii |
| NO _x | 1A3dii | 1A3dii | 1A3dii |
| PM10 | 1A3dii | 1A3dii | |
| PM2.5 | 1A3dii | 1A3dii | |
| BC | 1A3dii | | |
| CO | 1A3dii | | |

Source: ISPRA elaborations

Estimates of fuel consumption for domestic use, in the national harbours or for travel within two Italian destinations, and bunker fuels used for international travels are reported in Table 3.33.

An upward trend in emission levels is observed from 1990 to 2000, explained by the increasing number of ship movements. Nevertheless, the operational improvements in harbour activities and a reduction in ship domestic movements inverted the tendency in the last years.

Table 3.33 Marine fuel consumptions in domestic navigation and international bunkers (Gg) and pollutants emissions from domestic navigation (Gg)

| Consumptions and Emissions for NFR Subsector 1.A.3.d | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|----------|
| Gasoline for recreational craft (Gg) | 182.12 | 210.14 | 213.14 | 199.13 | 169.11 | 90.39 | 89.63 | 88.22 | 87.67 | 86.65 | 85.59 |
| Diesel oil for inland waterways (Gg) | 19.81 | 22.74 | 20.21 | 24.76 | 18.19 | 27.45 | 26.92 | 28.69 | 27.63 | 24.02 | 19.19 |
| Fuels used in domestic cruise navigation (Gg) | 778.06 | 706.38 | 811.37 | 739.97 | 725.35 | 525.27 | 558.15 | 533.24 | 577.86 | 621.10 | 667.01 |
| Fuel in harbours (dom+int ships) (Gg) | 748.46 | 692.95 | 818.48 | 758.89 | 743.90 | 538.70 | 546.88 | 592.64 | 592.64 | 636.98 | 684.06 |
| Fuel in international Bunkers (Gg) | 1,402.72 | 1,287.30 | 1,306.31 | 2,147.25 | 2,174.64 | 1,782.46 | 2,075.34 | 2,266.22 | 2,242.54 | 2,072.07 | 1,751.91 |
| Emissions from National Navigation (Gg) | | | | | | | | | | | |
| Emissions of NOx | 95.55 | 87.97 | 102.48 | 94.94 | 93.28 | 68.06 | 72.17 | 69.17 | 74.74 | 80.02 | 85.59 |
| Emissions of NMVOC | 46.11 | 52.42 | 50.17 | 43.28 | 31.82 | 15.31 | 14.92 | 14.12 | 13.81 | 13.39 | 13.00 |
| Emissions of SOx | 77.94 | 70.31 | 81.49 | 49.73 | 28.38 | 20.55 | 21.84 | 20.86 | 22.61 | 24.30 | 8.06 |
| Emissions of PM2.5 | 9.30 | 8.83 | 9.61 | 8.90 | 7.86 | 5.33 | 5.58 | 5.33 | 5.69 | 6.01 | 6.36 |
| Emissions of PM10 | 9.33 | 8.86 | 9.65 | 8.94 | 7.89 | 5.35 | 5.61 | 5.36 | 5.71 | 6.04 | 6.39 |
| Emissions of BC | 1.33 | 1.25 | 1.40 | 1.31 | 1.23 | 0.90 | 0.95 | 0.91 | 0.98 | 1.03 | 1.09 |
| Emissions of CO | 102.27 | 115.57 | 124.77 | 122.86 | 109.42 | 58.33 | 57.53 | 55.55 | 54.84 | 53.75 | 52.61 |

Source: ISPRA elaborations

3.7.4 QA/QC and Uncertainty

Basic data to estimate emissions are reconstructed starting from information on ship movements and fleet composition coming from different sources. Data collected in the framework of the national study from the local port authorities, carried out in 2009 (TECHNE, 2009), were compared with the official statistics supplied by ISTAT, which are collected from maritime operators with a yearly survey and communicated at international level to EUROSTAT. Differences and problems were analysed in details and solved together with ISTAT experts. Different sources of data are usually used and compared during the compilation of the annual inventory.

Besides, time series resulting from the recalculation have been presented to the national experts in the framework of an *ad hoc* working group on air emissions inventories. The group is chaired by ISPRA and includes participants from the local authorities responsible for the preparation of local inventories, sectoral experts, the Ministry of Environment, Land and Sea, and air quality model experts. Top-down and bottom-up approaches have been compared with the aim to identify the potential problems and future improvements to be addressed. There is also an ongoing collaboration and data exchange with regional environmental agencies on this issue.

3.7.5 Recalculations

A recalculation occurred for total fuel consumption from 2012. Moreover, the total number of ships arrived has been updated from 2017. The emission estimates have been consequently recalculated.

3.7.6 Planned improvements

Further improvements will include a verification of activity data on ship movements and emission estimates with regional environmental agencies, especially with those more affected by maritime pollution.

3.8 PIPELINE COMPRESSORS (NFR SUBSECTOR 1.A.3.E)

Pipeline compressors category (1.A.3e) includes all emissions from fuels delivered to the transportation by pipelines and storage of natural gas. Relevant pollutant emissions typical of a combustion process, such as SO_x, NO_x, CO and PM emissions, derive from this category. This category is not a key category.

Emissions from pipeline compressors are estimated on the basis of natural gas fuel consumption used for the compressors and the relevant emission factors. The amount of fuel consumption is estimated on the basis of data supplied for the whole time series by the national operators of natural gas distribution (SNAM and STOGIT) and refers to the fuel consumption for the gas storage and transportation; this consumption is part of the fuel consumption reported in the national energy balance in the consumption and losses sheet. Emission factors are those reported in the EMEP/EEA Guidebook for gas turbines (EMEP/CORINAIR, 2007). Emissions communicated by the national operators in their environmental reports are also taken into account to estimate air pollutants, especially SO_x, NO_x, CO and PM10.

Regarding QA/QC, fuel consumptions reported by the national operators for this activity are compared with the amount of natural gas internal consumption and losses reported in the energy balance as well as with energy consumption data provided by the operators to the emission trading scheme.

Starting from the length of pipelines, the average energy consumptions by kilometre are calculated and used for verification of data collected by the operators. Energy consumptions and emissions by kilometre calculated on the basis of data supplied by SNAM, which is the main national operator, are used to estimate the figures for the other operators when their annual data are not available.

No recalculations occurred with respect the previous submission.

In Table 3.34, nitrogen oxides, non-methane volatile organic compounds, sulphur oxides, particulate and carbon monoxide emissions (Gg) are reported.

Table 3.34 Emissions from pipeline compressors (Gg)

| Emissions for NFR Subsector 1.A.3.e | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|------|------|------|------|------|------|------|------|-------|-------|-------|
| NO _x | 2.89 | 4.18 | 2.96 | 2.37 | 1.71 | 0.36 | 0.46 | 0.55 | 0.60 | 0.43 | 0.38 |
| NM VOC | 0.02 | 0.03 | 0.04 | 0.04 | 0.05 | 0.02 | 0.03 | 0.03 | 0.03 | 0.03 | 0.03 |
| SO _x | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.005 | 0.005 | 0.005 |
| PM10 | 0.02 | 0.03 | 0.05 | 0.03 | 0.03 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 |
| CO | 1.26 | 1.38 | 1.03 | 0.60 | 0.61 | 0.23 | 0.30 | 0.35 | 0.21 | 0.18 | 0.16 |

Source: ISPRA elaborations

3.9 CIVIL SECTOR: SMALL COMBUSTION AND OFF-ROAD VEHICLES (NFR SUBSECTOR 1.A.4 - 1.A.5)

3.9.1 Overview

Emissions from energy use in the civil sector cover combustion in small-scale combustion units, with thermal capacity < 50 MW_{th}, and off road vehicles in the commercial, residential and agriculture sectors.

The emissions refer to the following categories:

- 1 A 4 a i Commercial / Institutional: Stationary
- 1 A 4 a ii Commercial / Institutional: Mobile
- 1 A 4 b i Residential: Stationary plants
- 1 A 4 b ii Residential: Household and gardening (mobile)
- 1 A 4 c i Agriculture/Forestry/Fishing: Stationary
- 1 A 4 c ii Agriculture/Forestry/Fishing: Off-road Vehicles and Other Machinery
- 1A 4 c iii Agriculture/Forestry/Fishing: National Fishing
- 1 A 5 a Other, Stationary (including military)
- 1 A 5 b Other, Mobile (Including military, land based and recreational boats)

In Table 3.35 the list of categories for small combustion and off-road vehicles identified as key categories by pollutant for 2020, 1990 and in the trend is reported.

Table 3.35 List of key categories by pollutant in the civil sector in 2020, 1990 and trend

| | Key categories in 2020 | | Key categories in 1990 | | Key categories in trend | | |
|-----------------|------------------------|-----------|------------------------|----------------------|-------------------------|-----------|----------------------|
| SO _x | 1 A 4 b i | 1 A 4 a i | | 1 A 4 b i | | 1 A 4 a i | |
| NO _x | 1 A 4 b i | 1 A 4 a i | 1 A 4 c ii | 1 A 4 c ii | | 1 A 4 a i | 1 A 4 b i 1 A 4 c i |
| NMVOC | 1 A 4 b i | 1 A 4 a i | | 1 A 4 b i 1 A 4 c ii | | 1 A 4 b i | 1 A 4 a i 1 A 4 c ii |
| CO | 1 A 4 b i | | | 1 A 4 b i 1 A 4 c ii | | 1 A 4 b i | |
| PM10 | 1 A 4 b i | | | 1 A 4 b i 1 A 4 c ii | | 1 A 4 b i | 1 A 4 c ii |
| PM2.5 | 1 A 4 b i | | | 1 A 4 b i 1 A 4 c ii | | 1 A 4 b i | 1 A 4 c ii |
| BC | 1 A 4 b i | | 1 A 4 c ii | 1 A 4 c ii 1 A 4 b i | | 1 A 4 b i | 1 A 4 c ii |
| Pb | 1 A 4 b i | | | | | | |
| Cd | 1 A 4 b i | | | 1 A 4 b i 1 A 4 a i | | | |
| PAH | 1 A 4 b i | | | 1 A 4 b i | | 1 A 4 b i | |
| DIOX | 1 A 4 b i | | | 1 A 4 a i 1 A 4 b i | | 1 A 4 b i | 1 A 4 a i |
| HCB | 1 A 4 b i | 1 A 4 a i | | | | 1 A 4 a i | 1 A 4 b i |
| PCB | 1 A 4 b i | | | | | 1 A 4 b i | |

3.9.2 Activity data

The Commercial / Institutional emissions arise from the energy used in the institutional, service and commercial buildings, mainly for heating. Additionally, this category includes all emissions due to wastes used in electricity generation as well as biogas recovered in landfills and wastewater treatment plant. In the residential sector the emissions arise from the energy used in residential buildings, mainly for heating and the sector includes emissions from household and gardening machinery. The Agriculture/ Forestry/ Fishing sector includes all emissions due to the fuel, including biogas from biodigestors, used in agriculture, mainly to produce mechanical energy, the fuel use in fishing and for machinery used in the forestry sector. Emissions from military aircraft and naval vessels are reported under 1A.5.b Mobile.

Emissions from 1.A.4.a ii are reported as IE, included elsewhere, because of they refer to road transport emissions of institutional and commercial vehicles. These emissions are estimated, and reported in 1.A.3.b, with a model (COPERT 5) which consider the vehicle fleet subdivided by technology and fuel and not by user. Emissions from 1.A.5.a are also reported as IE because they refer to stationary combustion in commercial and residential of military which are included and reported in 1.A.4.a i and 1.A.4.b i; also in this case the relevant energy statistics are not available by user.

The estimation procedure follows that of the basic combustion data sheet. Emissions are estimated from the energy consumption data that are reported in the national energy balance (MSE, several years (a)) and separating energy consumption between commercial/institutional, residential, agriculture and fishing,

according to the information available in the Joint Questionnaire OECD/IEA/EUROSTAT prepared by the Ministry of Economic Development and officially sent to the international organizations.

Emissions from 1.A.4.b Residential and 1.A.4.c Agriculture/Forestry/Fishing are disaggregated into those arising from stationary combustion and those from off-road vehicles and other machinery. The time series of fuel consumption for the civil sector are reported in Table 3.36.

Table 3.36 *Time series of fuel consumption for the civil sector*

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|---------|
| | TJ | | | | | | | | | | |
| 1 A 4 a i Commercial / Institutional: Stationary plants | 206,427 | 247,440 | 306,088 | 419,507 | 489,018 | 400,596 | 402,607 | 402,594 | 428,941 | 421,094 | 402,697 |
| 1 A 4 b i Residential: Stationary plants | 1,002,131 | 1,003,620 | 1,036,801 | 1,172,245 | 1,222,522 | 1,078,829 | 1,070,709 | 1,093,472 | 1,045,161 | 1,019,739 | 997,306 |
| 1 A 4 b ii Residential: Household and gardening (mobile) | 466 | 571 | 374 | 154 | 66 | 57 | 35 | 35 | 35 | 35 | 44 |
| 1 A 4 c i Agriculture/Forestry/Fishing: Stationary | 9,688 | 9,487 | 8,152 | 10,178 | 8,834 | 8,241 | 6,407 | 6,570 | 6,650 | 6,925 | 6,741 |
| 1 A 4 c ii Agriculture/Forestry/Fishing: Off-road Vehicles and Other Machinery | 96,536 | 101,928 | 94,668 | 95,869 | 84,461 | 81,263 | 83,116 | 83,543 | 88,497 | 83,276 | 83,991 |
| 1 A 4 c iii Agriculture/Forestry/Fishing: National Fishing | 8,413 | 9,651 | 8,584 | 10,464 | 7,731 | 6,194 | 6,918 | 6,662 | 7,046 | 6,630 | 6,688 |
| 1 A 5 b Other, Mobile (Including military, land based and recreational boats) | 14,840 | 20,814 | 11,595 | 16,947 | 9,001 | 6,388 | 7,183 | 4,531 | 4,754 | 6,317 | 8,733 |

3.9.3 Methodological issues

The Tier 2 methodology is applied to the whole category. Emissions are estimated for each fuel and category at detailed level and country specific emission factors are used for the key fuel and categories drivers of total emission trend.

More in detail, 1.A.4.a i, is key category in 2020 and in trend for SO_x, NO_x, NMVOC and HCB emissions as well as for cadmium and Dioxin in 1990, and dioxins in trend analysis. Most of these pollutants are due prevalently to emissions from waste incineration with energy recovery (more than 90% for HCB, PCB and HMs with the exception of Hg (74%) and Ni (83%), around 90% for NMVOC and SO_x and 28% of the total for NO_x). Emissions from waste combustion in incinerator with energy recovery have been calculated with a Tier 3 methodology from the database of incinerator plants which includes plant specific emission factors on the basis of their technology and measurements data (ENEA-federAmbiente, 2012). The methodology used to estimate emissions from incinerators is reported in the relevant paragraph on waste incineration in the waste sector, and in particular EFs are reported in Table 7.3. Up to 2009 emission factors have been estimated on the basis of a study conducted by ENEA (De Stefanis P., 1999), based on emission data from a large sample of Italian incinerators (FEDERAMBIENTE, 1998; AMA-Comune di Roma, 1996), legal thresholds (Ministerial Decree 19 November 1997, n. 503 of the Ministry of Environment; Ministerial Decree 12 July 1990) and expert judgements.

Waste management with incinerators is a commercial activity with recover of the energy auto-produced and emissions from these plants are allocated in the commercial / institutional category because of the final use of heat and electricity production. In fact, until the early 2000s, electricity and heat produced by incinerators have been prevalently used to satisfy the energy demand from connected activities: heating of buildings, domestic hot water and electricity for offices. This is still true in particular for industrial and hospital incinerators, meanwhile municipal solid waste incinerators have increased the amount of energy provided to the grid from the early 2000s until now, although only a small percentage of energy produced goes to the electricity grid (around 10%); the energy recovered by these plants is mainly used for district heating of commercial buildings or used to satisfy the internal energy demand of the plants. Since 2010, emission factors for urban waste incinerators have been updated on the basis of data provided by plants (ENEA-federAmbiente, 2012; De Stefanis P., 2012) concerning the annual stack flow, the amount of waste burned and the average concentrations of the pollutants at the stack and taking in account the abatement technologies in place. As the emission factors

are considerably lower than the old ones due to the application of very efficient abatement systems it was necessary to apply a linear smoothing methodology assuming a progressive application of the abatement systems between 2005 and 2010. In a similar way, emission factors for industrial waste incinerators have been updated from 2010 onwards on the basis of the 2019 EMEP/EEA Guidebook. Similarly, to municipal waste smoothing has been applied between 2005 and 2010 supposing a linear application of the abatement systems.

The other fuels driving emissions from this category are wood combustion, especially for NMVOC, and natural gas for NO_x while the trend of SO_x is driven by the decrease both of liquid fuel, as gasoil, fuel oil and kerosene, consumptions and their sulphur content which is also decreased according to European Union and national legislation. For what concerns wood combustion, the NMVOC average emission factor, as well as all the other pollutants, takes into account the different technologies used and is calculated on the basis of country specific emission factors and the ranges reported in the 2019 EMEP/EEA Guidebook; see paragraph 3.9.3.2 for details on methodology and emission factors. For natural gas and NO_x emissions, a Tier 2 methodology is used and country specific emission factors as described in the following paragraph 3.9.3.1. For the other fuels the default emission factors of EMEP/CORINAIR 2007 Guidebook have been used; it is planned to update these emission factors with those reported in the 2019 EMEP/EEA Guidebook and, according to the last review process, the update of EFs started from the current report with COVNM emissions. For gasoil, biogas and gasoline different emission factors are used for stationary engines and boilers.

Concerning the other pollutants, PM_{2.5} emissions from wood account for around 80% of the total 1.A.4.a i category; the other main fuel used for this category is biogas from landfills and wastewater treatment energy recovery, which account for around 10% of PM_{2.5} emissions of this category; an emission factor equal to 10 g/GJ is used. The other fuels have been estimated with EMEP/CORINAIR 2007 emission factors. For NO_x, in addition to waste fuel, see methodology in the waste chapter and in particular emission factors reported in Table 7.3, and natural gas, as described in the following paragraph 3.9.3.1, the other main fuel driving emission estimates is biogas from landfills and wastewater treatment energy recovery, accounting for 43% of NO_x emissions of this category in 2020, but for which no guidance is provided in the Guidebook. An emission factor equal to 1 kg/GJ has been used taking into account that the gas is burnt in stationary engines. These fuels plus wood that is also estimated with a Tier 2 account for more than 72% of total NO_x category emissions. HM and POP emissions from the sector are prevalently from waste incineration, estimated with country specific EFs, at technology level, and from wood combustion estimated also with country specific EFs in the range of 2019 EMEP/EEA Guidebook values.

For 1.A.4.b i, the category is key category in 2020 for SO_x, NO_x, NMVOC, CO, PM₁₀, PM_{2.5}, BC, Pb, Cd, PAH, Dioxin, HCB and PCB emissions. Most of these pollutants are also key in 1990 and for trend analysis. Most of these pollutants are due prevalently to emissions from wood combustion (more than 99% for PM, BC, PAH, Se, Zn, HCB and PCB emissions, 98% for dioxins, Cr and Pb, more than 90% CO and NMVOC, around 66% for SO_x and 43% of the total for NO_x) for which a Tier 2 is applied. Methodology and emission factors are described in paragraph 3.9.3.2.

For SO_x country specific and updated emission factors are used for wood, gasoil, residual oil, natural gas and LPG calculated on the basis of the maximum content of sulphur in these fuels; emissions from these fuels account for about 99.9% of SO_x category emissions.

A country specific methodology has been developed and applied to estimate NO_x emissions from gas powered plants and all emissions from wood combustion. More than 50% of the total emissions are due to the combustion of natural gas; methodology and country specific emission factors are described in the following paragraph 3.9.3.1. Biomass combustion accounts for around 43% of the total and methodology and country specific emission factors are also available in paragraph 3.9.3.2. For the other fuels, the default EMEP/CORINAIR 2007 Guidebook values have been used. In particular for liquid fuels (gasoil, kerosene and LPG) a default equal to 50 g/GJ is used. All these fuels cover more than 99% of total category emissions. According to the previous review process, an update of EFs occurs for NO_x, NMVOC and PM_{2.5} since 2000.

For 1.A.4.bii, 1.A.4.cii, 1.A.4.ciii and 1.A.5b emission estimates are calculated taking into account the relevant changes in emission factors along the time series due to the introduction of the relevant European Union Directives for off-road engines. Regarding mobile machinery used in agriculture, forestry and household, these sectors were not governed by any legislation until the Directive 97/68/EC (EC, 1997 [a]), which provides for a reduction in NO_x limits from 1st January 1999, and Directive 2004/26/EC (EC, 2004) which provide further reduction stages with substantial effects from 2011, with a following decreasing trend particularly in recent years. For engines with lower power as those used in forestry, household and gardening, the European Directives introduce emissions limits only starting from 2019 and 2021 so they have not had

effect up to now. Moreover, for the category 1.A.4.bii, 1.A.4.cii and 1.A.4.ciii, Pb emissions from 2002 are reported as 'NA', because of the introduction of unleaded liquid fuels in the market in 2002. In particular heavy metals contents values derive from the analysis about the physical - chemical characterization of fossil fuels used in Italy (Innovhub, Fuel Experimental Station, several years). According to the review (EEA, 2019), PCB, HCB and Dioxins emissions have been estimated and included in the inventory for 1.A.4.ciii category with the emission factors of Tier1 available in the 2019 EMEP/EEA Guidebook.

3.9.3.1 *NO_x emissions from gas powered plants in the civil sector*

A national methodology has been developed and applied to estimate NO_x emissions from gas powered plants in the civil sector, according to the EMEP/EEA Guidebook (EMEP/EEA, 2019).

On the basis of the information and data reported in available national studies for the year 2003, a distribution of heating plants in the domestic sector by technology and typology has been assessed for that year together with their specific emissions factors. Data related to heating plants, both commercial and residential, have been supplied for 2003 by a national energy research institute (CESI, 2005). In this study, for the residential sector, the sharing of single and multifamily houses plants by technology and a quantitative estimation of the relevant gas powered ones are reported, including their related NO_x emission factors. Domestic final consumption by type of plant, single or multifamily plants, has been estimated on the basis of data supplied by ENEA on their distribution (ENEA, several years).

Data reported by ASSOTERMICA (ASSOTERMICA, several years) on the number of heating plants sold are used for the years after 2003 to update the information related to the technologies. A linear regression, for the period 1995-2003, has been applied, while for the period 1990-1994, the technology with the highest emission factor has been assumed to be operating.

In Table 3.37 the time series of NO_x average emission factors for the relevant categories is reported.

Table 3.37 *Time series of NO_x emissions factor for the civil sector*

| EF NO _x | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|------|------|------|------|------|------|------|------|------|------|------|
| | g/Gj | | | | | | | | | | |
| 1 A 4 a i Commercial / Institutional: Stationary | 50 | 48.5 | 40.2 | 35.2 | 32.4 | 30.3 | 29.9 | 29.5 | 29.1 | 28.8 | 28.4 |
| 1 A 4 b i Residential: Stationary plants | 50 | 48.1 | 38.2 | 31.7 | 31.1 | 30.5 | 30.3 | 29.9 | 29.6 | 29.6 | 29.6 |

3.9.3.2 *Emissions from wood combustion in the civil sector*

A national methodology has been developed and applied to estimate emissions from wood combustion in the civil sector, according to the TIER 2 methodology reported in the EMEP/EEA Guidebook (EMEP/EEA, 2019). In the past years, several surveys have been carried out to estimate national wood consumption in the domestic heating and the related technologies used. In the estimation process, three surveys have been taken into account: the first survey (Gerardi and Perrella, 2001) has evaluated the technologies for wood combustion used in Italy for the year 1999, the second survey (ARPA, 2007) was related to the year 2006, while the third survey (SCENARI/ISPRA, 2013) was related to the year 2012.

For 2015 and 2019 information on the use of pellet, as available in the national energy balance, and on the relevant technologies, as provided by the industrial association, has been used to take in account the increase of pellet used for heating; the update has been developed taking in account also the results of the surveys on wood consumption and combustion technologies carried out by ISPRA (SCENARI/ISPRA, 2013) and by ISTAT (ISTAT, 2014).

The technologies assessed by the abovementioned surveys and the distribution of fuel combustion by technologies are reported in Table 3.38.

Table 3.38 *Distribution of wood combustion by technologies*

| Distribution of wood combustion by technologies | | | | | | | |
|---|------|------|------|------|------|------|------|
| | 1999 | 2006 | 2012 | 2015 | 2018 | 2019 | 2020 |
| | % | | | | | | |
| Fireplaces | 51.3 | 44.7 | 51.2 | 49.0 | 41.0 | 40.3 | 39.7 |

| Distribution of wood combustion by technologies | | | | | | | |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | 1999 | 2006 | 2012 | 2015 | 2018 | 2019 | 2020 |
| Stoves | 28.4 | 27.6 | 22.9 | 21.0 | 19.0 | 18.4 | 17.9 |
| Advanced fireplaces | 15.4 | 20.2 | 15.8 | 15.0 | 20.0 | 19.7 | 19.5 |
| Pellet stoves | 0 | 3.1 | 4.0 | 9.0 | 12.0 | 13.6 | 15.1 |
| Advanced stoves | 4.8 | 4.4 | 6.0 | 6.0 | 8.0 | 7.9 | 7.8 |

Average emission factors for 1999, 2006, 2012, 2015, 2018, 2019 and 2020 have been estimated at national level taking into account the technology distributions; for 1990 only old technologies (fireplaces and stoves) have been considered and linear regressions have been applied to reconstruct the time series from 1990 to 2006. For the years till 2011, emission factors from 2006 have been used in absence of further available information. The distribution of combustion technologies is updated, starting from this year, on an annual basis based on the sales data of the equipment by type.

For NMVOC, PAH, PM10 and PM2.5 emission factors the results of the experimental study funded by the Ministry of Environment and conducted by the research institute ‘Stazione Sperimentale dei Combustibili’ now Innovhub (SSC, 2012; INNOVHUB, 2021) have been used. This study measured and compared NO_x, CO, NMVOC, SO_x, TSP, PM10, PM2.5, PAH and Dioxin emissions for the combustion of different wood typically used in Italy as beech, hornbeam, oak, locust and spruce-fir, in open and closed fireplaces, traditional and innovative stoves, and pellet stoves. Emissions from certificated and not certificated pellets have been also measured and compared. In general, measured emission factors results in the ranges supplied by the EMEP/EEA Guidebook but for some pollutants and technologies results are sensibly different. In particular NMVOC emissions for all the technologies are close or lower to the minimum value of the range reported in the Guidebook, as well as PM emissions with exception of emissions from pellet stoves which are higher of the values suggested in the case of the use of not certificated pellet. For these pollutants the minimum values of the range in the Guidebook have been used when appropriate. For that concern PAH, measured emissions from open fireplaces are much lower than the minimum value of the range in the Guidebook while those from the advanced stoves are close to the superior values of the range for all the PAH compounds. In this case, for open fireplaces, experimental values have been used while for the other technologies the minimum or maximum values of the range in the Guidebook have been used as appropriate. For the other pollutants where differences with the values suggested by the Guidebook are not sensible, a more in-depth analysis will be conducted with the aim to update the emission factors used if needed. During 2020, a new experimental study funded by the Ministry of Environment and managed by ISPRA has been completed (INNOVHUB, 2021). This study regards the analysis on advanced appliances burning solid biomass (beech, fir and hornbeam, pellet A1, pellet A2). The pollutants that have been monitored are: CO, NO_x, SO₂, PM, PAH. The study also contains an interesting comparison between standard methodology and the BeReal method.

In Table 3.39 emission factors used for the Italian inventory are reported.

Table 3.39 Emission factors for wood combustion

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2018 | 2019 | 2020 |
|-------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | g/Gj | | | | | | | | |
| NO _x | 50 | 55 | 59 | 61 | 61 | 61 | 61 | 66 | 66 |
| CO | 6000 | 5791 | 5591 | 5427 | 5395 | 5010 | 5010 | 4633 | 4564 |
| NMVOC | 762 | 715 | 672 | 643 | 638 | 597 | 597 | 535 | 526 |
| SO ₂ | 10 | 11 | 12 | 13 | 13 | 13 | 13 | 14 | 14 |
| NH ₃ | 9 | 7 | 6 | 6 | 6 | 6 | 6 | 5 | 5 |
| PM10 | 507 | 465 | 428 | 409 | 406 | 392 | 392 | 352 | 348 |
| PM2.5 | 503 | 461 | 424 | 405 | 402 | 388 | 388 | 348 | 344 |
| BC | 40 | 37 | 35 | 34 | 34 | 33 | 33 | 31 | 31 |
| PAH | 0.25 | 0.24 | 0.23 | 0.23 | 0.22 | 0.21 | 0.21 | 0.19 | 0.19 |
| Dioxin (µg/GJ) | 0.48 | 0.47 | 0.45 | 0.44 | 0.43 | 0.40 | 0.40 | 0.37 | 0.36 |
| PCB | 0.00006 | 0.00006 | 0.00006 | 0.00006 | 0.00006 | 0.00006 | 0.00006 | 0.00006 | 0.00006 |
| HCB | 0.00001 | 0.00001 | 0.00001 | 0.00001 | 0.00001 | 0.00001 | 0.00001 | 0.00001 | 0.00001 |

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2018 | 2019 | 2020 |
|-------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| | g/Gj | | | | | | | | |
| As | 0.001 | 0.001 | 0.001 | 0.001 | 0.0005 | 0.0005 | 0.0005 | 0.0005 | 0.0005 |
| Cd | 0.002 | 0.002 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 | 0.001 |
| Cr | 0.001 | 0.002 | 0.003 | 0.003 | 0.003 | 0.003 | 0.003 | 0.003 | 0.003 |
| Cu | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| Hg | 0.0004 | 0.0004 | 0.0004 | 0.0004 | 0.0004 | 0.0004 | 0.0004 | 0.0004 | 0.0004 |
| Ni | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 |
| Pb | 0.04 | 0.04 | 0.04 | 0.04 | 0.04 | 0.04 | 0.04 | 0.03 | 0.03 |
| Se | 0.001 | 0.001 | 0.001 | 0.001 | 0.0005 | 0.0005 | 0.0005 | 0.0005 | 0.0005 |
| Zn | 0.10 | 0.10 | 0.10 | 0.09 | 0.09 | 0.09 | 0.09 | 0.09 | 0.09 |
| B(a)P | 0.07 | 0.07 | 0.07 | 0.07 | 0.07 | 0.06 | 0.06 | 0.06 | 0.06 |
| B(b)F | 0.09 | 0.08 | 0.08 | 0.08 | 0.08 | 0.07 | 0.07 | 0.07 | 0.07 |
| B(k)F | 0.04 | 0.04 | 0.04 | 0.04 | 0.03 | 0.03 | 0.03 | 0.03 | 0.03 |
| IND | 0.05 | 0.05 | 0.05 | 0.05 | 0.04 | 0.04 | 0.04 | 0.04 | 0.04 |

In 2014 the national Institute of Statistics (ISTAT) carried out a survey, funded by the Ministry of Economic Development and infrastructure (MSE), on the final energy consumption of households for residential heating which include the fuel consumption of solid biomass, as wood and pellets (ISTAT, 2014). In this regard the survey resulted in an official statistic for 2012 and 2013 of wood and pellet fuel consumption at national and regional level including the information on the relevant equipment. The resulting figure for 2013 doubled the value reported in the National Energy Balance for previous years which asked for the need to update the whole time series. An *ad hoc* working group has been established, involving ISPRA, MSE and the energy management system national operator (GSE), to reconstruct the complete time series of wood and pellet fuel consumption which has been recalculated and officially submitted to Eurostat in June 2015.

The methodology to recalculate consumption figures has taken in account the amount of wood harvested for energy purposes, the amount of wood biomass from pruning, import and export official statistics to estimate total wood consumption. A model to estimate the annual amount of wood for heating has been developed on the basis of the annual energy total biomass demand of households estimated considering the degree days time series, the number of households, the energy efficiency of equipment and fuel consumption statistics for the other fuels. As a consequence, time series for residential heating have been completely recalculated affecting the relevant pollutants and resulting in important recalculations at national total levels.

3.9.4 Time series and key categories

The time series of emissions for civil sector shows an increasing trend for several pollutants, except for SO_x and NO_x, due to a gradual shift from diesel fuel to gas, concerning SO_x, and to a replacement of classic boilers with those with low NO_x emission. Many other pollutants have a growing trend, as a consequence of the increase of wood combustion.

In particular the pollutants which are more affected by the increase of wood biomass in this category according to data available in the National Energy Balance are PM, PAH, NMVOC and CO. In particular for 1.A.4.c i the increasing trend of PAH in the last years is due to the increase of wood combustion for this category.

More in detail the decrease of SO_x emissions is the combination of the switch of fuel from gasoil and fuel oil to natural gas and LPG and the reduction in the average sulphur content of liquid fuels. The SO_x emission factors for 1990 and 2020 by fuels are shown in the following box.

| EMISSION FACTORS (kg/Gj) | | |
|---------------------------------|-------------|-------------|
| FUEL | 1990 | 2020 |
| steam coal | 0.646 | 0.646 |
| coke oven coke | 0.682 | 0.682 |
| wood and similar | 0.010 | 0.014 |
| municipal waste | 0.069 | 0.048 |
| biodiesel | 0.047 | 0.047 |
| residual oil | 1.462 | 0.146 |
| gas oil | 0.140 | 0.047 |

| EMISSION FACTORS (kg/Gj) | | |
|---------------------------------|-------------|-------------|
| FUEL | 1990 | 2020 |
| kerosene | 0.018 | 0.018 |
| natural gas | 0.0003 | 0.0003 |
| biogas | - | - |
| LPG | 0.0022 | 0.0022 |
| gas works gas | 0.011 | 0.011 |
| motor gasoline | 0.023 | 0.023 |

Time series of emissions is reported in Table 3.40.

Table 3.40 Time series of emissions in civil sector: small combustion and off-road vehicles

| | | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---------------------------|-----|---------|---------|---------|--------|---------|---------|---------|---------|---------|---------|---------|
| SO _x (Gg) | 1A4 | 96.47 | 42.66 | 26.37 | 22.84 | 12.12 | 10.32 | 10.32 | 10.15 | 10.43 | 9.99 | 9.81 |
| | 1A5 | 1.19 | 0.81 | 0.21 | 0.17 | 0.13 | 0.12 | 0.15 | 0.08 | 0.10 | 0.13 | 0.19 |
| NO _x (Gg) | 1A4 | 174.96 | 187.20 | 171.81 | 163.04 | 144.04 | 127.02 | 126.56 | 125.01 | 123.89 | 118.84 | 113.86 |
| | 1A5 | 11.16 | 11.99 | 7.24 | 13.50 | 6.11 | 3.29 | 3.28 | 2.36 | 2.05 | 2.73 | 3.48 |
| CO (Gg) | 1A4 | 1093.21 | 1088.79 | 1028.00 | 997.44 | 1716.14 | 1440.68 | 1392.68 | 1515.62 | 1331.32 | 1307.68 | 1244.95 |
| | 1A5 | 65.12 | 79.02 | 45.49 | 54.48 | 17.33 | 16.49 | 19.73 | 11.93 | 13.23 | 17.42 | 24.93 |
| PM ₁₀ (Gg) | 1A4 | 84.63 | 89.76 | 84.57 | 80.33 | 129.12 | 110.17 | 106.56 | 115.77 | 97.69 | 96.05 | 91.58 |
| | 1A5 | 1.27 | 1.54 | 0.90 | 1.60 | 0.81 | 0.47 | 0.49 | 0.34 | 0.31 | 0.42 | 0.55 |
| PM _{2.5} (Gg) | 1A4 | 83.81 | 89.07 | 83.93 | 79.63 | 127.78 | 108.96 | 105.39 | 114.49 | 96.50 | 94.87 | 90.45 |
| | 1A5 | 1.27 | 1.54 | 0.90 | 1.60 | 0.81 | 0.47 | 0.49 | 0.34 | 0.31 | 0.42 | 0.55 |
| BC (Gg) | 1A4 | 14.90 | 16.14 | 14.64 | 12.41 | 13.74 | 10.99 | 10.57 | 11.17 | 9.86 | 9.51 | 8.98 |
| | 1A5 | 0.72 | 0.82 | 0.49 | 0.92 | 0.46 | 0.25 | 0.25 | 0.18 | 0.16 | 0.21 | 0.27 |
| Pb (Mg) | 1A4 | 81.95 | 34.29 | 24.64 | 46.34 | 16.46 | 15.11 | 14.56 | 15.24 | 14.44 | 14.09 | 13.62 |
| | 1A5 | 16.34 | 4.22 | 1.16 | 0.00 | NA | 0.12 | 0.02 | 0.02 | 0.09 | 0.18 | 0.18 |
| Cd (Mg) | 1A4 | 1.51 | 1.21 | 1.74 | 2.62 | 0.66 | 0.55 | 0.53 | 0.55 | 0.51 | 0.49 | 0.47 |
| | 1A5 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| Hg (Mg) | 1A4 | 0.61 | 0.71 | 1.04 | 1.97 | 0.52 | 0.49 | 0.48 | 0.49 | 0.49 | 0.48 | 0.47 |
| | 1A5 | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA |
| PAH (Mg) | 1A4 | 32.06 | 35.44 | 35.91 | 39.08 | 67.42 | 56.10 | 54.33 | 59.39 | 52.33 | 51.40 | 48.72 |
| | 1A5 | 0.02 | 0.01 | 0.01 | 0.02 | 0.01 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 |
| HCB (Kg) | 1A4 | 2.55 | 3.64 | 6.98 | 7.14 | 5.11 | 3.80 | 3.91 | 4.09 | 4.05 | 3.52 | 3.46 |
| | 1A5 | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA |
| PCB (Kg) | 1A4 | 14.92 | 18.33 | 25.73 | 36.26 | 22.01 | 18.99 | 18.70 | 20.20 | 19.02 | 18.48 | 17.95 |
| | 1A5 | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA | NA |

3.9.5 QA/QC and Uncertainty

Basic data used in the estimation process are reported by Ministry of Economic Development in the National Energy Balance (MSE, several years (a)) and by TERNA (National Independent System Operator), concerning the waste used to generate electricity.

The energy data used to estimate emissions have different levels of accuracy:

- the overall sum of residential and institutional/service/commercial energy consumption is quite reliable and their uncertainty is comparable with data reported in the BEN; the amount of fuels used is periodically reported by main suppliers;
- the energy consumption for agriculture and fisheries is reported in energy statistics; data are quite reliable as they have special taxation regimes and they are accounted for separately;
- the energy use for military and off roads is reported in official statistics, but models are applied to estimate the energy use at a more disaggregated level.

3.9.6 Recalculation

Some recalculations affected 1A4 category in this submission.

Energy recovery from waste reported in the commercial heating has been updated from 2018 because of the update of activity data.

Minor recalculations occur from 2000 because of the update of NCV (Net Calorific Value) for gasoil and LPG.

3.9.7 **Planned improvements**

The updating of average emission factors is planned for future submission on the basis of the surveys on wood consumption and combustion technologies planned by ISTAT on fuel consumptions as well as from the results of an emission factor measurements campaign realized in Italy (ALTROCONSUMO, 2018), and the measurements campaign on advanced stoves completed by Innovhub. An in depth analysis of emission factors resulting from this experimental studies and their comparison with the values suggested by the last version of the EMEP/EEA Guidebook (EMEP/EEA, 2019) will be carried out and emission factors will be updated as needed.

3.10 FUGITIVE EMISSIONS (NFR SUBSECTOR 1.B)

3.10.1 Overview

Fugitive emissions arise during the stages of fuel production, from extraction of fossil fuels to their final use. Emissions are mainly due to leaks or other irregular releases of gases from the production and transformation of solid fuels, the production of oil and gas, the transmission and distribution of gas and from oil refining.

In Table 3.41 the list of categories for fugitive emissions identified as key categories by pollutant for 2020, 1990 and in the trend is reported.

Table 3.41 List of key categories by pollutant in the civil sector in 2020, 1990 and trend

| | Key categories in 2020 | Key categories in 1990 | Key categories in trend |
|-----------------|------------------------|------------------------|-------------------------|
| SO _x | 1 B 2 a iv | 1 B 2 a iv | 1 B 2 a iv, 1 B 2 c |
| NMVOG | | 1 B 2 a v | 1 B 2 a v |
| NH ₃ | | | 1 B 2 d |
| Hg | 1 B 2 d | 1 B 2 d | 1 B 2 d |

3.10.2 Methodological issues

In the following methodological issues including activity data and emission factors used are reported for each category and pollutant estimated in this sub sector.

Coal mining and handling (1B1a)

NMVOG emissions from coal mining have been estimated on the basis of activity data published on the national energy balance (MSE, several years [a]) which report the amount of coal production and emission factors provided by the EMEP/EEA Guidebook (EMEP/EEA, 2016).

PM emissions from storage of solid fuels have been estimated and included in this category. Activity data is the annual consumption of solid fuels published on the national energy balance (MSE, several years [a]) and emission factor are from the US EPA Guidebook.

Solid fuel transformation (1B1b)

NMVOG emissions from coke production have been estimated on the basis of activity data published in the national energy balance (MSE, several years [a]) and country specific emission factors calculated taking in account the information provided by the relevant operators in the framework of the EPRTR registry and the ETS. NO_x, SO_x and NH₃ emissions from coke production are estimated on the basis of data communicated by the national plants in the framework of the EPRTR and are reported under 1.A.1 c category. NH₃ emissions have been estimated on the basis of data communicated by the operators for the EPRTR registry from 2002. According to the review (EEA, 2019; EEA, 2020), PAH emissions from coke production has been also estimated with emission factor from the 2019 EMEP/EEA Guidebook (EMEP/EEA, 2019) and allocated in 1.B.1b.

Oil exploration and production (1B2a i)

NMVOG emissions have been calculated according with activity data published on national energy balance (MSE, several years [a]), data by oil industry association (UP, several years), data and emission factors provided by the relevant operators.

Oil transport and storage and refining (1B2a iv)

Fugitive emissions from oil refining are estimated starting from the total crude oil losses as reported in the national energy balance (MSE, several years [a]) and occur prevalently from processes in refineries.

This category is key for SO_x in 2020, in the base year and for the trend.

Emissions in refineries have been estimated on the basis of activity data published in the national energy balance (MSE, several years [a]) or supplied by oil industry association (UP, several years) and operators especially in the framework of the European Emissions Trading Scheme (EU-ETS). Fugitive emissions in refineries are mainly due to catalytic cracking production processes, sulphur recovery plants, flaring and emissions by other production processes including transport of crude oil and oil products. These emissions are then distributed among the different processes on the basis of average emission factors agreed and verified

with the association of industrial operators, Unione Petrolifera, and yearly updated, from 2000, on the basis of data supplied by the plants in the framework of the European Emissions Trading Scheme, Large Combustion Plant Directive and EPRTTR. SO_x, NO_x and PM emissions communicated by the plants in the framework of Large Combustion Plants directive are assumed to refer to combustion and are reported under 1.A.1b while the difference with the totals, communicated to the EPRTTR, are considered as fugitive emissions and reported in 1.B.2a iv. NMVOC are communicated by the operators for the EPRTTR registry as a total and the amount to be reported as fugitive is calculated subtracting by the total emission estimates for combustion activities and reduced for the implementation of losses control technology especially for transportation and storage of liquid fuels. ETS data are used to integrate and check emission data provided. Moreover fugitive emissions are also checked with the average emission factors provided by the relevant industrial association for each relevant process, as fluid catalytic cracking, sulphur recovery plant, and storage and handling of petroleum products. NH₃ emissions from refineries have been estimated on the basis of data communicated by the operators for the EPRTTR registry and distributed between combustion and fugitive emissions according to the emission factors available in the 2016 EMEP/EEA Guidebook.

Emissions from refineries of HM and POPs are all reported in 1.A.1b on the basis of data submitted in the PRTR framework at plant level; it is not possible at the moment distinguish combustion by fugitive emissions of HM and POPs.

Distribution of oil products (1B2a v)

This category is key for NMVOC in 1990 and for the trend. The category includes fugitive emissions from oil transport which have been calculated according with the amount of transported oil (MIT, several years) and emission factors published on the IPCC guidelines (IPCC, 2006). Most of the crude oil is imported in Italy by shipment and delivered at the refineries by pipelines as offshore national production of crude oil. The category includes also NMVOC fugitive emissions for gasoline distribution, storage and at service stations. Emission factors are estimated starting from the emissions communicated in the nineties by the operators and applying the implementation of the abatement technologies as regulated by the relevant European Union legislation. Emissions from distribution of gasoline have been reduced as a result of the application of the DM 16th May 1996 (Ministerial Decree 16 May 1996), concerning the adoption of devices for the recovery of vapours and of the applications of measures on deposits of gasoline provided by the DM 21st January 2000 (Ministerial Decree 21 January 2000).

Flaring in refineries (1B2c)

This category is key for SO_x for the trend. For what concern emissions from flaring in refineries, the emission factors for SO_x, NO_x, NMVOC and CO have been provided by the relevant industrial association and are assumed constant since 1990 with the exception of SO_x that are yearly estimated on the basis of the amount of sulphur not recovered by the operators and flared. Activity data, in terms of gas flared, is from 2005 derived by the ETS data at plant level.

Fugitive emissions from geothermal production (1B2d)

According to the review process NH₃, Hg and other heavy metals from geothermal production has been estimated and included in the emission inventory in the 2018 submission with a Tier 2 methodology. Hg from this category is key category for 2020, the base year and the trend while NH₃ is key for the trend.

Emissions are monitored by the Regional relevant environmental agency, ARPAT, where all the geothermal fields are located. Activity data, geothermal energy production, are published in the national energy balance (MSE, several years [a]) while emission data resulting by the monitoring are issued by ARPAT and reported from 2000 on yearly basis (ARPAT, several years). For earlier years emission factors of 2000 have been used.

3.10.2.1 Fugitive emissions from natural gas distribution (1.B.2b)

NMVOC fugitive emissions from the transport, storage and distributions (including housing) of natural gas (both in pipelines and in the distribution network) are calculated every year on the basis of fugitive natural gas emissions and the content of NMVOC in the gas distributed; NMVOC emissions due to transport and distribution are around 99% of the total. Emissions are calculated starting from methane emissions estimates, considering the annual average percentage of NMVOC in the natural gas distributed in Italy as in Table 3.42. The methodology and references are reported in detail in the NIR (ISPRA, 2022[a]). CH₄, CO₂ and NMVOC emissions have been estimated on the basis of activity data published by industry, the national authority, and

information collected annually by the Italian gas operators. Emission estimates take into account the information on: the amount of natural gas distributed supplied by the main national company (SNAM); length of pipelines, distinct by low, medium and high pressure and by type, cast iron, grey cast iron, steel or polyethylene pipelines as supplied by the national authority for the gas distribution (AEEG); natural gas losses reported in the national energy balance; methane emissions reported by operators, in their environmental reports (EDISON, SNAM, ENEL, Italgas). NMVOC and CO₂ emissions have been calculated considering CO₂ content in the leaked natural gas. Regarding exploration and production, an average emission factor, equal to 0.04 g/m³ gas produced, has been estimated on the basis of emission data communicated by the relevant companies for some years and applied to the whole time series.

The average natural gas chemical composition has been calculated from the composition of natural gas produced and imported. Main parameters of mixed natural gas, as calorific value, molecular weight, and density have been calculated as well. Data on chemical composition and calorific value are supplied by the main national gas providers for domestic natural gas and for each country of origin.

The following table shows average data for national pipelines natural gas.

Table 3.42 Average composition for pipelines natural gas and main parameters

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| HCV (kcal/m ³) | 9,156 | 9,193 | 9,215 | 9,261 | 9,325 | 9,303 | 9,351 | 9,340 | 9,334 | 9,336 | 9,340 |
| NCV (kcal/m ³) | 8,255 | 8,290 | 8,320 | 8,354 | 8,412 | 8,391 | 8,444 | 8,433 | 8,427 | 8,428 | 8,432 |
| Molecular weight | 17.03 | 17.19 | 17.37 | 17.45 | 17.46 | 17.34 | 17.53 | 17.44 | 17.34 | 17.29 | 17.33 |
| Density (kg/Sm ³) | 0.72 | 0.73 | 0.74 | 0.74 | 0.74 | 0.73 | 0.74 | 0.74 | 0.73 | 0.73 | 0.73 |
| CH ₄ (molar %) | 94.30 | 93.36 | 92.22 | 91.93 | 92.04 | 92.72 | 91.54 | 92.08 | 92.64 | 92.92 | 92.79 |
| NMVOC (molar %) | 3.45 | 4.09 | 4.84 | 5.35 | 5.74 | 5.26 | 6.17 | 5.93 | 5.62 | 5.49 | 5.63 |
| CO ₂ (molar %) | 0.22 | 0.20 | 0.18 | 0.49 | 0.75 | 0.70 | 0.65 | 0.67 | 0.74 | 0.64 | 0.88 |
| Other no carbon gas (molar %) | 2.03 | 2.34 | 2.76 | 2.24 | 1.48 | 1.32 | 1.64 | 1.32 | 1.00 | 0.95 | 0.70 |
| CH ₄ (weight %) | 88.83 | 87.14 | 85.16 | 84.53 | 84.54 | 85.81 | 83.79 | 84.72 | 85.68 | 86.23 | 85.88 |
| NMVOC (weight %) | 7.33 | 8.62 | 10.00 | 10.73 | 11.27 | 10.34 | 12.03 | 11.51 | 10.87 | 10.64 | 10.78 |
| CO ₂ (weight %) | 0.57 | 0.51 | 0.47 | 1.23 | 1.89 | 1.78 | 1.62 | 1.70 | 1.88 | 1.63 | 2.25 |
| Other no carbon gas (weight %) | 3.27 | 3.74 | 4.37 | 3.51 | 2.30 | 2.08 | 2.55 | 2.07 | 1.56 | 1.50 | 1.09 |

More in details, emissions are estimated separately for the different phases: transmission in primary pipelines and distribution in low, medium, and high pressure network, losses in pumping stations and in reducing pressure stations (including venting and other accidental losses) with their relevant emission factors, considering also information regarding the length of the pipelines and their type.

Emissions from low pressure distribution include also the distribution of gas at industrial plants and in residential and commercial sector; data on gas distribution are only available at an aggregate level thus not allowing a separate reporting. In addition, emissions from the use of natural gas in housing are estimated and included. Emissions calculated are compared and balanced with emissions reported by the main distribution operators. Finally, the emission estimates for the different phases are summed and reported in the most appropriate category (transmission/distribution).

Table 3.43 provides the trend of natural gas distribution network length for each pipeline material and the average CH₄ emission factor.

Table 3.43 Length of low and medium pressure distribution network (km) and network emission factors for CH₄ and NMVOC

| Material | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|----------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| Steel and cast iron (km) | 102,061 | 131,271 | 141,848 | 154,886 | 198,706 | 203,116 | 204,062 | 204,890 | 205,273 | 206,855 |
| Grey cast iron (km) | 24,164 | 22,784 | 21,314 | 15,080 | 4,658 | 2,398 | 2,163 | 2,088 | 2,063 | 2,061 |
| Polyethylene (km) | 775 | 8,150 | 12,550 | 31,530 | 49,663 | 56,943 | 57,883 | 59,368 | 59,358 | 59,593 |
| Total (km) | 127,000 | 162,205 | 175,712 | 201,496 | 253,027 | 262,457 | 264,108 | 266,346 | 266,693 | 268,509 |
| CH ₄ EF (kg/km) | 1,958 | 1,417 | 1,228 | 1,000 | 703 | 540 | 522 | 516 | 469 | 383 |

| NMVOC (kg/km) | EF | | | | | | | | | | |
|------------------|----|-----|-----|-----|-----|----|----|----|----|----|----|
| | | 162 | 140 | 144 | 127 | 94 | 65 | 75 | 70 | 59 | 47 |

3.10.3 Time series and key categories

The trend of fugitive emissions from solid fuels is related to the extraction of coal and lignite that in Italy is quite low. The decrease of NMVOC fugitive emissions from oil and natural gas is due to the reduction of losses for gas transportation and distribution, because of the gradual replacement of old grey cast iron pipelines with steel and polyethylene pipelines for low and medium pressure network as reported in the previous paragraph.

3.10.4 QA/QC and Uncertainty

Different data sources are used for fugitive emissions estimates: official statistics by Economic Development Ministry (MSE, several years [a], [b]), by Transport of Infrastructure Ministry (MIT, several years); national authorities (AEEG, several years; ISTAT, several years [a]), gas operators (ENI, several years; EDISON, several years; SNAM, several years), and industrial association for oil and gas (UP, several years).

CH₄ and NMVOC emissions from transmission and distribution of natural gas are verified considering emission factors reported in literature and detailed information supplied by the main operators (ENI, several years [b]; Riva, 1997).

3.10.5 Recalculation

Recalculations occurred because of revision of national average emission factors of losses from gas distribution on the basis of new information provided by the operators from 2019 and of average parameters of the natural gas consumed in Italy from 2008.

3.10.6 Planned improvements

No further improvements are planned for this category.

4 IPPU - INDUSTRIAL PROCESSES (NFR SECTOR 2)

4.1 OVERVIEW OF THE SECTOR

Emission estimates in this category include emissions from all industrial processes and also by-products or fugitive emissions, which originate from these processes. Where emissions are released simultaneously from the production process and from combustion, as in the cement industry, they are estimated separately and included in the appropriate categories, in sector 2 and in sector 1 category 1.A.2. This sector makes important contributions to the emissions of heavy metals, PAH, dioxins and PCB.

Regarding emissions of the main pollutants, in 2020, industrial processes account for 6.7% of SO₂ emissions, 0.8% of NO_x, 0.2% of NH₃, 4.8% of NMVOC and 3.4% of CO. About particulate matter, in 2020 this sector accounts for 7.02% of PM₁₀ emissions and 5.24% of PM_{2.5}. Industrial processes make a significant contribution to the total Italian emissions of heavy metals, despite significant reductions since 1990; particularly this sector accounts for 40.56% of Pb emissions, 32.43% of Cd and 44.24% of Hg. Regarding POPs emissions, 13.8% of PAH total emissions is emitted from industrial processes as well as 27.02% of dioxins and 71.24% of PCB.

In 2020, *iron and steel* sector (2C1) is a key category at level assessment for PM₁₀, PM_{2.5}, Pb, Cd, Hg, PAH, PCDD/F and PCB; emissions from *other chemical industry* (2B10a) is a key category source for SO₂ emissions. Emissions from *cement production* (2A1) is a key category source for PM₁₀. Emissions from 2G is a key category source for Cd too. Food and beverage industry (2H2) is a key category for NMVOC emissions. In 1990 emissions from *cement production* (2A1) is a key category source for PM₁₀ and CO; *other chemical industry* (2B10a) is a key category for Hg and Cd and *iron and steel production* (2C1) is a key category for CO, PM₁₀, PM_{2.5}, Cd, Hg, PAH, PCDD/F and PCB. At trend assessment, *iron and steel* sector is key category for Pb, , Hg, PAH and PCDD/F while *other chemical industry* (2B10a) is a key category for Hg and 2G is a key category for Cd emissions.

As requested by the TERT in the last review process, information has been included in this chapter about:

- the estimates for category 2A5b Construction and Demolition;
- the estimation process for PM_{2.5} from 2A1 Cement production;
- progress of investigations about 2A5a Quarrying and mining
- explanation of estimation for PAH from category 2C1 Iron and Steel, methods and data used
- method, AD, EF for PM₁₀, PM_{2.5}, BC, PCDD/F and HCB from 2C3 Aluminium production

4.2 METHODOLOGICAL ISSUES

Methodologies used for estimating emissions from this sector are based on and comply with the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007) and EMEP/EEA guidebook (EMEP/EEA, 2019), the IPCC Guidelines (IPCC, 1997; IPCC, 2006) and the Good Practice Guidance (IPCC, 2000). Included also in this sector are by-products or fugitive emissions, which originate from industrial processes.

There are different sources relevant to estimate emissions from this sector; activity data are provided by national statistics and industrial associations but a lot of information is supplied directly from industry. In fact, as for the energy sector, references derive from data collected in the framework of the national PRTR reporting obligation, the Large Combustion Plant directives and the European Emissions Trading Scheme. Other small plants communicate their emissions which are also considered individually. These processes have improved the efficiency in collecting data and the exchange of information. Whenever data cannot be straight used for the inventory compilation, they are taken into account as verification practice. Environmental Reports published by industrial associations are also considered in the verification process.

4.2.1 Mineral products (2A)

In this sector emissions from the following processes are estimated and reported: cement production, lime production, glass production, quarrying&mining and construction and demolition.

Cement production (2A1), is considerable for PM_{2.5} emissions and accounts for 1.48% of the total national emissions in 2020. SO₂ emissions from cement production results from fuel combustion during the manufacture of cement so in the present submission they are included under the energy sector 1A2f and not included under 2A1.

During the last 15 years, in Italy, changes in cement production sector have occurred, leading to a more stable structure confirming the leadership for the production in Europe. The oldest plants closed, wet processes were abandoned in favour of dry processes so as to improve the implementation of more modern and efficient technologies. Since 2011 Italy has become the second cement producer country in the EU 28 but the reduction in clinker production seems to have stopped, since 2016 clinker production at national level has kept almost the same. In 2020, 17 companies (54 plants of which: 30 full cycle and 24 grinding plants) operate in this sector: multinational companies and small and medium size enterprises (operating at national or only at local level) are present in the country. As for the localization of the operating plants: 42.6% is in northern Italy, 14.8% is in the central regions of the country and 42.6% is in the southern regions and in the islands (Federbeton/AITEC, 2021). In Italy different types of cement are produced; as for 2020 Federbeton/AITEC, the national cement association, has characterised the national production as follows: 71% is CEM II (Portland composite cement); 15% is CEM I (Portland ordinary cement); 12% is CEM IV (pozzolanic cement) and 2% is CEM III (blast furnace cement). Clinker production has been decreasing since 2007, although from 2016 to 2019 the production values have kept very close to the amount manufactured in 2016, in 2020 clinker production shows -6.0% compared to 2019; clinker demand in cement production was about 74% in 2020 (production of clinker out of production of cement). To estimate emissions from cement production, activity data on clinker/cement production are used as provided by ISTAT (ISTAT, several years up to 2008), MSE (MSE, several years since 2009 up to 2018) and facility reports in the framework of the Emissions Trading Scheme legislation.

In this category only PM₁₀ and PM_{2.5} emissions are reported separately from combustion while all the other pollutant emissions are included in the energy sector in 1.A.2f category.

Emission factor for PM₁₀ emissions is equal to 234 g/Mg of clinker for the whole time series and is calculated on the basis of plants emission data in the nineties. Emission factor for PM_{2.5} is equal to 130 g/Mg of clinker for the whole timeseries

Regarding SO₂ emissions, according to the EMEP/EEA Guidebook, in the present submission no sulphur oxides process emissions have been reported under 2A1 because it was clarified that SO₂ emissions originate from fuel combustion and, as far as energy aspects only are concerned, they have been allocated under 1A2f.

The remaining categories of mineral products (*lime production* (2A2)) industry represent less than 1% for each pollutant.

As regards 2A3 category *Glass production*, HM, PM and BC emissions are reported under 1A2f and emission factors have been provided by the research institute of the sectoral industrial association (Stazione Sperimentale del Vetro) distinguished by the different types of glass production. On the basis of the 2017 review process (EEA, 2017 [a]), the previous notation key has been replaced by the IE notation key.

About the 2A5 category, following the suggestions of the NECD review, more information has been added but different activities have to be dealt separately.

As regards 2A5a *Quarrying and mining of minerals other than coal*, there is no evidence of active mines of the main minerals as those indicated in the Guidebook (bauxite, copper, manganese and zinc). All these mines closed before 1990 for economic reasons. At the same time there is no available data to apply a Tier 1 on other mineral mines. The USGS Mineral yearbook provides info for Italy only for Feldspar, Gypsum, Pumice and Sand and gravel extraction. All the data are estimated and we are verifying the activity level with industry and local competent authorities. Moreover, it should be verified if the EFs available in the Guidebook are applicable to these national extractive activities because of the abatement technologies and the kind of mineral. A first rough estimation of emissions has been done using the spreadsheet available in the 2019 EMEP/EEA

Guidebook (2.A.5.a Quarrying and mining calculation model 2019 - Spreadsheet.xlsx), introducing production data for 2016 (last year available) as reported in the USGS database for crushed rocks, sand and gravels without modifying the other default values introduced (e.g. the number of plants) because at the moment these are unknown. Recycled aggregates have not been included in the calculation since crushed asphalt is recycled in specific plants and these emissions are already reported under the relevant category while for crushed concrete there is no specific information available in the USGS statistics and no other statistics are available that could be used alternatively. It should be noted that the concrete inert material in Italy goes prevalently to landfills. PM_{2.5} emission estimates results in 1.3 kt where most part of emissions are due to the internal transportation (1.1 kt), mostly due to resuspension of particles which do not seem to be included in the inventory (for road transport it is not). Further investigations are needed to clarify which mineral products have to be considered for the emissions estimation process.

As for the category 2A5b *Construction and Demolition*, the National Institute of Statistics provides information about residential and non-residential buildings, for both categories information about volumes (for the year from 1990-2000) or surfaces (usable living areas) could be retrieved to work out a timeseries of activity data and to develop estimates for TSP, PM₁₀ and PM_{2.5}. Emission factors for TSP (162 kg/1000 m²), PM₁₀ (81 kg/1000 m²) and PM_{2.5} (8 kg/1000 m²) are those included in the 2019 EMEP/EEA Guidebook.

For the category 2A5c *Storage, Handling and Transport of Mineral Products*, PM_{2.5} emissions have been estimated and reported in the sectoral categories 2A1 Cement Production and 2A2 Lime Production. The emissions from storage, handling and transport for other minerals than the aforementioned ones might not be included in the inventory because this potential under-estimate is likely to be below the threshold of significance.

4.2.2 Chemical industry (2B)

Emissions of this sector derive from organic and inorganic chemicals processes and are usually not significant except for SO_x emissions from the production of sulphuric acid and Hg emissions from chlorine production. Emission factors derive from data collected in the framework of the national EPER/E-PRTR register as well as from EMEP/EEA and EPA Guidebook.

As already mentioned, *other chemical industry* (2B10a) was key category for Hg emissions in 1990 and for SO_x emissions in 2020. Hg emissions are released from chlorine production facility with mercury cells process (EUROCHLOR, 1998). Total chlorine production in Italy amounted, in 1990, to 1,042,921 tonnes and reduced in 2020 to 240,740 tonnes. Activity production data are supplied by the National Institute of Statistics (ISTAT) and published in the official national statistics and since 2002 data have also been collected at facility level in the national EPER/E-PRTR register. To estimate emissions from 1990 to 2001, the average emission factor supplied by EUROCHLOR for western Europe chlor-alkali production plants (EUROCHLOR, 2001) has been used, while since 2002 emission data have been supplied directly by the production facilities in the framework of the national EPER/E-PRTR. The average emission factor decreased from 1.11 g Hg/t in 2002 to zero in 2018. The reduction observed in emissions for the last years is a consequence of both the conversion of production plants from the mercury cells process to the membrane technology and also the suspension of production at the existing facilities. In 2007 seven facilities carried out the chlor-alkali production: one facility had the membrane process in place, one facility was replacing mercury cells with membrane process while in the other five facilities the production was still based on the mercury cell process (Legambiente, 2007). In 2015 five facilities carried out chlor-alkali production: in four of them the membrane process was in place while one facility still operated the mercury cell process. In 2018 the four chlor-alkali facilities have the membrane process in place while the one with mercury cells was obliged to stop the production with this process and it is still in operation although the manufacturing process has been relying on the purchase of the intermediate products since then.

SO_x emissions reported in *other chemical industry* (2B10a) include emissions from sulphuric acid production and account for 4.97% of total SO_x emissions in 2020. Activity production data are supplied by the National Institute of Statistics (ISTAT) and published in the official national statistics and since 2004 data have also been collected at facility level in the national EPER/E-PRTR register. Emission factors from 1990 to 1994 and from 2002 are derived from emission data supplied directly by the production facilities in the framework of the CORINAIR inventory project and of the national EPER/E-PRTR, respectively.

On the basis of the 2017 review process NO_x, SO_x, CO, PM and BC emissions from 2B7 *Soda ash production* have been estimated. In Italy there is only one plant producing soda ash and it is in the framework of the EPRTR reporting. In particular, as regards PM emissions, the operator has never reported PM₁₀ emissions which implies that emissions are under the reporting threshold (50 t/year). As reported in the Guidebook measurements made in some plants indicate that more than 75% of the dust emitted is made of particle size > 10 µm and that the contribution of PM₁₀ is relatively low. Moreover the operator in its annual environmental report estimates TSP emissions (around 200 t/y) reporting explicitly that no PM₁₀ emissions occur. The achieved estimates, using the EMEP/EEA EFs, produced figures of around 20 Mg (PM₁₀) consistently with respect to the E-PRTR thresholds.

4.2.3 Metal production (2C)

The main activities in this sector are those regarding the *iron and steel* production. The main processes involved in iron and steel production are those related to sinter and blast furnace plants, to basic oxygen and electric furnaces and to rolling mills.

The sintering process is a pre-treatment step in the production of iron where fine particles of metal ores are agglomerated. Agglomeration of the fine particles is necessary to increase the passageway for the gases during the blast furnace process and to improve physical features of the blast furnace burden. Coke and a mixture of sinter, lump ore and fluxes are introduced into the blast furnace. In the furnace the iron ore is increasingly reduced and liquid iron and slag are collected at the bottom of the furnace, from where they are tapped. The combustion of coke provides both the carbon monoxide (CO) needed for the reduction of iron oxide into iron and the additional heat needed to melt the iron and impurities. The resulting material, pig iron (and also scrap), is transformed into steel in subsequent furnaces which may be a basic oxygen furnace (BOF) or electric arc furnace (EAF). Oxygen steelmaking allows the oxidation of undesirable impurities contained in the metallic feedstock by blowing pure oxygen. The main elements thus converted into oxides are carbon, silicon, manganese, phosphorus and sulphur.

In an electric arc furnace steel is produced from polluted scrap. The scrap is mainly produced by cars shredding and does not have a constant quality, even if, thanks to the selection procedures, the scrap quality becomes better year by year. The iron and steel cycle is closed by rolling mills with production of long products, flat products and pipes.

In 1990 there were four integrated iron and steel plants in Italy. In 2020, there are only two of the above mentioned plants remaining, one of which lacks BOF; oxygen steel production represents about 16.5% of the total production and the arc furnace steel the remaining 83.5% (FEDERACCIAI, several years). Currently, long products represent about 48% of steel production in Italy, flat products about 41%, and pipe the remaining 12%. Most of the flat production derives from only one integrated iron and steel plant while, in steel plants equipped with electric ovens almost all located in the northern regions, long products are predominantly produced (e.g carbon steel, stainless steels) and seamless pipes (only one plant) (FEDERACCIAI, several years).

Basic information for *Iron and steel production* derives from different sources in the period 1990-2020. Activity data are supplied by official statistics published in the national statistics yearbook (ISTAT, several years) and by the sectoral industrial association (FEDERACCIAI, several years).

For the integrated plants, emission and production data have been communicated by the two largest plants for the years 1990-1995 in the framework of the CORINAIR emission inventory, distinguished by sinter, blast furnace and BOF, and by combustion and process emissions. From 2000 production data have been supplied by all the plants in the framework of the ETS scheme, for the years 2000-2004 disaggregated for sinter, blast furnace and BOF plants, from 2005 specifying carbonates and fuels consumption. For 2002-2015 data have also been supplied by all the four integrated iron and steel plants in the framework of the EPER/E-PRTR registry but not distinguished between combustion and process. National experts have also been involved in the process of elaboration of the “monitoring and control plan” for the largest integrated plant in Italy in the framework of the IPPC permit. Qualitative information and documentation available on the plants allowed reconstructing their history including closures or modifications of part of the plants; additional qualitative information regarding the plants, collected and checked for other environmental issues or directly asked to the

plant, permitted to individuate the main driving of the emission trends for pig iron and steel productions. Emissions from lime production in steel making industries are reported in 1A2f Manufacturing Industries and Construction category.

In 2020, *iron and steel sector* (2C1) is key category for PM10, PM2.5, Pb, Cd, Hg, PAH, PCDD/F and PCB. In Table 4.1 relevant emission factors are reported.

Table 4.1 Emission factors for iron and steel for the year 2020

| | | PM10 [g/Mg] | PM2.5 [g/Mg] | Cd [mg/Mg] | Hg [mg/Mg] | Pb [mg/Mg] | PCB [mg/Mg] | PAH [mg/Mg] | PCDD/F [µg T-eq/Mg] |
|----------------------------------|--------------|----------------|-----------------|---------------|---------------|---------------|----------------|----------------|------------------------|
| Blast furnace charging | <i>Areal</i> | 60 | 37.5 | | | | | | |
| | <i>Point</i> | 2.8 | 1.8 | | | | | | |
| Pig iron tapping | <i>Areal</i> | 41.4 | 25.9 | 0.3 | 0.3 | 15 | | 950 | |
| | <i>Point</i> | 2.0 | 1.2 | 0.3 | 0.3 | | | | |
| Basic oxygen furnace | <i>Areal</i> | 62 | 54.3 | 25 | 3 | 850 | 3.6 | | |
| | <i>Point</i> | 10.7 | 9.4 | 2.3 | 1.7 | 49 | 3.6 | | |
| Electric arc furnace | | 124 | 108.5 | 50 | 150 | 3450 | 3.6 | 1.9 | 4.45 |
| Rolling mills | <i>Areal</i> | 59 | 45.9 | | | | | 125 | |
| | <i>Point</i> | 28.2 | 21.9 | | | | | 125 | |
| Sinter plant (except combustion) | <i>Areal</i> | 16 | 12.8 | | | | | | |
| | <i>Point</i> | 3.5 | 2.8 | | | | | | |

PM10 emission factors for integrated plants derive from personal communication of the largest Italian producer of pig iron and steel (ILVA, 1997) while PM10 emission factor for electric arc furnace derives from a sectoral study (APAT, 2003). The Emission factors manual PARCOM-ATMOS (TNO, 1992), the EMEP/Corinair Guidebook (EMEP/CORINAIR, 2006) and the IPPC BRef Report (IPPC, 2001) provide emission factors for heavy metals while a sectoral study (APAT, 2003) provides Cd emission factors for electric arc furnace. Recently, as reported above, work has begun to revise the EFs from the largest integrated plant (point sources) thanks to the analysis of the monitoring and control plans linked to the data collected in the framework of the IPPC permits. Emission factors derived from this survey on the IPPC permits have been applied since 2016 and relate both to combustion and process categories. Further information about combustion emissions is given in the combustion chapter. As regards the category 2C1, at the moment, EFs for PM emissions from blast furnace charging, pig iron tapping, basic oxygen furnace and sinter plant have been updated in the current submission. For BOF emissions Cd, Pb and Hg EFs have been updated too.

Regarding PAH emissions, for blast furnaces, results from measurements tell us that emission factor used for pig iron tapping covers also blast furnace charging while emissions from BOF are negligible.

To complete the category analysis, for 1A1c category and, in particular, for coke production according to the review (EEA, 2019) PAH emission factor has been disaggregated into those deriving from the combustion process and the fugitive ones and estimated with the emission factors in the Guidebook (EMEP/EEA, 2019).

Regarding POPs emissions, emission factors usually originate from EMEP/CORINAIR (EMEP/CORINAIR, 2007, EMEP/CORINAIR, 2006) except those relating to PAH and PCDD/PCDF from electric arc furnace that derive from direct measurements in some Italian production plants (ENEA-AIB-MATT, 2002). Dioxin emissions for sinter plant, and other sources within steelworks manufacturing oxygen steel occur during the combustion process and they are measured at the stack; emissions are therefore reported in the energy sector in 1.A.2a category. In 2020 the average emission factor is equal to 0.30 micrograms TEQ per Mg of sinter produced. EF is calculated yearly on the basis of measurements done in the two existing sinter plant in Italy. As regards HCB emissions, Italy reports HCB emissions from sintering production calculated with the 2006 Guidebook (“Sources of HCB emissions.pdf” does not distinguish between combustion and process) EF=0,032mg/Mg in 1A2a because in this case HCB emissions are clearly linked to the combustion activities. The 2016 Guidebook provides reference to the 2006 version.

As for other iron and steel activities, a series of technical meetings with the most important Italian manufacturers was held in the framework of the national PRTR in order to clarify methodologies for estimating POPs emissions. In the last years, a strict cooperation with some local environmental agencies allowed the acquisition of new data, the assessment of these data is still ongoing and improvements in emission estimates are expected for the next years. Thanks to the last review process in the framework of the NEC Directive (EEA, 2019) fugitive PAH emissions from coke oven (door leakage and extinction) have been estimated on the basis of 2019 EMEP/EEA Guidebook emission factors. As a consequence of the review process these emissions have been reported in 1B1b adopting the right allocation.

Emission factors used in 1990 estimates generally derive from Guidebook EMEP/CORINAIR.

The remaining categories of metal production industry represent about 0% for each pollutant because of the shutdown of several plants, in particular those linked to the non ferrous production.

As discussed during the recent review process (EEA, 2018; EEA, 2021), indeed, no plants for aluminium production by electrolysis have been working in Italy since 2012 and pollutants time series are reported, obviously, from 1990 to 2012, consequently no primary aluminium production has been occurring since 2012. Based on the Guidebook, emissions from primary aluminium production have been generally reported under 2C3. According to the Guidebook, most of the pollutants and EFs available for Tier 1 and Tier 2 for 2C3 category refer to primary aluminium production, since this activity stopped in Italy in 2012, notation key “IE” has been reported according to the last review process. Secondary aluminium production still occurs in Italy, activity data are provided annually by the statistical office of Assomet, the trade association dealing with non-ferrous metals. In general, emissions from secondary aluminium production are allocated under the energy sector (1A2b). Process emissions from secondary aluminium, including PM and POPs, are reported under the relevant category in the energy sector. Emission factor of Dioxin and PAH are country specific, while PM10 is from USEPA and the speciation of PM2.5 and BC refer to the value reported in the Guidebook for secondary aluminium. Moreover, HCB emissions from secondary aluminium production are not reported and are expected to be null because these emissions derive from the degassing of aluminium when hexachloroethane is used, but this compound is banned in Italy from '90s. Further information about emissions from secondary aluminium production are included in the paragraph devoted to category 1A2b included in Chapter 3, paragraph 3.3.1.2 and 3.3.2.2.

As for the production of lead, zinc and copper (2C5, 2C6 and 2C7a categories), at the moment SO_x, HM and PM emissions are reported in the energy sector because up to now there was no information to distinguish between energy and process emissions and, above all, these processes are considered combustion processes with contact, consequently, emissions are dependent on the combustion process. In the last year, thanks to the ETS data, it has been possible to separate CO₂ emissions in these two components and Italy is investigating the possibility of extension to other pollutants for the next submissions. In particular, in Italy no production of primary copper has ever occurred while, as regards lead and zinc, there is a sole integrated plant for the primary productions, and this makes it difficult to ensure a good breakdown. Consequently, the issue related to the allocation of emissions is not only about combustion and process but also about the different productions of different metals in the same factory. To resolve this issue, an in-depth investigation has been started with the aim to better specify the technology used on the basis of E-PRTR and IPPC permits. The first result of this investigation has been the update of certain EFs since 2014 (ISPRA, 2021). Anyway, for Pb, Cd and PCB the notation key IE has been added in the NFR because of the relevant emissions are reported in the energy sector.

Moreover in response to the review process Italy explained that the Hg emission factor for copper production in the EMEP/EEA Guidebook 2019 is not applicable because it refers to primary copper production while in Italy copper production between 1990 and 1998 was derived only from secondary technologies.

4.2.4 Other production (2G – 2H – 2I – 2L)

2G sector includes NMVOC emissions due to the *use of lubricants* as well as all potential emissions from the *use of tobacco and fireworks*. In 2H sector, non-energy emissions from *pulp and paper* as well as *food and drink* production, especially wine and bread, are reported. TSP emissions from wood processing are included and reported in 2I, Lead emissions from *batteries manufacturing* can be found in 2L sector.

Emissions from these categories are usually negligible except for NMVOC emissions from *food and drink* (2H2) accounting for 3.2% of the national total in 2020 and Cd from fireworks (2G), both are key categories in 2020. Emissions from this category refer to the processes in the production of bread, wine, beer and spirits. Activity data are derived from official statistics supplied by the National Institute of Statistics (ISTAT) and relevant industrial associations. Time series of bread production is reconstructed for the '90 years on the basis of family surveys from the national Institute of statistics (ISTAT) while from 1998 data are those reported in the PRODCOM statistics officially communicated by ISTAT to EUROSTAT. PRODCOM data collection has improved along the years producing more reliable figures. In the '00 years, bread production has changed from fresh artisanal production to a more industrial oriented production, without any impact on the total. For wine, beer and spirits the statistical information on activity data is much more reliable and their trends are driven by the seasonal variation (for wine) or market demand (for beer) while for spirits it is mostly driven by a change in the personal habits and relative consumptions. Emission factors are those reported in the EMEP/CORINAIR guidebook and, in lack of national information, they are assumed constant for the whole time series (CORINAIR, 1994; EMEP/CORINAIR, 2006).

Pulp and paper industry (2H1) referred to the acid sulphite and neutral sulphite semi-chemical processes up to 2007 and only to the neutral sulphite semi-chemical process for 2008 and 2009, while the kraft process was not present in Italy. Emissions of NO_x, NMVOC, SO_x and PM were estimated for those years on the basis of activity data provided by the two Italian production plants. In 2008 the bleached sulphite pulp production stopped and in 2009 the neutral sulphite semi-chemical pulp process plant also closed. So, for the IPPU inventory purposes, there was no production of pulp and paper after 2009 and consequently no emissions have been estimated. Acid sulphite process emissions are calculated for SO_x, NMVOC and NO_x on the basis of EFs available in the Best Available Techniques Reference Documents report (BRef report), for PM₁₀ on the basis of EF in the USEPA Guidebook (54% PST) while for PM_{2.5} and BC emission profiles reported in the EMEP/EEA 2016 Guidebook (Table 3.3) have been used. For neutral sulphite semi-chemical process the emission factors used through the time period referred for SO_x, NMVOC and NO_x to CORINAIR 1992, EMEP/CORINAIR Guidebook, and for NO_x, from 1996, data were communicated by the operator of the plant.

NMVOC emissions include emissions from chipboard production where activity data are those in the FAOSTAT database for particle board and the emission factor 500 g/Mg product is from “Corinair 1992 Default Emission Factors Handbook”.

Regarding 2G category (*other product use*) all potential emissions have been estimated both for the use of tobacco and fireworks; NMVOC, SO_x, NO_x, CO, NH₃, Cd, Pb, PM₁₀, PM_{2.5}, PCDD, Benzo(a)pyrene and PAH are estimated. For activity data, as regards fireworks, Eurostat data on import, export and production of fireworks have been used, while data on consumption of tobacco were collected from the Ministry of Health, observatory of tobacco smoking. Emission factors are those reported in the EMEP/EEA 2019 Guidebook (EMEP/EEA, 2019).

In 2I category TSP emissions from wood processing are reported. Considering that in Italy wood furniture production start from wood panels and sawnwood, emissions are estimated on the basis of statistics from the FAOSTAT database statistics for that kind of wood production and emission factor in the EMEP/EEA 2019 Guidebook equal to 1 kg/t of wood product.

In 2L category lead emissions from batteries manufacturing are reported. Activity data are provided by the non-ferrous metal industrial association (ASSOMET) and refer to the amount of lead used for the batteries production; the emission factor has been provided by the relevant industrial association (ANIE) calculated on the basis of average lead concentration to the chimney, equal to 0.2 mg/Nmc, the average flow (equal to 15 Nmc/h/tonnes Pb) and the annual number of hours.

4.3 TIME SERIES AND KEY CATEGORIES

The following sections present an outline of the main key categories, and relevant trends, in the industrial process sector. Table 4.2 reports the key categories identified in the sector.

Table 4.2 Key categories in the industrial processes sector in 2020

| | 2A1 | 2A2 | 2A5b | 2B1 | 2B2 | 2B3 | 2B6 | 2B7 | 2B10a | 2C1 | 2C2 | 2C3 | 2C7c | 2G | 2H1 | 2H2 | 2L |
|-----------------------|------|------|------|------|------|------|------|------|-------|-------|-----|-----|------|------|------|------|------|
| | | | | | | | | | | % | | | | | | | |
| SO_x | | | | 0.01 | | | 0.14 | 0.19 | 4.97 | 1.42 | | | | 0.01 | | | |
| NO_x | | | | 0.04 | 0.05 | 0.00 | 0.01 | 0.03 | 0.29 | 0.38 | | | | 0.02 | | | |
| NH₃ | | | | 0.00 | 0.00 | | | 0.07 | 0.02 | | | | | 0.08 | | | |
| NMVOC | | | | 0.01 | | | | | 0.20 | 0.32 | | | | 1.11 | 0.15 | 3.23 | |
| CO | | | | 0.00 | | | | 0.58 | 0.58 | 2.03 | | | | 0.21 | | | |
| PM10 | 1.89 | 0.69 | 0.98 | | | | 0.00 | 0.01 | 0.20 | 2.07 | | | | 1.17 | | 0.01 | |
| PM2.5 | 1.31 | 0.17 | 0.12 | | | | 0.00 | 0.02 | 0.10 | 2.16 | | | | 1.36 | | | |
| BC | 0.33 | 0.01 | | | | | 0.00 | 0.00 | 0.02 | 0.07 | | | | | | | |
| Pb | | | | | | | | | | 37.79 | | | | 1.38 | | | 1.39 |
| Cd | | | | | | | | | 1.20 | 22.42 | | | | 8.80 | | | |
| Hg | | | | | | | | | | 44.24 | | | | | | | |
| PAH | | | | | | | | | | 11.81 | | | | 0.01 | | | |
| Dioxin | | | | | | | | | | 27.01 | | | | 0.00 | | | |
| HCB | | | | | | | | | | | | | | | | | |
| PCB | | | | | | | | | | 71.24 | | | | | | | |

Note: key categories are shaded in blue

There is a general reduction of emissions in the period 1990 - 2020 for most of the pollutants due to the implementation of different directives at European and national level. A strong decrease is observed especially in the chemical industry due to the introduction of relevant technological improvements.

4.3.1 Mineral products (2A)

As mentioned above, TSP, PM10 and PM2.5 emission factors for cement production are set constant from 1990 to 2020 while in the present submission SO₂ emissions are set as not occurring along the whole timeseries as they originate from fuel combustion and consequently they are allocated under 1A2f. The trends of TSP, PM10 and PM2.5 emissions follow that of the activity data.

In Table 4.3, activity data, TSP, PM10 and PM2.5 emissions from cement production are reported.

Table 4.3 Activity data and PM10 emissions from cement production, 1990 – 2020 (Gg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|-------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Cement production [Gg] | 42,414 | 35,432 | 41,119 | 47,291 | 34,283 | 20,825 | 19,305 | 19,300 | 19,241 | 18,060 |
| Clinker production [Gg] | 29,785 | 28,778 | 29,816 | 33,122 | 25,239 | 15,527 | 14,822 | 14,820 | 15,119 | 13,389 |
| TSP emissions (Gg) | 7.74 | 7.48 | 7.75 | 8.61 | 6.56 | 4.04 | 3.85 | 3.85 | 3.93 | 3.48 |
| PM10 emissions [Gg] | 6.97 | 6.73 | 6.98 | 7.75 | 5.91 | 3.63 | 3.47 | 3.47 | 3.54 | 3.13 |
| PM2.5 emissions [Gg] | 3.9 | 3.7 | 3.9 | 4.3 | 3.3 | 2.0 | 1.9 | 1.9 | 2.0 | 1.7 |

In Table 4.4, activity data, TSP, PM10 and PM2.5 emissions from construction and demolition are reported.

Table 4.4 AD, estimates and significance on the national totals for TSP, PM10 and PM2.5 from 2A5b (1990-2020)

| Year | AD (m ²) | Emissions from 2A5b | | | National totals (sub. 2022) | | | Significance on the national totals | | |
|------|-------------------------|---------------------|-----------|----------|-----------------------------|-----------|-----------|-------------------------------------|------|-------|
| | | TSP | PM10 | PM2.5 | TSP | PM10 | PM2.5 | TSP | PM10 | PM2.5 |
| | | | Mg | | | Mg | | | % | |
| 1990 | 80,813.47 | 13,091.78 | 6,562.053 | 656.2053 | 364,742.5 | 302,491 | 230,416.4 | 3.6 | 2.2 | 0.3 |
| 1991 | 79,700.83 | 12,911.53 | 6,471.707 | 647.1707 | 373,446.7 | 309,790.8 | 238,838.2 | 3.5 | 2.1 | 0.3 |
| 1992 | 74,916.57 | 12,136.48 | 6,083.226 | 608.3226 | 362,379.1 | 300,683.5 | 231,239.5 | 3.3 | 2.0 | 0.3 |
| 1993 | 63,262.72 | 10,248.56 | 5,136.933 | 513.6933 | 360,830.4 | 300,346.1 | 232,630.4 | 2.8 | 1.7 | 0.2 |
| 1994 | 64,975.12 | 10,525.97 | 5,275.98 | 527.598 | 356,101.4 | 295,676.2 | 228,073.8 | 3.0 | 1.8 | 0.2 |
| 1995 | 76,047.5 | 12,319.69 | 6,175.057 | 617.5057 | 360,521 | 297,354.4 | 228,248.7 | 3.4 | 2.1 | 0.3 |
| 1996 | 68,788.19 | 11,143.69 | 5,585.601 | 558.5601 | 343,349.3 | 284,045.7 | 217,760.9 | 3.2 | 2.0 | 0.3 |
| 1997 | 58,331.85 | 9,449.759 | 4,736.546 | 473.6546 | 337,279 | 277,965.3 | 215,097.4 | 2.8 | 1.7 | 0.2 |
| 1998 | 59,221.3 | 9,593.851 | 4,808.77 | 480.877 | 337,634.8 | 277,339.7 | 214,523.3 | 2.8 | 1.7 | 0.2 |
| 1999 | 66,010.57 | 10,693.71 | 5,360.059 | 536.0059 | 334,550.1 | 273,519.4 | 211,472.2 | 3.2 | 2.0 | 0.3 |
| 2000 | 72,362.57 | 11,722.74 | 5,875.84 | 587.584 | 316,714.3 | 257,077.6 | 197,428.8 | 3.7 | 2.3 | 0.3 |
| 2001 | 66,898.31 | 10,837.53 | 5,432.143 | 543.2143 | 308,218.2 | 249,156.9 | 189,119.8 | 3.5 | 2.2 | 0.3 |
| 2002 | 78,205.57 | 12,669.3 | 6,350.292 | 635.0292 | 272,833.5 | 218,099.9 | 159,140.5 | 4.6 | 2.9 | 0.4 |
| 2003 | 71,556.18 | 11,592.1 | 5,810.362 | 581.0362 | 292,647 | 234,896.6 | 177,633.6 | 4.0 | 2.5 | 0.3 |
| 2004 | 77,529.29 | 12,559.74 | 6,295.378 | 629.5378 | 267,295.3 | 212,961.5 | 154,831 | 4.7 | 3.0 | 0.4 |
| 2005 | 73,925.27 | 11,975.89 | 6,002.732 | 600.2732 | 288,908.8 | 231,522 | 176,423.7 | 4.1 | 2.6 | 0.3 |
| 2006 | 69,903.27 | 11,324.33 | 5,676.146 | 567.6146 | 294,549.8 | 235,661.5 | 182,224.5 | 3.8 | 2.4 | 0.3 |
| 2007 | 68,171.56 | 11,043.79 | 5,535.53 | 553.553 | 321,534.1 | 258,858.1 | 206,979.5 | 3.4 | 2.1 | 0.3 |
| 2008 | 57,921.52 | 9,383.287 | 4,703.228 | 470.3228 | 330,902.2 | 268,375.6 | 219,567.9 | 2.8 | 1.8 | 0.2 |

| Year | Emissions from 2A5b | | | | National totals (sub. 2022) | | | Significance on the national totals | | |
|------|---------------------|-----------|-----------|----------|-----------------------------|-----------|-----------|-------------------------------------|------|-------|
| | AD | TSP | PM10 | PM2.5 | TSP | PM10 | PM2.5 | TSP | PM10 | PM2.5 |
| | (m ²) | Mg | | | Mg | | | % | | |
| 2009 | 43,831.92 | 7,100.771 | 3,559.152 | 355.9152 | 301,014.3 | 245,725.1 | 203,368.4 | 2.4 | 1.4 | 0.2 |
| 2010 | 38,285.37 | 6,202.229 | 3,108.772 | 310.8772 | 294,610.6 | 239,844.1 | 198,587.7 | 2.1 | 1.3 | 0.2 |
| 2011 | 35,681.11 | 5,780.34 | 2,897.306 | 289.7306 | 238,360.7 | 191,643.3 | 151,802.3 | 2.4 | 1.5 | 0.2 |
| 2012 | 28,012.68 | 4,538.054 | 2,274.629 | 227.4629 | 266,360.4 | 216,026.6 | 178,660.3 | 1.7 | 1.1 | 0.1 |
| 2013 | 19,930.55 | 3,228.75 | 1,618.361 | 161.8361 | 256,381.9 | 207,974.9 | 172,221.3 | 1.3 | 0.8 | 0.1 |
| 2014 | 18,248.71 | 2,956.291 | 1,481.795 | 148.1795 | 235,450.3 | 189,963.1 | 154,517.7 | 1.3 | 0.8 | 0.1 |
| 2015 | 18,273.12 | 2,960.245 | 1,483.777 | 148.3777 | 241,565 | 195,004.5 | 159,920.5 | 1.2 | 0.8 | 0.1 |
| 2016 | 19,986.58 | 3,237.826 | 1,622.91 | 162.291 | 235,598.4 | 189,505.1 | 154,939.1 | 1.4 | 0.9 | 0.1 |
| 2017 | 23,245.74 | 3,765.81 | 1,887.554 | 188.7554 | 244,330.5 | 196,480.1 | 162,151 | 1.5 | 1.0 | 0.1 |
| 2018 | 25,517.41 | 4,133.821 | 2,072.014 | 207.2014 | 223,792 | 178,374.3 | 143,674.6 | 1.8 | 1.2 | 0.1 |
| 2019 | 25,456.48 | 4,123.949 | 2,067.066 | 206.7066 | 220,648.8 | 176,050.5 | 137,844.6 | 1.9 | 1.2 | 0.1 |
| 2020 | 19,985.4 | 3,237.635 | 1,622.815 | 162.2815 | 206,580.5 | 165,742.3 | 133,179.6 | 1.6 | 1.0 | 0.1 |

4.3.2 Chemical industry (2B)

Other chemical industry (2B10a) was a key category for Hg emissions in 1990 and for SO_x in 2020 and for Hg at trend assessment. Hg emissions refer to chlorine production with mercury cells process; in Table 4.5, activity data and Hg emissions from chlorine production are reported. As reported in paragraph 4.1, to estimate emissions from 1990 to 2001, the average emission factor supplied by EUROCHLOR for western Europe chlor-alkali production plants has been used, while from 2002 emission data have been supplied directly from the production plants in the framework of the national EPER/E-PRTR reporting obligation. The average emission factor decreased from 1.11 g Hg/t in 2002 to zero in 2018. The reduction observed in Hg emissions for the last years is a consequence of the conversion of production plants from the mercury cells process to the membrane technology but it depends also on suspensions of production processes at some facilities.

Table 4.5 Activity data and Hg emissions from chlorine production, 1990 – 2020

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--------------------|-------|-------|-------|-------|--------|--------|--------|--------|--------|--------|
| Activity data [Gg] | 1,043 | 869.4 | 786.0 | 535.2 | 257.75 | 217.51 | 237.31 | 280.58 | 249.07 | 240.74 |
| Hg emissions [Mg] | 2,82 | 1,65 | 0,94 | 0,48 | 0,12 | 0,04 | 0,03 | - | - | - |

SO_x emissions are prevalently from carbon black production. Sulphuric acid production, titanium oxide, other sulphate and phthalic anhydride productions are other sources reported in 2B10a and emitting SO_x. Activity data and emission factors for these sources are collected at plant level on annual basis.

4.3.3 Metal production (2C)

Emission trend of HMs, PCB and PCDD/PCDF is driven mainly by the electric arc furnaces iron and steel production which increased from 15.1 Mt in 1990 to 19.6 Mt in 2008; in 2009, because of the economic crisis, steel production from electric arc has decreased substantially and since 2010 the production has increased again up to 19.0 Mt in 2019 and it decreased again in 2020 to 17.07 Mt.

In Table 4.6, activity data and HM, PCB and PCDD/PCDF emissions from electric arc furnace (EAF) and from the whole sector 2C1 are reported, but dioxins emissions from sinter plant are reported in the energy sector in 1.A.2f category. In 2020 average emission factor is equal to 0.30 micrograms TEQ per Mg of sinter produced. EF is calculated yearly on the basis of measurements done in the two existing sinter plant in Italy.

Table 4.6 Activity data and HMs, PCB and PCDD/PCDF emissions from electric arc furnace, 1990 – 2020

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|----------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Steel production EAF [kt] | 15,102 | 16,107 | 15,879 | 17,661 | 17,115 | 17,255 | 19,336 | 19,983 | 18,980 | 17,007 |
| Cd emissions EAF [Mg] | 1.1 | 1.1 | 0.8 | 0.9 | 0.9 | 0.9 | 1.0 | 1.0 | 0.9 | 0,9 |
| Cd emissions 2C1 [Mg] | 1.3 | 1.4 | 1.1 | 1.2 | 1.1 | 1.0 | 1.0 | 1.1 | 1.1 | 0,9 |
| Hg emissions EAF [Mg] | 2.3 | 2.4 | 2.4 | 2.6 | 2.6 | 2.6 | 2.9 | 3.0 | 2.8 | 2,6 |
| Hg emissions 2C1 [Mg] | 2.3 | 2.5 | 2.4 | 2.7 | 2.6 | 2.6 | 2.9 | 3.0 | 2.9 | 2,6 |
| Pb emissions EAF [Mg] | 52.1 | 55.6 | 54.8 | 60.9 | 59.0 | 59.5 | 66.7 | 68.9 | 65.5 | 58,7 |
| Pb emissions 2C1 [Mg] | 61.1 | 65.7 | 64.1 | 71.0 | 66.5 | 63.7 | 67.0 | 69.2 | 65.8 | 58,9 |
| PCB emissions EAF [kg] | 54.4 | 58.0 | 57.2 | 63.6 | 61.6 | 62.1 | 69.6 | 71.9 | 68.3 | 61,2 |
| PCB emissions 2C1 [kg] | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 | 0,1 |
| PCDD/F emissions EAF [g T-eq] | 67.2 | 71.7 | 70.7 | 78.6 | 76.2 | 76.8 | 86.0 | 88.9 | 84.5 | 75.7 |
| PCDD/F emissions 2C1 [g T-eq] | 67.2 | 71.7 | 70.7 | 78.6 | 76.2 | 76.8 | 86.0 | 88.9 | 84.5 | 75.7 |

For Pb and Hg, the same EFs have been used for the whole time series (derived by the EMEP/CORINAIR Guidebook), while for Cd a national emission factor, equal to 50 mg/t, was available thanks to a sectoral study (APAT, 2003) and refers to the years after 1997.

This study shows range < 1-54 mg/t and the value set to 50 mg/t was chosen for conservative reason being more consistent with the old one; this value should include technology progresses occurred in the iron and steel production activities in those years. Lacking information for the years backwards, the default CORINAIR EF was used.

For PCB and PCDD/Fs, emission factors are constant from 1990 to 2020 and emission trends are ruled by the activity data.

For SO₂ and PM emissions from lead, zinc and copper production they are included and reported in the energy relevant sector. In Italy there is a sole integrated plant for the primary production of zinc and lead and this makes it difficult to ensure a good breakdown between the energy and the process sectors and the activities. During the latest year more information about the plant has been supplied taking advantage of a direct contact with the facility through the E-PRTR registry but it was not sufficient to split the emissions. Thanks a deep survey on this category based on the analysis of IPPC permits and data available from the monitoring and control plan, the estimates for Pb, Cd, Zn, PM, NO_x e SO₂ have been revised since 2014. The analysis of this documentation will carry out further improvement such as, for example, a better allocation of emissions between combustion and process but also between the zinc and the lead production.

Following the decision 2012/17 of the Executive Body of the Convention on Long Range Transboundary Air Pollution, that requests Italy to submit information concerning the status and details of its work to improve the emission inventory of PAH, Italy in recent years has reviewed the estimates regarding PAH major sources. In the 2013 submission different recalculations have been performed in the energy and waste sector, emissions from iron and steel production have been revised in the 2014 submission. The most important update regards pig iron tapping emission factor that considers, from 2000 onwards, the abatement due to fabric filters and the relevant EF derived from the Guidebook EMEP/CORINAIR 2006 (0.95 g/Mg). Investigations on the largest integrated plant in Italy confirmed the installation of fabric filters on each point of emission related to pig iron tapping (MATTM, 2011). As regards EAF too, EF was update on the basis of a sectoral study (APAT, 2003) which reports the development of abatement technologies in the '90s in Italy and the consequent evolution in

the plants with the installation of fabric filters; but in this case the update is referred to 1990-1999 because the EF used in previous submissions concerned already abated emissions.

In Table 4.7, activity data and PAH emissions from integrated plants and from the whole sector 2C1 are reported.

Table 4.7 *Steel production data and PAH emissions from integrated plants, 1990 – 2020*

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|---------------------------|--------|--------|--------|--------|-------|-------|-------|-------|-------|-------|
| Pig iron production [Gg] | 11,852 | 11,678 | 11,209 | 11,424 | 8,555 | 5,051 | 5,071 | 4,845 | 4,619 | 3,406 |
| Steel production BOF [Gg] | 10,365 | 11,664 | 10,744 | 11,688 | 8,635 | 4,763 | 4,732 | 4,520 | 4,211 | 3,371 |
| PAH emissions i.p.* [Mg] | 41.9 | 41.3 | 11.7 | 12.1 | 9.2 | 5.8 | 5.9 | 5.7 | 5.4 | 4.1 |
| PAH emissions 2C1 [Mg] | 44.9 | 44.5 | 14.3 | 15.1 | 11.9 | 8.2 | 8.6 | 8.3 | 8.0 | 6.4 |

*i.p.: integrated plants

4.3.4 Other production (2G – 2H – 2I – 2L)

Emissions from these categories are usually negligible except for Cd emissions from use of *fireworks* accounting for 8.8% in 2020 of cadmium national totals and for NMVOC emissions from *food and drink* (2H2) accounting for 3.2% of the national total. Emissions from this last category refer to the processes in the production of bread, wine, beer and spirits. Emission factors are assumed constant for the whole time series. In Table 4.8, activity data and NMVOC emissions from sector 2H2 are reported.

Table 4.8 *Activity data and NMVOC emissions from sector 2H2, 1990 – 2020*

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|--|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Activity data - Bread [Gg] | 4,153 | 3,882 | 3,565 | 4,109 | 4,161 | 3,877 | 3,749 | 3,311 | 3,796 | 4,666 |
| Activity data – Wine [10 ⁶ dm ³] | 5,521 | 5,620 | 5,409 | 5,057 | 4,673 | 5,073 | 4,610 | 5,660 | 5,219 | 5,433 |
| Activity data – Beer [10 ⁶ dm ³] | 1,215 | 1,199 | 1,258 | 1,280 | 1,281 | 1,429 | 1,567 | 1,642 | 1,725 | 1,583 |
| Activity data – Spirits [10 ⁶ dm ³] | 268 | 232 | 206 | 161 | 115 | 98 | 101 | 100 | 108 | 111 |
| NMVOC emissions [Gg] | 31.7 | 29.2 | 26.8 | 27.5 | 25.9 | 24.3 | 23.6 | 22.0 | 24.5 | 28.3 |

4.4 QA/QC AND VERIFICATION

Activity data and emissions reported under EU-ETS and the national EPER/EPRTTR register are compared to the information provided by the industrial associations. The general outcome of this verification step shows consistency among the information collected under different legislative frameworks and information provided by the relevant industrial associations.

Every five years emissions referring to 1990-2015 are disaggregated at regional and provincial level and figures are compared with results obtained by regional bottom up inventories. From 2015 onwards the disaggregation at local level takes place every four years, so up to now the disaggregation of the national inventory covers estimates for the years: 1990, 1995, 2000, 2005, 2010, 2015 and 2019. PM10 emissions

disaggregated at local level are also used as input for air quality modelling. The distribution of PM10 emissions from the *industrial processes* sector at NUTS3 level for 2019 is reported in Figure 4.1; methodologies are described in the relevant publication (ISPRA, 2009) which is currently under revision to accommodate results and updates related to 2019 estimates.




Figure 4.1 PM10 emissions from industrial processes in 2019 (t)

4.5 RECALCULATIONS

4.5.1 Mineral industry (2A)

Recalculations occur for 1990-2019 for cement production because SO_x emission estimates for cement production result from fuel combustion and must be allocated under 1A2f. Recalculations occur also for PM10, PM2.5 along the whole time series because this submission includes PM emission estimates from category

2.A.5b construction and demolition. Recalculations occur for NMVOC emissions from lubricants use because of the update of the activity data for 2012-2019.

4.5.2 Chemical industry (2B)

Recalculations occur in 2019 for NO_x from phthalic anhydride because of the update of the EF.

4.5.3 Metal industry (2C)

Minor recalculations occur for Cd, Hg and Pb emissions in 2016-2019 because of the update of EFs for 2C1.

4.5.4 Other product use (fireworks and tobacco) (2G)

No recalculations occurred from this source category.

4.5.5 Other industrial processes (2H)

Recalculations occur in category 2H1 for NMVOC in the years 2013-2019 because of the update of the activity data (chipboard) for the same years.

4.6 PLANNED IMPROVEMENTS

Following the suggestions of the last review processes further investigations are under way for the category 2A5a in order to identify reliable activity data and corresponding emission factors.

As for 2C1, additional data collected in the context of the environmental authorization of the largest Italian (and European) iron&steel facility, allow for the review of the emissions of all pollutants starting from 2015, based on the measurement data available at installation and chimney level. The processing of those data has been going on and the revision of the estimates is foreseen for the next submission.

Activities 2C3, 2C5, 2C6 and 2C7 are under investigations to allocate emissions between combustion and process.

5 IPPU - SOLVENT AND OTHER PRODUCT USE (NFR SECTOR 2)

5.1 OVERVIEW OF THE SECTOR

In this sector all non-combustion emissions from other industrial sectors than manufacturing and energy industry are reported.

Emissions are related to the use of solvent in paint application, degreasing and dry cleaning, chemical products, manufacture and processing and other solvent use, including emissions from road paving with asphalt and asphalt roofing activities.

NMVOC emissions are estimated from all the categories of the sector as well as PM for polyester and polyvinylchloride processing, in the chemical product category, and for asphalt processes and PAH emissions from the preservation of wood in the other solvent use.

The categories included in the sector are specified in the following.

- 2D3a Domestic solvent use includes emissions from the use of solvent in household cleaning and car care products as well as cosmetics.
- 2D3b Road paving with asphalt includes emissions from the production and use of asphalt for road paving.
- 2D3c Asphalt roofing includes emissions from the manufacturing of roofing products and the blowing of asphalt.
- 2D3d1 Decorative coating includes emissions from paint application for construction and buildings, domestic use and wood products.
- 2D3d2 Industrial coating includes emissions from paint application for manufacture of automobiles, car repairing, coil coating, boat building and other industrial paint application.
- 2D3e Degreasing includes emissions from the use of solvents for metal degreasing and cleaning.
- 2D3f Dry cleaning includes emissions from the use of solvent in cleaning machines.
- 2D3g Chemical products, manufacture and processing covers the emissions from the use of chemical products such as polyurethane and polystyrene foam processing, manufacture of paints, inks and glues, textile finishing and leather tanning.
- 2D3h Printing includes emissions from the use of solvent in the printing industry
- 2D3i Other product use addresses emissions from glass wool enduction, printing industry, fat, edible and non-edible oil extraction, preservation of wood, application of glues and adhesives, vehicles dewaxing.

According to the review process we are still exploring if Hg emissions from fluorescent tubes occur in Italy. No other emissions from the sector occur.

NMVOC emissions from 2D3a, 2D3d, 2D3g, 2D3h and 2D3i are key categories in 2020; the same categories, except 2D3h plus 2D3e, were also key categories in 1990. For the trend 1990-2020, 2D3a, 2D3d and 2D3g result as key categories.

The sector accounts, in 2020, for 42.4% of total national NMVOC emissions, whereas in 1990 the weight out of the total was equal to 31.08%. Total sectoral NMVOC emissions decreased by 38.5%, between 1990 and 2020.

PM10 emissions account for 1.49%, while PM2.5, BC and PAH emissions are also estimated but they account for less than 1%.

In Figure 5.1 the share of NMVOC emissions of the sector is reported for the years 1990 and 2020.




Figure 5.1 Share of NMVOC emissions for the solvent use sector in 1990 and 2020

5.2 METHODOLOGICAL ISSUES

The sector is characterized by a multitude of activities which implies that the collection of activity data and emission factors is laborious. A lot of contacts have been established in different sectors with industrial associations and documentation has been collected even though improvements are still needed especially in some areas.

Emissions of NMVOC from solvent use have been estimated according to the methodology reported in the EMEP/EEA guidebook, applying both national and international emission factors (Vetrella, 1994; EMEP/CORINAIR, 2007; EMEP/EEA, 2016). Country specific emission factors provided by several accredited sources have been used extensively, together with data from the national EPER/PRTR registry; in particular, for paint application (Offredi, several years; FIAT, several years), solvent use in dry cleaning (ENEA/USLRMA, 1995), solvent use in textile finishing and in the tanning industries (Techne, 1998; Regione Toscana, 2001; Regione Campania, 2005; GIADA, 2006). Basic information from industry on percentage reduction of solvent content in paints and other products has been applied to EMEP/EEA emission factors in order to evaluate the reduction in emissions during the considered period.

A more detailed description is reported for the 2020 key categories of NMVOC emissions in the following sections. A description of CO emissions estimation process for asphalt roofing is also included.

5.2.1 Domestic solvent use (2D3a)

The category comprises a lot of subcategories whose emissions, specifically NMVOC, originate from the use of solvent in household cleaning and car care products as well as cosmetics.

Emissions from this category have been calculated using a detailed methodology, based on VOC content per type of consumer product. Emissions from domestic solvent use comprise emissions from the use of products for household and cleaning and for cosmetics which are derived as described in the following.

Activity data

Activity data are expressed as the sum, in tonnes, of household and cleaning products and cosmetics.

Household and cleaning products: data are communicated by the National Association of Detergents and Specialties for industry and home care (Assocasa, several years) either by personal communications or

Association Reports and refer to the consumption of soaps and detergents and cleaning and maintenance products.

Cosmetics: data are the sum of cosmetics products in aerosol form and other cosmetics.

Figures of cosmetics in aerosol form are provided by the Italian Aerosol Association (AIA, several years [a] and [b]) and refer to the number of pieces of products sold for personal care (spray deodorants, hair styling foams and other hair care products, shaving foams, and other products). These figures are then converted in tonnes by means of the capacity of the different cosmetics containers.

Figures for other cosmetics products are derived by the Production Statistics Database (Prodcom) supplied by the National Institute of Statistics (ISTAT, several years [a] and [b]) by difference with the previous aerosol data.

Time series of cosmetics production is reconstructed by means of the annual production index, considering the year 2000 as the base year because this is the year where production national statistics and Prodcom data coincide. The next step is the calculation of apparent consumption taking into account import-export data derived by the National Association of Cosmetic Companies (UNIPRO, several years). Since these figures also include aerosol cosmetics, the amount of aerosol cosmetics is subtracted.

Final consumption is therefore estimated.

Emission factors

NMVOC emission factors are expressed in percentage of solvent contained in products.

Household and cleaning products: figures are communicated by the relevant industrial association, ASSOCASA, by personal communications. For leather, shoes, wood etc. and car maintenance products, figures are taken from BiPro Association. For insecticides and disinfectants, emission factors derive from national studies at local level.

Cosmetics: for aerosol cosmetics, the emission factor is communicated by the Italian Aerosol Association for the year 2004 and supposed constant from 1995. For other cosmetics, information from BiPro has been considered (EC report 'Screening study to identify reductions in VOC emissions due to the restrictions in the VOC content of products', year 2002 (EC, 2002)), and supposed constant from 1996.

5.2.2 Asphalt Roofing (2D3c)

The category includes CO and NMVOC emissions from the application of asphalt to insulate (waterproofing) the roof of buildings. As requested by the last review process, further information is included about the estimation of CO emissions. The emission factors for CO and NMVOC emissions from asphalt roofing are set constant along the whole timeseries and are equal to 38 g CO/10³ m² and 0.085 g NMVOC/10³ m². The activity data are yearly provided by the national association of producers while the EF is taken from the EMEP/EEA Guidebook with the assumption that 1000 m² of asphalt material corresponds to 4 Mg. The following table describes AD, EF and emissions for CO and NMVOC between 1990 and 2020.

Table 5.1 AD, EF and emissions for CO and NMVOC from asphalt roofing, 1990-2020.

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|---|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Activity data – Asphalt roofing [10 ⁶ m ²] | 200.0 | 200.0 | 200.0 | 204.0 | 204.0 | 152.2 | 127.7 | 134.4 | 122.9 | 117.6 |
| EF CO –[g CO/10 ³ m ²] | 38 | 38 | 38 | 38 | 38 | 38 | 38 | 38 | 38 | 38 |
| CO emissions [Gg] | 0.007 | 0.007 | 0.007 | 0.009 | 0.007 | 0.006 | 0.005 | 0.005 | 0.005 | 0.005 |
| EF NMVOC [g NMVOC/10 ³ m ²] | 0.085 | 0.085 | 0.085 | 0.085 | 0.085 | 0.085 | 0.085 | 0.085 | 0.085 | 0.085 |
| NMVOC emissions [Gg] | 0.017 | 0.017 | 0.017 | 0.020 | 0.017 | 0.013 | 0.011 | 0.011 | 0.010 | 0.010 |

5.2.3 Decorative coating (2D3d1)

The category includes NMVOC emissions from the application of paint for construction and buildings, domestic use and wood products.

Activity data on the consumption of paint for construction and buildings and related domestic use are provided by the Ministry of Productive Activities for 1990 and 1991 (MICA, 1999) and updated on the basis of production figures provided annually by the National Institute of Statistics (ISTAT, several years [a] and [b]).

From 2007 onwards, data are also provided by SSOG (Stazione Sperimentale per le industrie degli Oli e dei Grassi, *Experimental Station for Oils and Fats Industries*), which collects information and data regarding national production and imports for paint categories set out in the directive 2004/42/EC on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products. The purpose of this directive is to limit the total content of VOCs in certain paints and varnishes and vehicle refinishing products in order to prevent or reduce air pollution resulting from the contribution of VOCs to the formation of tropospheric ozone. The directive sets maximum VOCs content limit values for some paints and varnishes.

As for emission factors, those for construction and buildings are taken from the EMEP/EEA guidebook and are considered constant till 2009, whereas the default values for domestic use vary in consideration of the different share between solvent and water content in paint throughout the years. In particular, the variation of emission factor from 1990 to 2000 is equal to 35% - 65% up to 25% - 75% in 2000, on the basis of qualitative information supplied by industry on the increase of water based paints products in the market. From 2010, emission factors are calculated taking into account maximum VOC content limit values for paint and varnishes set out in Annex II A of Directive 2004/42/EC and data collected by SSOG. The comparison of national emission estimates for this category with those produced by IIASA for 2010 resulted in similar values.

On the other hand, information on activity data and emission factors for emissions from wood products are provided by the national association of wood finishing (Offredi, several years). Emission factors have been calculated for 1990, 1998 and 2003 on the basis of information provided by the industrial association distinguishing the different type of products which contain different solvent percentages. Data have been supplied also for the years 2005 and 2006. Actually, we are keeping constant the 2006 value unless the association provides us with updated information. For previous years, values have been interpolated.

In this category, emissions from paint application in wood are one of the biggest contributors to national NMVOC emissions and the relevant share has grown considerably in recent years. NMVOC emissions due to the use of paint and other products except from industrial coating could not be controlled properly in the past since the EU Directive 2004/42/EC entered into force. This directive, transposed into the Italian legislation in 2004, sets out maximum VOC content for many paint, varnishes and vehicle refinishing products that had to be achieved in two steps. The early limit values, to be respected from 2007 till 2009, did not lead to a significant reduction of NMVOC emissions, while the latest values, that had to be respected from 2010 onwards, brought to a significant decrease.

5.2.4 Industrial coating (2D3d2)

The category includes emissions from paint application for manufacture of automobiles, car repairing, coil coating, boat building and other industrial paint application.

Activity data on the number of vehicles are provided by the National Automobile Association (ACI, several years) in the Annual Statistical Report and the emission factors are those reported by the main automobile producers on the relevant activity in their environmental reports and communicated from 2003 in the framework of E-PRTR.

For the paint used in car repairing, activity data are provided by the Ministry of Productive Activities for 1990 and 1991 (MICA, 1999) and updated on the basis of production figures provided annually by the National Institute of Statistics (ISTAT, several years [a] and [b]). The default emission factor (provided by the EMEP guidebook) used from 1990 to 1995 equal to 700 g/kg paint is also confirmed by the European guidelines for car repairing provided by the Conseil Europeen de l'Industrie des Peintures (CEPE, 1999). The reduction of

the emission factor in 1999 (13% of 1995) is applied on the basis of information on different shares between solvent and water based paint throughout the years provided by the national study PINTA, *Piano nazionale di tutela della qualità dell'aria* (ENEA, 1997). From 1996 to 1999 the reduction is linear. From 1999 to 2006 the value is kept constant. From 2007 onwards emission factors have been calculated taking into account the maximum VOC content limit values for paint and varnishes set out in Annex II B of Directive 2004/42/EC and data collected by SSOG. The Italian implied emission factor is the weighted average of the different products used in this activity where data are collected at detailed level and communicated within the European Directive. The trend is driven by the increase in the last years of the use of primers and special finishes. Similar trend is noted for the construction and building and domestic paints where the variability is mainly due to the percentage of solvent based paint product used out of the total paints.

Concerning coil coating, boat building and other industrial paint application, activity data are provided by the Ministry of Productive Activities for 1990 and 1991 (MICA, 1999) and updated annually by the National Institute of Statistics (ISTAT, several years [a] and [b]). Emission factors are taken from the EMEP guidebook considering the national legislation where relevant.

Emission factors of the other industrial paint application from 1990 to 1995 are constant and derive from the 1999 EMEP/CORINAIR guidebook. The reduction of the emission factor from 1996 to 2004 is applied on the basis of information on different share of paints throughout the years provided by the national study PINTA. From 2010, the value of the 1999 Guidebook has been chosen considering the further reduction of the sector (in PINTA, the reduction for 2005 with respect to 1995 is equal to 37%, and for 2010 64%; considering the default emission factor 250 g/kg of paint, the reduction is equal to 53%).

NMVOC emissions from this category have been decreasing constantly since the nineties, when all industrial installations have been subjected to permits from local authorities. Since then, most of the installations have to comply with emission limit values and technological requirements imposed at regional level, taking into account the EU directives on industrial emissions (i.e. Directive 99/13/EC on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations (EC, 1999)) and often going beyond the European legislation. With regard to car repairing the emission cut from 2007 onwards is mainly due to the maximum contents of VOC set by EU Directive 2004/42/EC (EC, 2004).

5.2.5 Degreasing (2D3e)

NMVOC emissions have been estimated for this category. The emission factor used is equal to 1000 g NMVOC/kg solvent and refers to the percentage of NMVOC emitted per tones of solvent used and it is not directly comparable to those provided in the Guidebook (700 g/kg cleaning product). According to the information provided by the National Industrial Association, due to technological improvements, the amount of solvent used in the products decreased during the period whereas it has been assumed that the percentage of NMVOC emissions remains constant. The EF used in the inventory comes from the “Corinair 1992 Default Emission Factors Handbook” but, taking in consideration the comments from the last reviews and lacking information about abatement technologies, it has been assumed that all the solvent goes into the atmosphere; so the EF is equal to 1000 g NMVOC/kg solvent from the last submission. Activity data, solvent used, are also provided by the relevant industrial association (Federchimica, several years). According to the review process we are verifying how to apply the emission factors available in the EMEP/EEA Guidebook that refers to the volumes of cleaning products used instead of the solvent used but the information on the amount of cleaning products in the sector is not yet available.

Table 5.2 AD, EF and emissions for NMVOC from metal degreasing 2D3e, 1990-2020.

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|---------------------------------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Activity data – Degreasing [Gg] | 52.8 | 32.8 | 25.9 | 22.2 | 19.1 | 16.4 | 15.4 | 15.0 | 14.5 | 14.1 |
| EF NMVOC [g NMVOC/kg] | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 | 1,000 |
| NMVOC emissions [Gg] | 52.8 | 32.8 | 25.9 | 22.2 | 19.1 | 16.4 | 15.4 | 15.0 | 14.5 | 14.1 |

5.2.6 Dry cleaning (2D3f)

Concerning dry cleaning, activity data, equal to 30,000 machines, remain unchanged throughout the time series and the emission factor is calculated based on the allocation of machines to closed-circuit (CCM) and open-circuit (OCM). Different amounts of solvent are used in these machines and have different emission factors. The emission factors are calculated assuming that in 1990 the closed-circuit machines were 60%, 90% in 1995 and in up to 100% in 1999.

The average consumption of solvent per machine is equal to 258 kg/year for CCM and 763 kg/year for OCM, as derived from a national study by ENEA/USL-RMA (ENEA/USL-RMA, 1995). It is assumed that only perchlorethylene is used. These values are multiplied by the emission factors of the Guidebook EMEP, referred to the amount of solvent consumed (equal to 0.4 and 0.8 kg/kg of solvent, for CCM and OCM, respectively) and then the average annual emission factor was calculated based on the percentage distribution of closed and open circuit machines.

5.2.7 Chemical products, manufacture and processing (2D3g)

The category comprises emissions from the use of chemical products such as polyester, polyurethane, polyvinylchloride and polystyrene foam processing, manufacture of paints, inks and glues, textile finishing and leather tanning.

Activity data for polystyrene and polyurethane are derived from the relevant industrial associations, and ISTAT (ISTAT, several years [a] and [b]), whereas emission factors are from the EMEP/CORINAIR guidebook. For what concerns polyurethane, the relevant national industrial association has communicated that the phase out of CFC gases occurred in the second half of nineties and the blowing agent currently used is penthane. Because of manufacturing plant have abatement system in place PM emissions could all be considered as PM_{2.5}.

As for polyvinylchloride (PVC), activity data and emission factors are supplied in the framework of the national PRTR. NMVOC emissions are entirely attributed to the phase of PVC production; no use of solvents occurs in the PVC processing. This information has been provided by the relevant industrial plant, EVC Italy, in 2001. Because of manufacturing plant have abatement system in place PM emissions could all be considered as PM_{2.5}.

For the other categories, activity data are provided by the relevant industrial associations and by ISTAT, while emission factors are taken from the EMEP/CORINAIR guidebook considering national information on the solvent content in products supplied by the specific industrial associations.

As regard rubber processing, emission factors for the first years of nineties have been provided by the industrial association. The use of the Swedish emission factor from 1997 was justified in lack of other updated data.

For the glues manufacturing category, emission factors for 1990 are derived from the 1992 EMEP/CORINAIR guidebook. The trend of emission factor is estimated on the basis of the trend of the emission factor for consumption of glue (as indicated by the industrial association). From 1995 to 2004, the industrial association communicated data on consumption and solvent content by product. The reductions from 2000 are based on the assumptions of PINTA. From 2004 the emission factor has been assumed constant in lack of updated information. For previous years, values have been interpolated.

As regards leather tanning, emission factor for 1990 is from Legislative Decree 152/2006, equal to the maximum VOC content limit value (150 g/m²). For 2000 and 2003, emission factors have been calculated on the basis of emission figures derived by the national studies on the major leather tanning industries and statistical production.

As regards asphalt blowing and possible PAH and Benzo(a)pyrene emissions as suggested by the 2016 EMEP/EEA Guidebook, according to the relevant industrial association PAH emissions are negligible because all the asphalt blowing plants have abatement filter system of PM and afterburners of gas. Moreover, these

plants should respect national environmental legislation not exceeding at the stack more than 0.1mg/Nm³ for total PAH. For this pollutant the relevant notation key NE has been used.

As requested by the last review process, Italy has revised PM10 and PM2.5 emissions coming from polyester and polyvinylchloride processing. No PM emission factors are provided in the EMEP EEA 2019 GB for these activities, so after checking with the relevant industries the values of the EFs have been differentiated for the activities concerned with PM10 assumed as 80% of TSP and PM2.5 set equal to 60% of TSP (as suggested in the 2019 EMEP/EEA GB, 2B Chemical industry, paragraph 3.2.2.1, pag.14).

Table 5.3 EF and emissions for PM2.5 from metal Chemical products 2D3g, 1990-2020.

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|---|------|------|------|------|------|------|------|------|------|------|
| EF PM10 polyester proc. [g PM10/Mg] | 92 | 92 | 92 | 92 | 92 | 92 | 92 | 92 | 92 | 92 |
| EF PM10 polyvinylchloride proc. [g PM10/Mg] | 24.0 | 20.8 | 9.2 | 8.8 | 8.8 | 8.8 | 8.8 | 8.8 | 8.8 | 8.8 |
| EF PM2.5 polyester proc. [g PM2.5/Mg] | 69 | 69 | 69 | 69 | 69 | 69 | 69 | 69 | 69 | 69 |
| EF PM2.5 polyvinylchloride proc. [g PM2.5/Mg] | 18 | 15.6 | 6.9 | 6.6 | 6.6 | 6.6 | 6.6 | 6.6 | 6.6 | 6.6 |
| Total PM10 emissions [Gg] | 0.03 | 0.03 | 0.02 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| Total PM2.5 emissions [Gg] | 0.02 | 0.02 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |

5.2.8 Other product use (2D3i)

The category includes NMVOC emissions from the application of glues and adhesives, which account for most of emissions from the category, emissions from fat, edible and non-edible oil extraction and minor emissions from glass wool enduction.

Activity data and emission factors for the application of glues and adhesives had been provided by the relevant industrial association up to 2004. After that period, activity data have been updated on the basis of information by ISTAT (ISTAT, several years [a] and [b]) whereas the emission factor is considered constant in absence of further information.

For fat, edible and non-edible oil extraction activity data derive from the FAOSTAT database (<http://faostat.fao.org>) whereas default emission factors do not change over the period.

5.3 TIME SERIES AND KEY CATEGORIES

The sector accounts, in 2020, for about 42.4% of total national NMVOC emissions. PM, BC and PAH emissions are also estimated in this sector but they account for less than 1%.

NMVOC emissions from the use of solvent decreased from 1990 to 2020 of about 38.5%, from 619 Gg in 1990 to 375 Gg in 2020, mainly due to the reduction of emissions in paint application, in degreasing and dry cleaning and in other product use. The general reduction observed in the emission trend of the sector is

due to the implementation of the European Directive 1999/13/EC (EC, 1999) on the limitation of emissions of volatile organic compounds due to the use of organic solvents, entered into force in Italy in January 2004, and the European Directive 2004/42/EC (EC, 2004), entered into force in Italy in March 2006, which establishes a reduction of the solvent content in products. Moreover, the reduction of emissions from paint application, is also due to the implementation of the Italian Legislative Decree 161/2006.

Figure 5.2 shows emission trends from 1991 to 2020 with respect to 1990 by sub-sector.




Figure 5.2 Trend of NMVOC emissions from 1991 to 2020 as compared to 1990

The main source of emissions is *paint application* (2D3d) where NMVOC emissions derive mainly from wood application and construction and building. The second source of emissions is *domestic solvent use* (2D3a), mostly for the consumption of cosmetics, followed by *chemical products and other product use* (2D3g), especially for emissions deriving from polyurethane processing, paints manufacturing and leather tanning.

Table 5.1 represents the pollutants estimated in the sector and the key categories identified.

Table 5.1 Key categories in the IPPU - Solvent and other product use sector in 2020

| | 2D3a | 2D3b | 2D3c | 2D3d | 2D3e | 2D3f | 2D3g | 2D3h | 2D3i |
|-----------------|-------|------|------|-------|------|------|------|------|------|
| SO _x | | | | | | | | | |
| NO _x | | | | | | | | | |
| NH ₃ | | | | | | | | | |
| NMVOC | 14.32 | 0.98 | 0.00 | 15.23 | 1.59 | 0.35 | 5.68 | 2.02 | 3.17 |
| CO | | | 0.00 | | | | | | |
| PM10 | | 1.46 | 0.03 | | | | 0.01 | | |
| PM2.5 | | 0.24 | 0.01 | | | | 0.01 | | |
| BC | | 0.12 | 0.00 | | | | | | |
| Pb | | | | | | | | | |
| Cd | | | | | | | | | |
| Hg | | | | | | | | | |
| PAH | | | | | | | | | 0.02 |
| Dioxin | | | | | | | | | |
| HCB | | | | | | | | | |
| PCB | | | | | | | | | |

Note: key categories are shaded in blue

In Table 5.2 and 5.3 activity data and emission factors used to estimate emissions from the sector are reported at SNAP code level.

A strong decrease in the content of solvents in the products in the nineties is observed.

| | | | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--------------|---|--------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| 06 01 | Paint application | | | | | | | | | | | | |
| 06 01 01 | Paint application : manufacture of automobiles | <i>vehicles</i> | 2.865.857 | 2.521.355 | 2.770.104 | 1.766.930 | 1.310.425 | 1.326.711 | 1.433.047 | 1.499.956 | 1.443.870 | 1.293.241 | 1.118.952 |
| 06 01 02 | Paint application : car repairing | <i>Mg paint</i> | 22.250 | 17.850 | 24.276 | 23.475 | 19.479 | 25.395 | 32.521 | 35.217 | 38.728 | 37.416 | 16.899 |
| 06 01 03 | Paint application : construction and buildings (except item 06.01.07) | <i>Mg paint</i> | 111.644 | 120.736 | 125.928 | 163.455 | 168.358 | 158.661 | 159.823 | 157.265 | 161.199 | 196.105 | 177.383 |
| 06 01 04 | Paint application : domestic use (except 06.01.07) | <i>Mg paint</i> | 420.000 | 420.000 | 420.000 | 420.000 | 420.000 | 420.000 | 420.000 | 420.000 | 420.000 | 420.000 | 420.000 |
| 06 01 05 | Paint application : coil coating | <i>Mg paint</i> | 14.500 | 14.500 | 14.500 | 14.500 | 14.500 | 14.500 | 14.500 | 14.500 | 14.500 | 14.500 | 14.500 |
| 06 01 06 | Paint application : boat building | <i>Mg paint</i> | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 |
| 06 01 07 | Paint application : wood | <i>Mg paint</i> | 150.000 | 150.000 | 140.000 | 140.000 | 123.250 | 80.000 | 75.000 | 80.000 | 75.000 | 75.000 | 70.000 |
| | Other industrial paint application | <i>Mg paint</i> | 125.000 | 125.000 | 125.000 | 125.000 | 125.000 | 125.000 | 125.000 | 125.000 | 125.000 | 125.000 | 125.000 |
| 06 02 | Degreasing, dry cleaning and electronics | | | | | | | | | | | | |
| 06 02 01 | Metal degreasing | <i>Mg solvents</i> | 52.758 | 32.775 | 25.895 | 22.237 | 19.095 | 16.398 | 15.906 | 15.429 | 14.966 | 14.517 | 14.081 |
| 06 02 02 | Dry cleaning | <i>machines</i> | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 |
| 06 03 | Chemical products manufacturing or processing | | | | | | | | | | | | |
| 06 03 01 | Polyester processing | <i>Mg product</i> | 179.852 | 197.882 | 168.704 | 112.188 | 89.638 | 94.389 | 96.522 | 109.510 | 109.510 | 109.510 | 109.510 |
| 06 03 02 | Polyvinylchloride processing | <i>Mg product</i> | 617.600 | 575.600 | 405.285 | 348.497 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 06 03 03 | Polyurethane processing | <i>Mg product</i> | 145.700 | 230.633 | 350.187 | 175.278 | 196.585 | 196.585 | 196.585 | 196.585 | 196.585 | 196.585 | 196.585 |
| 06 03 04 | Polystyrene foam processing (c) | <i>Mg product</i> | 85.004 | 80.400 | 90.200 | 35.200 | 33.692 | 46.800 | 36.200 | 51.200 | 44.100 | 51.600 | 42.600 |
| 06 03 05 | Rubber processing | <i>Mg product</i> | 671.706 | 700.859 | 810.124 | 831.187 | 607.667 | 545.989 | 557.079 | 613.364 | 631.709 | 603.318 | 622.427 |
| 06 03 06 | Pharmaceutical products manufacturing | <i>Mg product</i> | 80.068 | 88.094 | 104.468 | 106.861 | 110.183 | 120.907 | 126.068 | 131.052 | 137.321 | 136.202 | 135.174 |
| 06 03 07 | Paints manufacturing | <i>Mg product</i> | 697.129 | 747.417 | 900.683 | 964.631 | 891.882 | 851.450 | 770.497 | 940.682 | 905.943 | 885.576 | 943.407 |
| 06 03 08 | Inks manufacturing | <i>Mg product</i> | 87.527 | 110.667 | 132.256 | 132.521 | 133.979 | 108.600 | 102.949 | 92.708 | 115.533 | 112.953 | 120.329 |
| 06 03 09 | Glues manufacturing | <i>Mg product</i> | 111.683 | 266.169 | 302.087 | 331.770 | 317.560 | 249.152 | 259.393 | 242.425 | 279.154 | 286.629 | 276.060 |
| 06 03 10 | Asphalt blowing | <i>Mg product</i> | 77.248 | 70.336 | 77.408 | 88.896 | 65.000 | 25.000 | 21.000 | 20.000 | 20.000 | 18.000 | 16.000 |
| 06 03 12 | Textile finishing | <i>1000 m2</i> | 1.332.679 | 1.301.105 | 1.173.047 | 987.705 | 831.236 | 631.573 | 643.518 | 612.500 | 614.774 | 543.469 | 518.522 |
| 06 03 13 | Leather tanning | <i>1000 m2</i> | 173.700 | 183.839 | 200.115 | 157.891 | 186.824 | 162.500 | 169.668 | 244.724 | 173.387 | 139.831 | 129.350 |
| 06 04 | Other use of solvents and related activities | | | | | | | | | | | | |
| 06 04 01 | Glass wool enduction | <i>Mg product</i> | 105.029 | 119.120 | 139.421 | 129.958 | 115.332 | 86.929 | 86.498 | 87.208 | 98.805 | 99.552 | 88.319 |
| 06 04 02 | Mineral wool enduction | <i>Mg product</i> | 0 | 11.000 | 18.000 | 20.500 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 06 04 03 | Printing industry | <i>Mg ink</i> | 73.754 | 91.667 | 100.690 | 111.550 | 98.206 | 79.604 | 79.106 | 79.142 | 98.626 | 96.424 | 102.721 |
| 06 04 04 | Fat, edible and non edible oil extraction | <i>Mg product</i> | 5.070.398 | 7.560.387 | 6.539.796 | 7.939.548 | 7.088.890 | 6.147.566 | 5.307.964 | 5.514.461 | 4.886.301 | 6.980.365 | 6.936.497 |
| 06 04 05 | Application of glues and adhesives | <i>Mg product</i> | 98.500 | 234.751 | 266.996 | 292.687 | 280.150 | 219.801 | 228.836 | 213.866 | 246.269 | 252.863 | 243.540 |
| 06 04 08 | Domestic solvent use (other than paint application)(k) | <i>Mg product</i> | 1.938.779 | 2.282.020 | 2.410.338 | 2.767.759 | 2.614.274 | 2.265.605 | 2.226.736 | 2.248.707 | 2.346.757 | 2.376.579 | 2.748.918 |
| 06 04 09 | Vehicles dewaxing | <i>vehicles</i> | 2.540.597 | 1.740.212 | 2.361.075 | 2.238.344 | 1.972.070 | 1.594.259 | 1.849.608 | 1.994.407 | 1.945.120 | 1.949.554 | 1.441.385 |

Table 5.2 Activity data in the IPPU - Solvent and other product use sector

| | | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|----------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| Paint application | | | | | | | | | | | | |
| Paint application : manufacture of automobiles | <i>g/vehicles</i> | 8.676 | 6.296 | 4.833 | 4.065 | 2.854 | 3.037 | 3.080 | 3.023 | 2.870 | 2.614 | 2.685 |
| Paint application : car repairing | <i>g/Mg paint</i> | 700.000 | 700.000 | 605.500 | 605.500 | 497.810 | 617.377 | 587.616 | 567.101 | 551.450 | 527.187 | 434.596 |
| Paint application : construction and buildings (except item 06.01.07) | <i>g/Mg paint</i> | 300.000 | 300.000 | 300.000 | 300.000 | 200.000 | 152.412 | 149.364 | 222.552 | 220.326 | 209.310 | 198.844 |
| Paint application : domestic use (except 06.01.07) | <i>g/Mg paint</i> | 126.450 | 113.100 | 99.750 | 99.750 | 67.710 | 54.360 | 54.360 | 75.720 | 73.050 | 72.516 | 77.370 |
| Paint application : coil coating | <i>g/Mg paint</i> | 200.000 | 200.000 | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 | 10.000 |
| Paint application : boat building | <i>g/Mg paint</i> | 750.000 | 750.000 | 622.500 | 475.417 | 340.000 | 340.000 | 340.000 | 340.000 | 340.000 | 340.000 | 340.000 |
| Paint application : wood | <i>g/Mg paint</i> | 446.500 | 425.000 | 406.300 | 390.750 | 377.250 | 354.000 | 342.500 | 340.000 | 339.500 | 333.250 | 313.750 |
| Other industrial paint application | <i>g/Mg paint</i> | 530.000 | 530.000 | 439.900 | 337.583 | 250.000 | 250.000 | 250.000 | 250.000 | 250.000 | 250.000 | 250.000 |
| Degreasing, dry cleaning and electronics | | | | | | | | | | | | |
| Metal degreasing | <i>g/Mg solvents</i> | 1.000.000 | 1.000.000 | 1.000.000 | 1.000.000 | 1.000.000 | 1.000.000 | 1.000.000 | 1.000.000 | 1.000.000 | 1.000.000 | 1.000.000 |
| Dry cleaning | <i>g/machines</i> | 306.000 | 154.000 | 103.000 | 103.000 | 103.000 | 103.000 | 103.000 | 103.000 | 103.000 | 103.000 | 103.000 |
| Chemical products manufacturing or processing | | | | | | | | | | | | |
| Polyester processing | <i>g/Mg product</i> | 325 | 325 | 325 | 325 | 325 | 325 | 325 | 325 | 325 | 325 | 325 |
| Polyvinylchloride processing | <i>g/Mg product</i> | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Polyurethane processing | <i>g/Mg product</i> | 120.000 | 110.000 | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 |
| Polystyrene foam processing (c) | <i>g/Mg product</i> | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 | 60.000 |
| Rubber processing | <i>g/Mg product</i> | 12.500 | 10.000 | 8.000 | 8.000 | 8.000 | 8.000 | 8.000 | 8.000 | 8.000 | 8.000 | 8.000 |
| Pharmaceutical products manufacturing | <i>g/Mg product</i> | 55.000 | 55.000 | 55.000 | 55.000 | 55.000 | 55.000 | 55.000 | 55.000 | 55.000 | 55.000 | 55.000 |
| Paints manufacturing | <i>g/Mg product</i> | 15.000 | 15.000 | 15.000 | 13.110 | 10.863 | 9.524 | 11.134 | 10.175 | 10.748 | 10.769 | 10.375 |
| Inks manufacturing | <i>g/Mg product</i> | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 | 30.000 |
| Glues manufacturing | <i>g/Mg product</i> | 20.000 | 5.041 | 3.603 | 2.806 | 2.806 | 2.806 | 2.806 | 2.806 | 2.806 | 2.806 | 2.806 |
| Asphalt blowing | <i>g/Mg product</i> | 544 | 544 | 544 | 544 | 544 | 544 | 544 | 544 | 544 | 544 | 544 |
| Textile finishing | <i>g/1000 m2</i> | 296 | 296 | 296 | 296 | 296 | 296 | 296 | 296 | 296 | 296 | 296 |
| Leather tanning | <i>g/1000 m2</i> | 150.000 | 150.000 | 125.000 | 105.378 | 82.267 | 71.000 | 71.000 | 71.000 | 71.000 | 71.000 | 71.000 |
| Other use of solvents and related activities | | | | | | | | | | | | |
| Glass wool enduction | <i>g/Mg product</i> | 800 | 800 | 800 | 800 | 800 | 800 | 800 | 800 | 800 | 800 | 800 |
| Mineral wool enduction | <i>g/Mg product</i> | 300 | 300 | 300 | 300 | 300 | 300 | 300 | 300 | 300 | 300 | 300 |
| Printing industry | <i>g/Mg ink</i> | 234.649 | 228.190 | 184.332 | 174.227 | 174.227 | 174.227 | 174.227 | 174.227 | 174.227 | 174.227 | 174.227 |
| Fat, edible and non edible oil extraction | <i>g/Mg product</i> | 790 | 704 | 706 | 691 | 700 | 701 | 718 | 693 | 715 | 741 | 740 |
| Application of glues and adhesives | <i>g/Mg product</i> | 600.000 | 151.230 | 108.086 | 84.190 | 84.190 | 84.190 | 84.190 | 84.190 | 84.190 | 84.190 | 84.190 |
| Domestic solvent use (other than paint application)(k) | <i>g/Mg product</i> | 60.117 | 52.262 | 42.356 | 46.153 | 42.172 | 34.483 | 32.598 | 33.802 | 37.080 | 38.263 | 46.140 |
| Vehicles dewaxing | <i>g/vehicles</i> | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 |

Table 5.3 NMVOC Emission factors in the IPPU - Solvent and other product use sector

5.4 QA/QC AND VERIFICATION

Data production and consumption time series for some activities (paint application in constructions and buildings, polyester processing, polyurethane processing, pharmaceutical products, paints manufacturing, glues manufacturing, textile finishing, leather tanning, fat edible and non-edible oil extraction, application of glues and adhesives) are checked with data acquired by the National Statistics Institute (ISTAT, several years [a], [b] and [c]), the Sectoral Association of the Italian Federation of the Chemical Industry (AVISA, several years) and the Food and Agriculture Organization of the United Nations (FAO, several years). For specific categories, emission factors and emissions are also shared with the relevant industrial associations; this is particularly the case of paint application for wood, some chemical processes and anaesthesia and aerosol cans.

In the framework of the MeditAIRaneo project, ISPRA commissioned to Techne Consulting S.r.l. a survey to collect national information on emission factors in the solvent sector. The results, published in the report “*Rassegna dei fattori di emissione nazionali ed internazionali relativamente al settore solventi*” (TECHNE, 2004), have been used to verify and validate emission estimates. In 2008, ISPRA commissioned to Techne Consulting S.r.l. another survey to compare emission factors with the last update figures published in the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007). The results are reported in “*Fattori di emissione per l'utilizzo di solventi*” (TECHNE, 2008) and have been used to update emission factors for polyurethane and polystyrene foam processing activities.

In addition, for paint application, data communicated from the industries in the framework of the EU Directive 2004/42, implemented by the Italian Legislative Decree 161/2006, on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products have been used as a verification of emission estimates. These data refer to the composition of the total amount of paints and varnishes (water and solvent contents) in different subcategories for interior and exterior use and the total amount of products used for vehicle refinishing and they are available from the year 2007.

Verifications of the emissions from the sector occurred in 2012, on account of the bilateral independent review between Italy and Spain and the revision of national estimates and projections in the context of the National emission ceilings Directive for the EU Member States and the Gothenburg Protocol of the Convention on Long-Range Transboundary Air Pollution (CLRTAP). The analysis by category did not highlight the need of major methodological revisions of the sector; an additional source of emissions was added affecting only NMVOC emissions.

Furthermore, every five years (for inventory years from 1990 to 2015) and every four years for inventory years from 2015 onwards ISPRA carries out emission estimates at NUTS level which is the occasion of an additional check with local environmental agencies.

The distribution of NMVOC emissions from the *solvent and other product use* sector at NUTS3 level for 2019 is reported in Figure 5.3; methodologies are described in the relevant publication (ISPRA, 2009) which ISPRA is going to update in 2022.




Figure 5.3 NMVOC emissions from solvent and other product use in 2019 (t)

5.5 RECALCULATIONS

Minor recalculations occurred because of the update of activity data in the manufacture of fat edible, in 2019.

5.6 PLANNED IMPROVEMENTS

Specific developments will regard the improvement of emission factors for some relevant categories. In particular, several improvements are planned with the aim to update the status of technologies in this sector where main challenges regard the availability of data collected from the industry. Main focus will be on metal degreasing and leather production where the EFs used need to be updated.

6 AGRICULTURE (NFR SECTOR 3)

6.1 OVERVIEW OF THE SECTOR

The agriculture sector is responsible for the largest part of NH₃ emissions, and contributes also to PM₁₀, PM_{2.5}, BC, TSP, NO_x, NMVOC, CO, SO₂, heavy metals (As, Cr, Cu, Ni, Se, Zn, Pb, Cd, Hg), Dioxins, PAH and HCB emissions. Italy estimates agricultural emissions for manure management (3B), agricultural soils (3D) including the use of pesticides, and field burning of agricultural wastes (3F). NO_x emissions are reported as NO₂.

In 2020, key categories level was identified for NH₃ emissions (3B1a, 3B1b, 3B3, 3B4gii, 3Da1 and 3Da2a), for NMVOC emissions (3B1a and 3B1b), for NO_x emissions (3Da1), for PM₁₀ emissions (3Dc) and for HCB emissions (3Df). In 1990 similar figures were obtained except for NH₃ emissions 3B4gii and NO_x emissions which were not key categories and PM₁₀ emissions 3B4gii which were key categories. For the trend analysis, key categories were related to NH₃ emissions (3B1a, 3B3, 3B4a, 3B4gi, 3B4gii, 3Da2a and 3Da2c), NO_x emissions (3Da2a), NMVOC emissions (3B1a and 3B1b), PM₁₀ emissions (3Dc) and HCB (3Df).

In 2020, NH₃ emissions from the agriculture sector were 345.0 Gg (95.1% of national emissions) where 3B, 3D and 3F categories represent 54.0%, 41.0% and 0.1% of total national emissions. The trend of NH₃ from 1990 to 2020 shows a 23.6% decrease due to the reduction in the number of animals, the diffusion of best environmental practices in manure management in relation to housing, storage and land spreading systems, the decrease of cultivated surface/crop production and use of N-fertilisers. A representation of the contribution by source of agriculture NH₃ emissions for 1990 and 2020 is shown in Figure 6.1.




Figure 6.1 Share of NH₃ emissions in the agriculture sector for 1990 and 2020

Agricultural official statistics are mainly collected from the National Institute of Statistics, ISTAT. Most important activity data (number of animals, N-fertilizers, agricultural surface and production, milk production) are available on-line: <http://dati.istat.it/>. ISTAT has a major role in the comprehensive collection of data through structural (such as the Farm Structure Survey, FSS) and conjunctural surveys, and the general agricultural census¹. For consistency reasons the same agricultural official statistics are used for UNFCCC and UNECE/CLRTAP emission inventory.

ISPRA participates to the Agriculture, Forestry, and Fishing Quality Panel, which has been established to monitor and improve national statistics. This is the opportunity to get in touch with experts from the Agriculture Service from ISTAT in charge for main agricultural surveys. In this way, data used for the inventory is continuously updated according to the latest information available.

Agricultural statistics reported by ISTAT are also published in the European statistics database² (EUROSTAT). The verification of statistics is part of the QA/QC procedures; therefore, as soon as outliers are identified ISTAT and category associations are contacted.

In Table 6.1 the time series of animal categories is shown.

¹ The last census was conducted in 2010 and data are available at the link <http://dati-censimentoagricoltura.istat.it/>

² <http://ec.europa.eu/eurostat/data/database>

Table 6.1 *Time series of animals*

| Year | Dairy cattle | Non-dairy cattle | Buffalo | Sheep | Goats | Horses | Mules/a sses | Swine | Rabbits | Poultry | Fur animals |
|------|--------------|------------------|---------|------------|-----------|---------|--------------|-----------|------------|-------------|-------------|
| | heads | | | | | | | | | | |
| 1990 | 2,641,755 | 5,110,397 | 94,500 | 8,739,253 | 1,258,962 | 287,847 | 83,853 | 6,949,091 | 14,893,771 | 173,341,562 | 325,121 |
| 1995 | 2,079,783 | 5,189,304 | 148,404 | 10,667,971 | 1,372,937 | 314,778 | 37,844 | 6,625,890 | 17,110,587 | 184,202,416 | 220,000 |
| 2000 | 2,065,000 | 4,988,000 | 192,000 | 11,089,000 | 1,375,000 | 280,000 | 33,000 | 6,828,000 | 17,873,993 | 176,722,211 | 230,000 |
| 2005 | 1,842,004 | 4,409,921 | 205,093 | 7,954,167 | 945,895 | 278,471 | 30,254 | 7,484,162 | 20,504,282 | 174,667,361 | 200,000 |
| 2010 | 1,746,140 | 4,086,317 | 365,086 | 7,900,016 | 982,918 | 373,324 | 46,475 | 7,588,658 | 17,957,421 | 175,912,339 | 125,000 |
| 2011 | 1,754,981 | 4,142,544 | 354,402 | 7,942,641 | 959,915 | 373,327 | 50,966 | 7,602,093 | 17,549,225 | 174,787,108 | 160,000 |
| 2012 | 1,857,004 | 3,885,606 | 348,861 | 7,015,729 | 891,604 | 395,913 | 59,865 | 7,254,621 | 17,465,477 | 174,647,830 | 165,000 |
| 2013 | 1,862,127 | 3,984,545 | 402,659 | 7,181,828 | 975,858 | 393,915 | 63,166 | 7,111,607 | 16,548,690 | 176,919,056 | 170,000 |
| 2014 | 1,830,990 | 3,925,080 | 369,349 | 7,166,020 | 937,029 | 390,886 | 67,016 | 7,269,295 | 16,435,598 | 175,563,904 | 175,000 |
| 2015 | 1,826,484 | 3,954,864 | 374,458 | 7,148,534 | 961,676 | 384,767 | 70,872 | 7,266,945 | 15,760,502 | 177,391,671 | 180,000 |
| 2016 | 1,821,764 | 4,108,003 | 385,121 | 7,284,874 | 1,026,263 | 388,324 | 74,215 | 7,102,896 | 15,207,274 | 178,690,367 | 160,000 |
| 2017 | 1,791,120 | 4,158,273 | 400,792 | 7,215,433 | 992,177 | 367,561 | 72,455 | 7,185,630 | 14,000,931 | 178,635,180 | 180,000 |
| 2018 | 1,693,332 | 4,229,872 | 401,337 | 7,179,158 | 986,255 | 367,561 | 72,455 | 7,085,003 | 12,089,836 | 175,021,627 | 145,000 |
| 2019 | 1,643,117 | 4,331,830 | 402,286 | 7,000,880 | 1,058,720 | 367,561 | 72,455 | 7,098,664 | 11,755,922 | 175,520,313 | 145,000 |
| 2020 | 1,638,382 | 4,354,633 | 407,027 | 7,034,164 | 1,065,712 | 367,561 | 72,455 | 7,118,709 | 11,010,203 | 178,906,532 | 145,000 |

As for poultry, since no annual statistics on the number of animals are available, the following methodology was followed. For 1990 the ISTAT data from the Agricultural Census have been used; for the years 1991-1999, the number of heads was estimated on the basis of the annual decreases/increases in the production of heads and meat supplied by UNA (National Union of Poultry, which later became UNAITALIA); for 2000 and 2010 ISTAT data from the Censuses of Agriculture for laying hens and broilers were used; for the period 2001-2009 and since 2011 data on the number of broilers and laying hens have been updated, as described below; for the other poultry category, since 1998 the data have been estimated on the basis of UNAITALIA data. Data on turkeys derive from the ISTAT statistics on the Census and the FSS survey.

As stated, data on the number of broilers and laying hens in the period 2001-2009 and since 2011 have been updated. The estimation methodology involved successive steps. Firstly, ISTAT data from the Census and SPA surveys (available for the years 2000, 2005, 2007, 2010, 2013 and 2016) were taken into account; on the basis of these data the number of heads was estimated for the missing years from 2001, assuming a linear trend. The second step involved estimating the number of animals since 2001 on the basis of production data provided by UNAITALIA. The annual variation in production was multiplied by the number of animals in the 2000 Census. The third step was to calculate the average of the two time series calculated in the previous steps.

In Table 6.2 the nitrogen content of N-fertilisers by type applied to soils is shown together with the differentiated EFs. Detailed figures for “other nitrogenous fertilizers” are reported from 1998 because disaggregated official statistics from ISTAT were available only from that year (ENEA, 2006).

Table 6.2 *Time series of N content by fertilisers and relevant emission factors*

| Type of fertilizers | Emission factor | | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|-----------------|---------------------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| | % of N applied | g NH ₃ /kg N applied | | | | | | | | | | | |
| Nitrogen content (t N yr ⁻¹) | | | | | | | | | | | | | |
| Ammonium sulphate | 7.58% | 92 | 50,762 | 61,059 | 36,698 | 27,855 | 32,568 | 16,986 | 18,064 | 16,174 | 16,624 | 10,267 | 17,830 |
| Calcium cyanamide | 1.32% | 16 | 3,310 | 507 | 3,003 | 2,357 | 4,958 | 3,046 | 2,803 | 2,958 | 2,696 | 2,655 | 2,845 |
| Nitrate (*) | 1.27% | 15.41 | 46,657 | 52,769 | 48,701 | 58,427 | 32,964 | 40,157 | 29,247 | 27,869 | 32,477 | 22,332 | 28,097 |
| CAN | 0.66% | 8 | 112,565 | 139,253 | 112,541 | 109,445 | 71,261 | 51,200 | 54,081 | 51,848 | 55,804 | 44,102 | 55,339 |
| Urea | 13.09% | 159 | 291,581 | 321,196 | 329,496 | 317,814 | 209,829 | 266,154 | 321,594 | 261,767 | 241,209 | 247,199 | 295,134 |
| Other nitric nitrogen | 1.32% | 16 | - | - | 3,204 | 5,219 | 3,332 | 1,189 | 1,513 | 1,001 | 1,221 | 1,784 | 2,155 |
| Other ammoniacal nitrogen | 1.32% | 16 | - | - | 6,278 | 18,069 | 12,412 | 7,035 | 8,423 | 6,868 | 7,460 | 7,335 | 9,036 |
| Other amidic nitrogenous | 13.09% | 159 | - | - | 6,988 | 17,420 | 15,366 | 11,796 | 18,246 | 19,944 | 17,982 | 24,331 | 29,366 |
| Phosphate nitrogen | 5.52% | 67 | 112,237 | 99,468 | 77,916 | 69,758 | 45,837 | 35,054 | 33,240 | 42,937 | 35,555 | 34,648 | 44,533 |
| Potassium nitrogen | 1.81% | 22 | 3,937 | 2,876 | 5,291 | 12,289 | 15,955 | 9,077 | 13,361 | 10,503 | 10,751 | 7,922 | 11,353 |
| NPK nitrogen | 5.52% | 67 | 138,018 | 101,528 | 113,897 | 106,384 | 64,462 | 50,174 | 49,829 | 47,416 | 45,749 | 36,686 | 45,730 |

| Type of fertilizers | Emission factor | | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---------------------|-----------------|---------------------------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| | % of N applied | g NH ₃ /kg N applied | | | | | | | | | | | |
| Organic mineral | 1.32% | 16 | 444 | 20,960 | 38,688 | 34,809 | 19,085 | 25,986 | 20,385 | 33,555 | 27,477 | 27,581 | 36,033 |
| Total | | | 759,510 | 799,614 | 782,701 | 779,846 | 528,029 | 517,854 | 570,786 | 522,840 | 495,005 | 466,842 | 577,451 |

(*) includes ammonium nitrate < 27% and ammonium nitrate > 27% and calcium nitrate. The emission factor g NH₃/kg N applied is a weighted average of factors 16 and 14 g NH₃/kg N applied of ammonium nitrate and calcium nitrate.

6.2 METHODOLOGICAL ISSUES

Methodologies used for estimating national emissions from this sector are based on and conform to the *EMEP/EEA Guidebook* (EMEP/EEA, 2019), the *2006 IPCC Guidelines* (IPCC, 1997; IPCC, 2006) and the *IPCC Good Practice Guidance* (IPCC, 2000). Consistency among methodologies for the preparation of the agricultural emission inventory under the UNFCCC and UNECE/CLRTAP is guaranteed through an operational synergy for activity data collection, inventory preparation and reporting to international conventions and European Directives. Information reported in the *National Inventory Report/Common Reporting Format (NIR/CRF)* for the GHG inventory is coherent and consistent with information reported in the *Informative Inventory Report/Nomenclature for Reporting (IIR/NFR)*.

Factor 1.214 (= 17/14) was used to convert ammonia nitrogen to ammonia and factor 3.286 (= 46/14) was used to convert nitrous nitrogen to nitrogen dioxide.

6.2.1 Manure management (3B)

For 3B category, Italy has estimated emissions for pollutants recommended in the *2019 EMEP/EEA Guidebook* (NH₃, NO_x, NMVOC, PM10 and PM2.5). A detailed and updated description of the methodologies for the estimation of NH₃ emissions, as well as of national specific circumstances and reference material, is provided in sectoral reports (APAT, 2005; C3ndor *et al.*, 2008; C3ndor, 2011), and in the NIR (ISPRA, several years [a]). Detailed information on activity data sources, methods and EFs by pollutant for 3B category is shown in Table 6.3.

Table 6.3 Activity data sources, methods and emission factors by pollutant for manure management

| NFR code | Animal category | Method | Activity data | Emission Factor |
|--------------------------------|--|--|---------------|--|
| 3B1a, 3B1b | Cattle | T2 (NH ₃ , NO _x , NMVOC), T1 (PM10, PM2.5) | NS | CS (NH ₃ , NO _x), D (PM10, PM2.5), T2 (NMVOC) |
| 3B4a, 3B2, 3B4d, 3B4e, 3B4f | Buffalo, Sheep, Goats, Horses, Mules and Asses | T2 (NH ₃ , NO _x , NMVOC), T1 (PM10, PM2.5) | NS, IS | CS (NH ₃ , NO _x), D (PM10, PM2.5), T2 (NMVOC) |
| 3B3 | Swine | T2 (NH ₃ , NO _x , NMVOC), T1 (PM10, PM2.5) | NS | CS (NH ₃ , NO _x), D (PM10, PM2.5), T2 (NMVOC) |
| 3B4gi, 3B4gii, 3B4giii, 3B4giv | Poultry | T2 (NH ₃ , NO _x , NMVOC), T1 (PM10, PM2.5) | AS | CS (NH ₃ , NO _x), D (PM10, PM2.5), T2 (NMVOC) |
| 3B4h | Other | T2 (NH ₃ , NO _x , NMVOC), T1 (PM10, PM2.5) | NS | CS (NH ₃ , NO _x), D (PM10, PM2.5), T2 (NMVOC) |

NS=national statistics; IS= International statistics (FAO); AS= category association statistics (UNAITALIA); CS=country-specific; D=Default (from EMEP/EEA Guidebook)

Concerning the 3B category, the estimation procedure for NH₃ emissions consists in successive subtractions from the quantification of nitrogen excreted annually for each livestock category. This quantity can be divided in two different fluxes, depending on whether animals are inside (housing, storage and manure application) or outside the stable (grazing). More in detail, part of the nitrogen excreted in housing volatilizes during the settle of manure in the local farming and it is calculated with the relevant emission factor in housing for the different livestock; this amount is therefore subtracted from the total nitrogen excreted to derive the amount of nitrogen

for storage. During storage another fraction of nitrogen is lost (calculated with the relevant emission factor for storage), which is then subtracted to obtain the amount of nitrogen available for the agronomic spreading. Losses occurring during the spreading are finally calculated with the specific emission factor for spreading. For the nitrogen excreted in the pasture losses due to volatilization calculated with the relevant emission factor for grazing by livestock only occur at this stage (CRPA, 2006[a]).

The manure application source is reported in 3Da2a *Animal manure applied to soils* and the animal grazing source is reported in 3Da3 *Urine and dung deposited by grazing animals*.

As regards the animal grazing, the percentage of grazing animals is equal to (CRPA, 1997, CRPA, 2006[a]): 5% for dairy cattle, 2.2% for other cattle, 2.9% for buffalo, 60% for equines, 90% for sheep and goats.

The excretion rates (CRPA, 2018; CRPA, 2006[a]; GU, 2006; Xiccato *et al.*, 2005), slurry/solid manure production and average weights (CRPA, 2018; CRPA, 2006[a]; GU, 2006; Regione Emilia Romagna, 2004) were updated with country specific information. Other improvements of country specific EFs were obtained with research studies (CRPA, 2010[b]; CRPA, 2006 [a], [b]; CRPA, 2000). Average weight and N excretion rate for NH₃ estimations are reported in Table 6.4.

Table 6.4 Average weight and nitrogen excretion rates from livestock categories in 2020

| Category | Weight kg | Housing | Grazing | Total |
|------------------|--------------|--|---------|---------------|
| | | kg N head ⁻¹ yr ⁻¹ | | |
| Non-dairy cattle | 386.9 | 50.62 | 1.42 | 52.04 |
| Dairy cattle | 602.7 | 103.61 | 5.45 | 109.06 |
| Buffalo | 497.2 | 57.49 | 1.72 | 59.21 |
| Other swine (*) | 88.1 | 13.45 | - | 13.45 |
| Sow (*) | 172.1 | 27.66 | - | 27.66 |
| Sheep | 47.1 | 1.62 | 14.58 | 16.20 |
| Goats | 45.1 | 1.62 | 14.58 | 16.20 |
| Horses | 550.0 | 20.00 | 30.00 | 50.00 |
| Mules and asses | 300.0 | 20.00 | 30.00 | 50.00 |
| Poultry | 1.9 | 0.49 | - | 0.49 |
| Rabbit | 1.6 | 1.02 | - | 1.02 |
| Fur animals | 1.0 | 4.10 | - | 4.10 |

(*) Other swine and sows are sources that represent the 'swine' category

6.2.1.1 Dairy cattle (3B1a)

Following the update of the gross energy intake (GE), based on the estimation of the parameters digestibility (DE) of diet and methane conversion factor (Ym), the excreted nitrogen value of dairy cows was updated for the whole time series. Excreted nitrogen is in fact calculated from GE using equations 10.31-10.33 of the 2006 IPCC Guidelines. In addition, the percentage for protein in diet has been updated from the previous submission. This parameter is used with GE in the estimation of excreted nitrogen.

As regards the DE e Ym parameters, the estimation methodology, set out in the 2019 IPCC Guidelines, allows, from data on average annual milk production per cow and per production level (low "<5000 kg/head/year", medium "5000-8500 kg/head/year" and high ">8500 kg/head/year") and information on animal diets, to calculate an average value of DE and Ym for dairy cows. On the basis of data from the Italian Livestock Breeders' Association (AIA) on average annual milk production and the number of dairy cows in production, by region and breed, the distribution of animals was calculated according to the three productivity levels identified by the 2019 IPCC Guidelines, for the years 2004-2019. The AIA carries out milk productivity checks on behalf of the Ministry of Agriculture and each year the sample of animals checked is about 50% of the number of animals reared. The difference in cow numbers between the AIA total and the ISTAT total (used for emission estimates) was attributed to the low production level. The DE values assigned to the three production levels (low, medium, high) are 62, 65 and 70.11 respectively and were identified in collaboration with the CRPA dairy cow feeding experts. The value 62 is the minimum value of the range indicated in Table 10.12 for low producing cows. The value 65 is lower than the average value of the range indicated in Table 10.12 for medium producing cows. The value 70.11 for high-producing cows is a weighted average of two values: the first is 65 (corresponding to diets with DE \geq 70 and NDF \geq 35) and was attributed to 27% of the high-producing cows fed without silage fodder; the second is 72 (corresponding to diets with DE \geq 70 and NDF \leq 35) and was attributed to 73% (=100-27%) of the high-producing cows fed silage fodder. With reference to the 27% of cows, this value includes cows whose milk is intended for the production of Parmigiano Reggiano (17% of total cows), and cows fed with good quality dry and green fodder (e.g. for the production of

Trentingrana PDO (Protected Designation of Origin), Latte Fieno STG (Traditional Speciality Guaranteed) and other mountain cheeses; 10% of total cows). In support of the choices made for high productivity values, mention is made of a study published in 2020 (Gislon et al, 2020) carried out on eight Italian Friesian cows in multiparous lactation, with high productivity, using a 4 × 4 replicated Latin square pattern. The experimental design of the square involves all cows receiving all diets (with adaptation periods between each), so we have 2 groups of 4 cows that rotated 4 times on as many diets. The number of observations for each diet is 32. The cow effect is nullified because they all receive all diets and the results obtained are therefore irrefutable and highly representative, according to CRPA experts. Four diets, based on the following forages (% of dry matter, DM; neutral detergent fiber content, NDF, expressed as % of DM), were tested: corn silage (CS, 49.3; 32.8 NDF), alfalfa silage (AS, 26.8; 27.1 NDF), wheat silage (WS, 20.0; 33.7 NDF), and a typical hay-based Parmigiano Reggiano cheese production diet (PR, 25.3 of both alfalfa and Italian ryegrass hay; 36.7 NDF). The lowest DM digestibility was observed for the PR diet (64.5%) and the highest for the CS diet (73.3%); AS and WS diets showed intermediate values (71.4 and 70.3% respectively). PR diet is associated with diets with DE \geq 70 and NDF \geq 35 in table 10.12 of the 2019 IPCC Guidelines and the other three diets are associated with diets with DE \geq 70 and NDF \leq 35 in the same table. For the year 2020, the percentages of dairy cows according to the three productivity levels are 78.9%, 12.3% and 8.8%. The digestibility values associated with these productivity levels are, as mentioned earlier, 70.11%, 65% and 62% respectively. With these data, the average digestibility value of the diets consumed by dairy cows was calculated as 68.77%. The weighted average value of Ym for the year 2020 is 5.98% of gross energy intake. This value was calculated from the percentage distribution of dairy cows according to the three productivity levels and using the default factors given in Table 10.12 of the 2019 IPCC Guidelines. These values are: for lactating phase, 6.5 for low producing cows, 6.3 for medium producing cows, 6.0 and 5.7 for high producing cows; for dry phase, 7 for low producing cows, 6.3 for medium and high producing cows. From the two values for high producing cows in lactating phase, a weighted average value of 5.78 was calculated for the year 2020, with the distribution of cows according to diet type, shown above: 27% of high-productivity cows are associated with diets with DE \geq 70 and NDF \geq 35 (Ym 6.0); 73% of high-productivity cows are associated with diets with DE \geq 70 and NDF \leq 35 (Ym 5.7). As regards the percentage for protein in diet, mentioned above, on the basis of data from around 500 samples of rations (unifeed) of lactating and dry dairy cows, analysed by the CRPA's zootechnical feed service for the three-year period 2017-2019, from all over Italy, the crude protein of the ration was updated. The data were obtained by weighing the values expressed as % of the dry matter of the ration with the average annual lactation period (equal to 305 days) and the dry period (equal to 60 days). The value obtained, 14.22, was used for the time series from 2010 onwards, as indicated by the CRPA experts. For the previous years, the previous figure of 15.32 was left until 2000, and an average value of 14.5 was used for the intermediate years between 2000 and 2010.

In Table 6.5 the animal waste management system (AWMS) distribution and EFs used are reported. EF was multiplied by the percentage of the nitrogen excreted in housing equal to 95% of the total, assuming that 5% is excreted in grazing. The value is a weighted average based on country specific emission factors and the distribution of livestock housing has been assumed in the following main housing systems reported in Table 6.5 (based on a 1998 CRPA survey carried out in Lombardy, Emilia Romagna and the centre of Italy and on ISTAT statistics of 2003 and on 2010 Agricultural Census). Between 2005 and 2010 a gradual transition to the updated distribution of housing systems has been assumed for the intermediate years taking in account the gradual penetration of systems to ensure animal welfare.

Table 6.5 AWMS distribution and EF by manure management system for the dairy cattle category

| Emission factors by manure management system | 1990 | 2003 | 2005 | 2010 | 2013 |
|--|-------------|-------------|-------------|-------------|-------------|
| Housing | | | | | |
| cubicle house: 14.3 N-NH ₃ kg/head/year (Bonazzi et al, 2005) | 14.6% | 14.6% | 14.6% | 27.9% | 27.9% |
| loose housing on bedding: 15.7 N-NH ₃ kg/head/year (Bonazzi et al, 2005) | 9.2% | 9.2% | 9.2% | 42.6% | 42.6% |
| tied cows: 12.9 N-NH ₃ kg/head/year (Bonazzi et al, 2005) | 76.2% | 76.2% | 76.2% | 29.5% | 29.5% |
| EF N-NH₃ kg/head/year | 13.4 | 13.4 | 13.4 | 14.5 | 14.5 |
| Storage | | | | | |

| | liquid manure = 36% | liquid manure = 36% | liquid manure = 36% | liquid manure = 48% | liquid manure = 48% |
|---|------------------------|------------------------|------------------------|------------------------|------------------------|
| liquid manure | | | | | |
| Tanks (for liquid manure): 23% of N at storage (Bonazzi et al, 2005) | 40.0% | 75.5% | 75.5% | 82.3% | 70.1% |
| Lagoons (for liquid manure): 32.2% (multiplication factor equal to 1.4 respect to tanks) | 50.0% | 12.5% | 12.5% | 2.5% | 1.7% |
| covered storage (for liquid manure): | 10.0% | 12.5% | 12.5% | 15.2% | 28.3% |
| covered tanks high reduction: 4.6% (reduction of 80% compared to tanks) | | 1.0% | 1.0% | 3.0% | 4.0% |
| covered tanks medium reduction: 9.2% (reduction of 60% compared to tanks) | | 1.0% | 1.0% | 3.0% | 4.0% |
| covered tanks low reduction: 13.8% (reduction of 40% compared to tanks) | 10.0% | 10.5% | 10.5% | 8.9% | 16.2% |
| biogas: no emission | | | | 0.33%(1) | 4.1%(1) |
| solid storage: 14.2% of N at storage (Regione Emilia Romagna, 2001) | solid manure = 64% | solid manure = 64% | solid manure = 64% | solid manure = 52% | solid manure = 52% |

(1) Data were calculated from the results of the CRPA study on assessing the emission effects of livestock processing (CRPA, 2018). The study estimated that in 2020 18% of cattle manure would be sent to anaerobic digestion and 48% of digestate tanks would be covered. Based on these results, it was assumed that: in 2010, 3% of manure went to digesters and 11% were covered digestate tanks; in 2013 these values become 15% and 27% respectively. This trend was assumed on the basis of the exponential growth in the last ten years of anaerobic digesters.

As regards the manure storage (see Table 6.5), emission factors are expressed as a percentage of the nitrogen contained in manure to storage. Emission factors used for tanks is derived from national literature (Bonazzi et al, 2005) and emission factors for lagoons and covered storage have been estimated applying an increase (for lagoons) and a reduction (for covered storage) to tanks EF (as referenced in CRPA, 2006[a] and CRPA, 2006[b]).

The proportion of liquid system (considering liquid system= liquid system + digesters) and solid storage (considering solid storage= solid storage + digesters), reported in the CRF (*Common Reporting Format* for the GHG inventory) refer to the nitrogen excreted and not to the amount of animal waste. The proportion reported in the Table 6.5 refer to the manure production according to the type of housing.

EFs for lagoons and covered storage have been provided by CRPA (CRPA, 2006[a]). For lagoons, they have a high exposure area relative to their capacity and represent a higher emission type than the tank. Considering the volumes of the two types of storage, an increase in the surface of slurry in the lagoons with respect to the tanks can be estimated equal to 40%. Since ammonia emissions are estimated to be proportional to the surface of slurry exposed to air, emissions from lagoons will be approximately 40% higher than those of the tanks (CRPA, 1997). For covered storage, the emission reduction has been assumed on the basis of the ILF-BREF document (EC, 2003) and the Ammonia Guidance Document (Bittman S. et al, 2014), as reported in CRPA (CRPA, 2006[a]; 2018).

A linear emission reduction in the period 1990-2003 has been estimated to assess the dynamics of evolution of storage systems from the values available in 1990 and 2003, as reported by CRPA (CRPA, 2006[a]). In 2003 respect to 1990 an increase of storage in tanks with respect to lagoons as well as a small increase of covered storage is observed as available in the Table 6.5. On the basis of ISTAT statistics on storage systems as 2010 Agricultural Census and 2013 Farm Structure Survey, an update of emission factors from manure storage for cattle category has been estimated. A gradual transition to the updated emission factors has been assumed for the intermediate years (for the period 2005-2010 and 2010-2013) taking in account the gradual penetration of the abatement technologies.

On the basis of the study for the evaluation of the effects on emissions of livestock management practices carried out by CRPA for the emission scenarios for 2020 and 2030 (CRPA, 2018), NH₃ emissions from storage for cattle have been modified considering the average distribution of the covered tanks related to the different ammonia emission reduction efficiencies.

EFs for manure storage reported in the Table 6.5 have been multiplied by the percentage of nitrogen remaining after housing emissions and the result has been multiplied by the nitrogen excreted in housing to obtain emissions from storage. Emissions have been divided by total heads to obtain the EF kg/head reported in the Table 6.8 for the year 2020.

Regarding emission factors for cattle, the evolution of different abatement technologies along the period is considered in the EFs used for NH₃ estimation for housing, storage and land spreading systems. Improvements in the abatement technologies are based on the results of both the IIASA questionnaire for the implementation

of RAINS scenarios in 2003 and an *ad hoc* survey conduct in the 2005 by CRPA (CRPA, 2006 [a], [b]) and on ISTAT statistics such as 2010 Agricultural Census, 2013 and 2016 Farm Structure Survey.

6.2.1.2 Swine (3B3)

Activity data of swine population (3B3) reported in the IIR/NFR are different from data reported in the NIR/CRF. In fact, piglets (swine less than 20 kg) are included in the swine population in the NIR/CRF for the estimation of CH₄ emission from enteric fermentation, while they are not included in the number of the NFR templates because the NH₃ EF used for sows takes into account the emissions from piglets, thus ensuring the comparability of the implied emission factors. For NH₃ estimations average weighted emission factors for each category (other swine and sows) are calculated taking in account the relevant emission factors of the abatement technologies for each manure system. The implemented abatement technologies for the years 1990, 2003 and 2005 are reported in Table 6.6.

Table 6.6 Abatement technologies for the swine category

| Livestock category | 1990 | 2003 | 2005 | 2010 | 2013 |
|--|--|--|--|--|---|
| Housing | | | | | |
| fattening swine | 55% Partly-slatted floor (PSF); 20% Fully-slatted floor (FSF); 25% solid floor | 55% PSF; 25% FSF; 20% solid floor | 26% FSF; 39% PSF; 12% FSF + vacuum system (VS); 4% FSF + with flush canals; 7% FSF + with flush tubes; 5% PSF + VS; 6% PSF + with flush canals; 1% PSF + with flush tubes | Same distribution for the year 2005 | Same distribution for the year 2005 |
| gestating sows (75% of the total sows) | 65% FSF; 35% PSF | 50% FSF; 50% PSF | 26% FSF; 52% PSF; 5% FSF + vacuum system (VS); 5% FSF + with flush canals; 7% FSF + with flush tubes; 2% PSF + VS; 2% PSF + with flush canals; 1% PSF + with flush tubes | Same distribution for the year 2005 | Same distribution for the year 2005 |
| lactating sows (25% of the total sows) | 75% FSF+ deep collection pit; 25% sloping floor | 65% FSF+ deep collection pit; 35% sloping floor | 52% FSF + deep collection pit; 39% sloping floor; 3% with flush; 6% mechanical removal | Same distribution for the year 2005 | Same distribution for the year 2005 |
| weaners 6-20 kg | 80% FSF + deep collection pit; 20% sloping floor | 70% FSF+ deep collection pit; 30% sloping floor | 63% FSF + deep collection pit; 14% sloping floor; 7% FSF + VS; 11% FSF with flush tubes; 2% FSF + scraper; 2% PSF + VS; 1% PSF + deep collection pit | Same distribution for the year 2005 | Same distribution for the year 2005 |
| Livestock category | | | | | |
| Storage | | | | | |
| swine | 61% lagoons; 36% tanks; 3% covered storage: | 54% lagoons; 43% tanks; 3% covered storage: | 46% lagoons; 51% tanks; 3% covered storage: | 10% lagoons; 79% tanks; 11% covered storage: | 7% lagoons; 67% tanks; 25% covered storage: |
| covered tanks high reduction: reduction of 80% compared to tanks | | | | 1% | 3% |
| covered tanks | | 1% | 1% | 4% | 5% |

| Livestock category | 1990 | 2003 | 2005 | 2010 | 2013 |
|--|------|------|------|----------|----------|
| medium reduction: 60% compared to tanks | | | | | |
| covered tanks low reduction: 40% compared to tanks | 3% | 2% | 2% | 6% | 17% |
| biogas: no emission | | | | 0.33%(1) | 0.81%(1) |

| Livestock category | 1990 | 2003 | 2005 | 2010 | 2013 | 2016 |
|-----------------------|------|---|---|--|--|--|
| Land spreading | | | | | | |
| swine | 100% | 80% broadcasting 10% low efficiency 10% high efficiency | 78% broadcasting 11% low efficiency 11% high efficiency | 70% broadcasting 17% low efficiency 6% medium efficiency 7% high efficiency | 48% broadcasting 30% low efficiency 12% medium efficiency 11% high efficiency | 36% broadcasting 27% low efficiency 26% medium efficiency 11% high efficiency |

(1) Data were calculated from the results of the CRPA study on assessing the emission effects of livestock processing (CRPA, 2018). The study estimated that in 2020 3% of swine manure would be sent to anaerobic digestion and 48% of digestate tanks would be covered. Based on these results, it was assumed that: in 2010 and 2013 the percentage of manure sent to digesters remains at 3%, while the percentage of covered digestate tanks changes, becoming 11% and 27% respectively.

Regarding emission factors for swine, the evolution of different abatement technologies along the period is considered in the EFs used for NH₃ estimation for housing, storage and land spreading systems. Improvements in the abatement technologies are based on the results of both the IIASA questionnaire for the implementation of RAINS scenarios in 2003 and an *ad hoc* survey conduct in the 2005 by CRPA (CRPA, 2006 [a], [b]). Furthermore, an update of emission factors from manure storage and land spreading for swine category has been estimated on the basis of ISTAT statistics on manure storage systems and land spreading techniques such as 2010 Agricultural Census, 2013 and 2016 Farm Structure Survey. A gradual transition to the updated emission factors has been assumed for the intermediate years (for the period 2005-2010, 2010-2013 and 2014-2016) taking in account the gradual penetration of the abatement technologies.

On the basis of the study for the evaluation of the effects on emissions of livestock management practices carried out by CRPA for the emission scenarios for 2020 and 2030 (CRPA, 2018), NH₃ emissions from storage for swine have been modified considering the average distribution of the covered tanks related to the different ammonia emission reduction efficiencies.

6.2.1.3 Poultry (3B4g)

As regards 3B4gi (laying hens) and 3B4gii (Broilers) categories, NH₃ emissions show different trends. The different trend for the laying hens is due to the evolution of different abatement technologies along the period, that are considered in the EFs used for NH₃ estimation for housing, storage and land spreading systems. Emission factors used for each of the different abatement technologies for laying hens (as referenced in CRPA, 2006[a] and CRPA, 2006[b]) are reported in Table 6.7.

Table 6.7 AWMS distribution, abatement technologies and EF by manure management system for the laying hens category

| Emission factors by manure management system | 1990 | 2003 | 2005 | 2010 |
|---|------|------|------|------|
| Housing | | | | |
| open manure storage under cages (for liquid manure) (RS) = 0.220 kg NH ₃ /head/year (EC, 2003) | 100% | 20% | 11% | 4% |

| Emission factors by manure management system | 1990 | 2003 | 2005 | 2010 |
|--|-------------|-------------|-------------|-------------|
| deep pit = 0.162 kg NH ₃ /head/year (ENEA, 2003) | | 24% | | |
| vertical tiered cages with manure belts and forced air drying = 0.06 kg NH ₃ /head/year (ENEA, 2003) [reduction in ammonia emissions of 73% compared to RS] | | 56% | 74% | 50% |
| vertical tiered cages with manure belt and whisk-forced air drying = 0.088 kg NH ₃ /head/year (EC, 2003) [reduction in ammonia emissions of 60% compared to RS] | | | 2% | |
| aerated open manure storage (deep-pit or high rise systems and canal house) = 0.154 kg NH ₃ /head/year (EC, 2003) [reduction in ammonia emissions of 30% compared to RS] | | | 10% | 11% |
| vertical tiered cages with manure belt and drying tunnel over the cages = 0.044 kg NH ₃ /head/year (EC, 2003) [reduction in ammonia emissions of 80% compared to RS] | | | 3% | |
| Loose housing with outdoor access (RS) = 0.3 kg NH ₃ /head/year (Bittman S. et al, 2014) | | | | 7% |
| Loose housing without outdoor access = 0.18 kg NH ₃ /head/year (Bittman S. et al, 2014; our assumptions) | | | | 28% |
| Storage | | | | |
| liquid manure = 16% (percentage of nitrogen to storage) (Nicholson et al, 2004) | 100% | 20% | 11% | 4% |
| solid manure = 7.3% (ENEA, 2003) | | 80% | 89% | 96% |
| Land spreading | | | | |
| liquid manure = 37.1% of TAN applied (TAN/TKN = 35%) (CRPA, 2006[a]) [broadcasting] | 100% | 5% | 9% | 9% |
| low efficiency = 7.8% (bandspreading and incorporation within 6 hours for liquid manure) [reduction of 40% compared to broadcasting] | | 50% | 65% | 65% |
| high efficiency = 2.6% (shallow and deep injection for liquid manure) [reduction of 80% compared to broadcasting] | | 45% | 26% | 26% |
| solid manure = 67% of TAN applied (TAN/TKN = 21%) (Nicholson et al, 2004; CRPA, 2006[a]) [broadcasting] | | 10% | 9% | 10% |
| low efficiency = 11.0% (incorporation within 12-24 hours for solid manure) [reduction of 20% compared to broadcasting] | | 40% | 37% | 43% |
| high efficiency = 2.8% (incorporation within 4 hours for solid manure) [reduction of 80% compared to broadcasting] | | 50% | 54% | 46% |

Emission factors used for each of the different techniques for housing are derived from ILF BREF of IPPC (EC, 2003) and a study at national level on ammonia emissions from laying hens (ENEA, 2003). In 2010, on the basis of the housing distribution collected from the 2010 Agricultural Census and emission factors and abatement systems data reported in the Guidance from the UNECE Task Force on Reactive Nitrogen (Bittman S. et al, 2014) emission factors have been updated. Between 2005 and 2010 a gradual transition to the updated distribution of housing systems has been assumed for the intermediate years taking in account the gradual penetration of systems to ensure animal welfare.

As regards the manure storage, emission factors are expressed as a percentage of the nitrogen contained in manure to storage. Emission factors used for liquid manure is derived from Nicholson et al (Nicholson et al, 2004) and emission factors for solid manure is from ENEA (ENEA, 2003). On the basis of the 2010 Agricultural Census conducted by ISTAT, an update of emission factors from manure storage for laying hens category has been estimated. A gradual transition to the updated emission factors has been assumed for the intermediate years (for the period 2005-2010) taking in account the gradual penetration of the abatement technologies. EFs for manure storage reported in Table 6.7 have been multiplied by the amount of nitrogen remaining after housing emissions.

For land spreading, emissions have been estimated by CRPA (CRPA, 2006[a] and CRPA, 2006[b]). As regards the liquid manure, the amount of N-NH₄ emissions, in percentage of the applied ammoniacal nitrogen, have been assumed equal to those of the cattle slurry due to the lack of data (CRPA, 2006[a]). As regards the solid manure, the amount of N-NH₄ emissions, in percentage of the applied ammoniacal nitrogen, were equal to 67% (Nicholson et al, 2004; CRPA, 2006[a]). In 2003 and 2005 the evolution of different improvements technologies based on the results of both the IIASA questionnaire for the implementation of RAINS scenarios and a survey conduct by CRPA, has been implemented in the EFs used. For the period 1900-2003, a linear emission reduction has been estimated and applied. The efficiency of reduction techniques has been estimated

on the basis of the UNECE document Control techniques for preventing and abating emissions of ammonia (as referenced in CRPA, 2006[a] and CRPA, 2006[b]). EFs for land spreading reported in Table 6.7 have been multiplied by the amount of nitrogen remaining after storage emissions.

As regards broilers, only a slight improvement on spreading system has occurred. From 1995 a chicken-dung drying process system has been introduced for laying hens and improved along the period.

As recommends by the 2019 and 2020 NECD review (EEA, 2019; EEA, 2020), emissions of NO_x, NH₃, PM and NMVOC from turkeys have been estimated and reported in category 3B4giii.

Regarding emission factors for poultry, the evolution of different abatement technologies along the period is considered in the EFs used for NH₃ estimation for housing, storage and land spreading systems. Improvements in the abatement technologies are based on the results of both the IIASA questionnaire for the implementation of RAINS scenarios in 2003 and an *ad hoc* survey conduct in the 2005 by CRPA (CRPA, 2006 [a], [b]) and on ISTAT statistics such as 2010 Agricultural Census, 2013 and 2016 Farm Structure Survey.

Average emission factors for NH₃ per head are reported in Table 6.8.

Table 6.8 NH₃ emission factors for manure management for the year 2020

| Category | Housing | Storage | Land spreading <i>kg NH₃ head⁻¹ yr⁻¹</i> | Grazing | Total |
|------------------|---------|---------|--|---------|--------------|
| Non-dairy cattle | 8.17 | 6.96 | 5.98 | 0.14 | 21.25 |
| Dairy cattle | 16.73 | 13.70 | 11.55 | 0.53 | 42.50 |
| Buffalo | 9.28 | 9.91 | 7.79 | 0.17 | 27.15 |
| Other swine (*) | 2.38 | 1.62 | 1.12 | | 5.12 |
| Sow (*) | 4.86 | 3.33 | 2.31 | | 10.49 |
| Sheep | 0.22 | 0.28 | 0.32 | 0.71 | 1.53 |
| Goats | 0.22 | 0.25 | 0.33 | 0.71 | 1.50 |
| Horses | 3.21 | 4.43 | 1.97 | 2.91 | 12.51 |
| Mules and asses | 3.21 | 4.43 | 1.97 | 2.91 | 12.51 |
| Laying hens | 0.13 | 0.05 | 0.04 | | 0.22 |
| Broilers | 0.08 | 0.05 | 0.02 | | 0.15 |
| Turkeys | 0.25 | 0.15 | 0.09 | | 0.49 |
| Other poultry | 0.09 | 0.05 | 0.02 | | 0.16 |
| Rabbit | 0.34 | 0.13 | 0.06 | | 0.54 |
| Fur animals | 1.37 | | 0.30 | | 1.67 |

(*) Other swine and sows are sources that represent the 'swine' category

NH₃ emissions from digesters biogas facilities (in particular due to different phases of the process: during storage of feedstock on the premises of the biogas facility, during the liquid–solid separation of the digestate, during storage of the digestate) have been estimated on the basis of the amount of nitrogen in manure feeding anaerobic digesters and the tier 1 emission factor derived by the EMEP/EEA Guidebook (EMEP/EEA, 2019). On the basis of CRPA data on measurements of nitrogen quantities in livestock manure (downstream of releases to housing and storage) per animal category and type of manure, the nitrogen quantities in livestock manure sent to anaerobic digestion were estimated. The coefficients, expressed as g N/kg manure, were calculated gross of losses and then the losses to housing were deducted. The resulting coefficients were then multiplied by the quantities of manure sent for anaerobic digestion. The whole time series was updated. NH₃ emissions from digesters biogas facilities have been subtracted from manure management category (for cattle, swine and poultry categories) and allocated in the anaerobic digestion at biogas facilities (5B2 of the waste sector). As requested during the 2019 and 2020 NECD reviews (EEA, 2019; EEA, 2020), the amount of total feedstock, livestock manure and nitrogen in manure treated by biogas facilities are shown in Table 6.9.

Table 6.9 Total feedstock, animal manure and nitrogen in manure treated by biogas facilities

| Year | Total feedstock (t) | Animal manure in total feedstock (t) | Nitrogen in animal manure (kg) |
|-------------|---------------------|--------------------------------------|--------------------------------|
| 1990 | - | - | - |
| 1995 | n.a. | 273,179 | 1,440,550 |
| 2000 | n.a. | 165,256 | 870,566 |
| 2005 | n.a. | 1,075,851 | 5,405,336 |
| 2006 | n.a. | 1,910,684 | 10,022,720 |
| 2007 | 6,395,717 | 2,214,577 | 11,695,580 |

| Year | Total feedstock (t) | Animal manure in total feedstock (t) | Nitrogen in animal manure (kg) |
|------|---------------------|--------------------------------------|--------------------------------|
| 2008 | 5,848,368 | 1,887,038 | 9,861,437 |
| 2009 | 4,634,551 | 1,684,000 | 8,728,895 |
| 2010 | 6,496,988 | 1,761,932 | 8,580,530 |
| 2011 | 12,566,873 | 3,836,365 | 19,974,670 |
| 2012 | 15,428,001 | 7,062,440 | 37,695,458 |
| 2013 | 27,655,508 | 12,704,785 | 68,647,391 |
| 2014 | 30,856,283 | 14,166,069 | 76,571,624 |
| 2015 | 29,477,553 | 13,566,441 | 73,188,472 |
| 2016 | 29,637,431 | 13,638,212 | 73,518,651 |
| 2017 | 30,301,334 | 13,944,626 | 75,172,834 |
| 2018 | 30,235,418 | 13,915,377 | 75,055,782 |
| 2019 | 32,446,567 | 14,891,895 | 80,410,913 |
| 2020 | 32,708,097 | 15,031,251 | 81,151,281 |

n.a. = not available

Because of multiple substrates fed to bio-digesters, the following average characteristics of the feedstock, reported in Table 6.10, as supplied by CRPA, are considered for the Italian bio-digesters in order to calculate the total amount of feed from animal manure anaerobic digestion (CRPA, 2018).

Table 6.10 Percentages of different substrates for anaerobic digestion feedstock

| Type of feed | Units | animal manure | energy crops | agro-industrial by-products |
|--|---------------|---------------|--------------|-----------------------------|
| Animal manure only | % in the feed | 100 | 0 | 0 |
| Animal manure + energy crops + agro-industrial by-products | % in the feed | 28 | 52 | 20 |
| Animal manure + energy crops | % in the feed | 38 | 62 | 0 |
| Animal manure + agro-industrial by-products | % in the feed | 69 | 0 | 31 |
| Energy crops + agro-industrial by-products | % in the feed | 0 | 81 | 19 |

Source: CRPA

On the basis of the information reported above and in consideration of the typical feed of the bio-digesters the average parameters for animal manure, energy crops and agro-industrial by-products are those reported in Table 6.11. The biogas methane content is generally reported to range from 50% to 65%, for the inventory purposes and according to CRPA methane content is assumed to be 55%. As regards the average volatile solids content, values for animal manure and agro-industrial by-products have been changed based on the recent study of CRPA (CRPA, 2018).

Table 6.11 Average parameters of different substrates for anaerobic digestion feedstock

| Parameters | Units | animal manure | energy crops | agro-industrial by-products |
|------------------------------------|-----------------------------|---------------|--------------|-----------------------------|
| Average biogas producing potential | m ³ biogas/kg VS | 0.4 | 0.6 | 0.6 |
| Average CH ₄ content | % | 55 | 55 | 55 |
| Average volatile solids content | kg/t feed | 139 | 280 | 237 |

Source: CRPA

For further information on the method of estimating the quantity of manure sent to digesters and the amount of nitrogen stored in digesters, see the information and data reported in the NIR (see paragraphs 5.3.2 *Methodological issues* in chapter 5 and A7.2 Manure management (3B) in annex 7).

The percentage of nitrogen lost through N-NH₃ emissions from anaerobic digesters was subtracted from the percentage of nitrogen left after emissions during housing and storage, reducing the amount of nitrogen used at the spreading. The amount of nitrogen used at the spreading also includes the digestate.

For NO_x emissions (during storage) tier 2 method reported in the EMEP/EEA Guidebook (EMEP/EEA, 2019) was used for calculations. EFs by livestock category and manure type derived from the EMEP/EEA Guidebook (EMEP/EEA, 2019) are based on nitrogen mass-flow approach built from country specific data on nitrogen excretion and solid/liquid distribution of manure.

For NMVOC emissions a tier 2 method was used for calculations. Tier 2 NMVOC EFs are those reported in the EMEP/EEA Guidebook (EMEP/EEA, 2019). However, the emission factors are very high and therefore more detailed analyzes will be carried out. As requested during the 2020 NECD reviews (EEA, 2020), NMVOC emissions from turkeys are estimated and reported in the 3B4giii category.

For particulate matter emissions a tier 1 method was used for calculations. EFs for PM10 and PM2.5 are derived from the EMEP/EEA Guidebook (EMEP/EEA, 2019; EMEP/CORINAIR, 2006), modified on the basis of the Italian animal breeding characteristics and weight parameters (Córdoba *et al.*, 2008; Córdoba, 2011). For swine and poultry, emission factors have been updated from 2010, estimating a gradual transition to the updated emission factors from 2005, reflecting changes in manure management systems recorded by ISTAT surveys (FSS and the agricultural census). From 2010 PM emission estimates are based on emission factors provided by the 2019 EMEP/EEA Guidebook. These emission factors are based on studies conducted between 2006 and 2016 which include scientific works conducted in Italy. These studies have suggested that Takai's emission factors suggested in the 2006 EMEP/CORINAIR Guidebook are too high and do not represent current particulate emission levels. A gradual transition to the updated emission factors has been assumed for the intermediate years (2004-2010) taking in account the gradual penetration of the abatement technologies. As requested during the 2020 NECD reviews (EEA, 2020), PM emissions from turkeys are estimated and reported in the 3B4giii category.

PM emissions from turkeys, sheep, goats, mules and asses and fur animals are also estimated. Average emission factors for PM per head are reported in Table 6.12.

Table 6.12 PM emission factors for manure management for the year 2020

| Category | PM10 | PM2.5 |
|------------------|--|-------|
| | <i>kg PM head⁻¹ yr⁻¹</i> | |
| Non-dairy cattle | 0.318 | 0.210 |
| Dairy cattle | 0.657 | 0.428 |
| Buffalo | 0.502 | 0.328 |
| Other swine (*) | 0.190 | 0.008 |
| Sow (*) | 0.230 | 0.012 |
| Sheep | 0.053 | 0.016 |
| Goats | 0.051 | 0.015 |
| Horses | 0.242 | 0.154 |
| Mules and asses | 0.137 | 0.086 |
| Laying hens | 0.033 | 0.002 |
| Broilers | 0.024 | 0.002 |
| Turkeys | 0.109 | 0.020 |
| Other poultry | 0.050 | 0.008 |
| Rabbit | - | - |
| Fur animals | 0.003 | 0.002 |

(*) Other swine and sows are sources that represent the 'swine' category

6.3 AGRICULTURAL SOILS (3D)

For agricultural soils, estimations of NH₃ emissions account for the direct application of synthetic N-fertilizers (3Da1), animal manure applied to soils (3Da2a), sewage sludge applied to soils (3Da2b), other organic fertilisers applied to soil (3Da2c), animal grazing (3Da3) and N fixed by leguminous crops (3De). For the same sources, emissions of NO_x were estimated (except for 3De *Cultivated crops*). *Crop residues applied to soils* (3Da4) and *Indirect emissions from managed soils* (3Db) emissions have not been estimated as in the guidelines there is insufficient information. PM10 and PM2.5 emissions from the Farm-level agricultural operations including storage, handling and transport of agricultural products have been estimated and reported in 3Dc category. NMVOC emissions from animal manure applied to soils, animal grazing and cultivated crops have been estimated and reported in 3Da2a, 3Da3 and 3De categories respectively. HCB emissions from the use of pesticides have been estimated and reported in 3Df category.

NH₃ emissions from synthetic N-fertilizer (3Da1) are based on the guidebook methodology (EMEP/EEA, 2019), which provides different EFs by type of fertilizers taking into account climatic conditions and pH of the soil (EFs in Table 6.2). A tier 2 method has been implemented for 3Da1 source. NH₃ emissions from synthetic N-fertilizers are obtained with the amount of the N content by type of fertilizer multiplied by the specific EFs. Emissions have been calculated on the basis of EFs for temperate climate and normal pH factors

according to the IPCC climate zones classification and the definition available in the 2002 EMEP/CORINAIR Guidebook for which Italy is defined with large areas of acidic soils (soil pH below 7.0) and with some calcareous soils (or managed with soil pH above 7.0).

In 2011 a validation of EFs and estimations was carried out considering the results of a research study that estimated, at NUTS 2 level, emissions for the use of synthetic N-fertilizers considering type of cultivation, altitude, and climatic conditions (CRPA, 2010[b]; C ndor and Valli, 2011).

Based on the comparison with ASSOFERTILIZZANTI and ISTAT experts on the time series of synthetic fertiliser use, the nitrate data in the years 2009-2011 were revised for the current submission. In addition, nitrate data (quantity and nitrogen content) were recalculated to include the estimated CAN fertiliser. The emission factor of NH₃ for the use of CAN from the EMEP/EEA Guidebook (EMEP/EEA, 2019) has been used.

NO_x emission factor for synthetic N-fertilizer is equal to 0.04 kg NO₂/kg fertiliser N applied (EMEP/CORINAIR, 2019).

The method for estimating NH₃ emissions from animal manure applied to soils (3Da2a) is described in 3B (tier 2). On the basis of ISTAT statistics on spreading systems such as 2010 Agricultural Census, 2013 and 2016 Farm Structure Survey, an update of emission factors from land spreading for cattle, swine, laying hens and broilers categories have been estimated. A gradual transition to the updated emission factors has been assumed for the intermediate years (for the period 2005-2010, 2010-2013 and 2013-2016) taking in account the gradual penetration of the abatement technologies. For NO_x emissions (during spreading) a tier 2 method was used for calculations. EFs by livestock category and manure type derived from the EMEP/EEA Guidebook (EMEP/EEA, 2019) are based on nitrogen mass-flow approach. For NMVOC emissions a tier 2 method was used for calculations. Tier 2 NMVOC EFs are those reported in the EMEP/EEA Guidebook (EMEP/EEA, 2019).

Concerning the sludge spreading (3Da2b), the total production of sludge from urban wastewater plants, as well as the total amount of sludge used in agriculture and some parameters such as N content, are communicated from 1995 by the Ministry for the Environment, Land and Sea from 1995 (MATTM, several years[a]) in the framework of the reporting commitments fixed by the European Sewage Sludge Directive (EC, 1986) transposed into the national Legislative Decree 27 January 1992, n. 99. From 1990 to 1994 activity data and parameters were reconstructed, as reported in detail in the Chapter 7 of the National Inventory Report on the Italian greenhouse gas inventory (ISPRA, several years [a]).

The amount of sewage N applied was calculated using the amount of sewage sludge (expressed in tons of dry matter) and the N content of sludge. The dry matter contained in sludge at national level is assumed to be 25% of total sludge. In Table 6.13, the total amount of sewage sludge production as well as sludge used in agriculture and nitrogen content in sludge is reported. The default NH₃ EF (0.13 kg NH₃/kg N applied) and NO_x EF (0.04 kg NO₂/kg N applied) are from EMEP/EEA Guidebook (EMEP/EEA, 2019).

Table 6.13 *Sludge spreading activity data and parameters, 1990 – 2020*

| Year | Sewage sludge production (t) | Sewage sludge used in agriculture (t) | Sewage sludge used in agriculture (t of dry matter) | N concentration in sludge (% dry matter) | Total N in sludge (t) |
|-------------|-------------------------------------|--|--|---|------------------------------|
| 1990 | 3,272,148 | 392,658 | 98,164 | 5.2 | 5,071 |
| 1995 | 2,437,024 | 630,046 | 157,512 | 5.2 | 8,137 |
| 2000 | 3,402,017 | 869,696 | 217,424 | 5.0 | 10,954 |
| 2005 | 4,298,576 | 862,970 | 215,742 | 4.1 | 8,874 |
| 2010 | 3,697,625 | 992,859 | 248,215 | 4.0 | 10,040 |
| 2011 | 3,572,092 | 1,196,634 | 299,159 | 3.7 | 11,119 |
| 2012 | 3,446,558 | 1,096,380 | 274,095 | 4.7 | 12,864 |
| 2013 | 3,074,108 | 814,178 | 203,545 | 4.0 | 8,053 |
| 2014 | 3,154,060 | 804,623 | 201,156 | 4.1 | 8,301 |
| 2015 | 3,069,302 | 888,899 | 222,225 | 3.7 | 8,303 |
| 2016 | 3,183,919 | 765,639 | 191,410 | 3.9 | 7,410 |
| 2017 | 3,183,641 | 691,375 | 172,844 | 4.1 | 7,116 |
| 2018 | 3,137,372 | 693,434 | 173,358 | 4.2 | 7,247 |
| 2019 | 3,254,796 | 673,670 | 168,417 | 4.1 | 6,977 |

| Year | Sewage sludge production (t) | Sewage sludge used in agriculture (t) | Sewage sludge used in agriculture (t of dry matter) | N concentration in sludge (% dry matter) | Total N in sludge (t) |
|------|------------------------------|---------------------------------------|---|--|-----------------------|
| 2020 | 3,224,910 | 651,972 | 162,993 | 4.1 | 6,759 |

As regards the other organic fertilisers applied to soil (3Da2c) category, the use of other organic N fertilisers, including compost and organic amendments, and N content are provided by ISTAT (as reported in the paragraph 6.1). The default NH₃ EF (0.08 kg NH₃/kg waste N applied) and NO_x EF (0.04 kg NO₂/kg N waste applied) are from EMEP/EEA Guidebook (EMEP/EEA, 2019).

For 3Da3 the time series of the quantity of N from animal grazing is the same as that reported in the NIR and in the relevant CRF tables. The method for estimating NH₃ emissions is described in 3B (tier 2). The default NO_x EF is from EMEP/EEA Guidebook (EMEP/EEA, 2019). For NMVOC emissions a tier 2 method was used for calculations. Tier 2 NMVOC EFs are those reported in the EMEP/EEA Guidebook (EMEP/EEA, 2019).

Nitrogen input from N-fixing crops (3De) has been estimated starting from data on surface and production for N-fixing crops and forage legumes; nitrogen input from N-fixing crops (kg N yr⁻¹) is calculated with a country-specific methodology. Peculiarities that are present in Italy were considered: N-fixing crops and legumes forage. Nitrogen input is calculated with two parameters: cultivated surface and nitrogen fixed per hectare (Erdamn 1959 in Giardini, 1983). Emissions are calculated using the default emission factor 1 kg N-NH₃/ha (EMEP/CORINAIR, 2006). In Table 6.14, cultivated surface from N-fixing species (ha yr⁻¹) and N fixed by each species (kg N ha⁻¹ yr⁻¹) are shown.

NMVOC emissions from cultivated crops have been estimated and reported in 3De category. The method (tier 1) for estimating NMVOC emissions from cultivated crops (3De) is described in 3D chapter of the EMEP/EEA Guidebook (EMEP/EEA, 2019). The default NMVOC EF is from EMEP/EEA Guidebook (EMEP/EEA, 2019). Hectares of wheat, rape, rye crops surface and total grass surface were considered as activity data according to the methodology EMEP/EEA Guidebook (EMEP/EEA, 2019).

Table 6.14 Cultivated surface (ha) and nitrogen fixed by each variety (kg N ha⁻¹ yr⁻¹)

| N fixed | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2018 | 2019 | 2020 | |
|--|------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|------------------|
| kg N ha ⁻¹ yr ⁻¹ | | | | | ha | | | | | |
| Bean, f.s. | 40 | 29,096 | 23,943 | 23,448 | 23,146 | 19,027 | 17,059 | 18,368 | 18,253 | 17,915 |
| Bean, d.s. | 40 | 23,002 | 14,462 | 11,046 | 8,755 | 7,001 | 5,870 | 6,411 | 5,587 | 5,541 |
| Broad bean, f.s. | 40 | 16,564 | 14,180 | 11,998 | 9,484 | 8,487 | 7,914 | 7,985 | 7,624 | 7,372 |
| Broad bean, d.s. | 40 | 104,045 | 63,257 | 47,841 | 48,507 | 52,108 | 42,157 | 50,421 | 60,007 | 61,982 |
| Pea, f.s. | 50 | 28,192 | 21,582 | 11,403 | 11,636 | 8,691 | 14,940 | 15,559 | 16,197 | 16,154 |
| Pea, d.s. | 72 | 10,127 | 6,625 | 4,498 | 11,134 | 11,692 | 11,181 | 17,916 | 22,926 | 20,766 |
| Chickpea | | | | | | | | | | |
| a | 40 | 4,624 | 3,023 | 3,996 | 5,256 | 6,813 | 11,167 | 26,024 | 20,999 | 18,579 |
| Lentil | 40 | 1,048 | 1,038 | 1,016 | 1,786 | 2,458 | 3,099 | 5,417 | 5,861 | 5,612 |
| Vetch | 80 | 5,768 | 6,532 | 6,800 | 7,142 | 7,560 | 7,795 | 7,827 | 7,827 | 7,827 |
| Lupin | 40 | 3,303 | 3,070 | 3,300 | 2,500 | 3,401 | 3,358 | 3,337 | 3,337 | 3,337 |
| Soya | | | | | | | | | | |
| bean | 58 | 521,169 | 195,191 | 256,647 | 152,331 | 159,511 | 308,979 | 326,587 | 273,332 | 256,134 |
| Alfalfa | 194 | 987,000 | 823,834 | 810,866 | 779,430 | 745,128 | 667,325 | 695,492 | 719,073 | 715,642 |
| Clover | | | | | | | | | | |
| grass | 103 | 224,087 | 125,009 | 114,844 | 103,677 | 102,691 | 119,942 | 124,375 | 127,087 | 127,270 |
| Total | | 1,958,025 | 1,301,746 | 1,307,702 | 1,164,784 | 1,134,567 | 1,220,786 | 1,305,719 | 1,288,110 | 1,264,131 |

(*) f.s.=fresh seed; d.s.=dry seed

PM10 and PM2.5 emissions from the Farm-level agricultural operations including storage, handling and transport of agricultural products have been estimated and reported in 3Dc category. The method (tier 1) for estimating PM10 and PM2.5 emissions is described in 3D chapter of the EMEP/EEA Guidebook (EMEP/EEA, 2019). The default PM10 and PM2.5 EFs are from EMEP/EEA Guidebook (EMEP/EEA, 2019). Hectares of total arable crop surface have been used as activity data for PM emissions according to the methodology EMEP/EEA Guidebook (EMEP/EEA, 2019).

HCB emissions from the use of pesticides (3Df) have been estimated. HCB emissions result from the use of HCB as pesticide but also by the use of other pesticides which contain HCB as an impurity. For the period 1996-2001, data are from the database of pesticides contained in the National agricultural information system

(*Sistema informativo agricolo nazionale - SIAN*³). For the period 2002-2008, SIAN data have been elaborated by Provincial Agency for the Protection of the Environment of the Autonomous Province of Trento⁴. From 2009 activity data have been processed by the Service for risks and environmental sustainability of technologies, chemical substances, production cycles and water services and for inspection activities of ISPRA on the basis of data provided by ISTAT related to substances chlorothalonil, picloram, lindane and chlortal-dimetile which are the active ingredients of pesticides containing HCB.

The availability of data allows estimating emissions from pesticides where HCB is found as an impurity, as in lindane, DCPA, chlorothalonil and Picloram. Emissions from the use of HCB as a pesticide were not estimated. As result from the 2020 NEC review, the HCB emissions from the use of pesticides have been revised. On the basis of the amount of HCB contained in these pesticides (lindane: 0.005%; DCPA: 0.004%; chlorothalonil: 0.004%; Picloram: 0.005%) and according to the EMEP/EEA Guidebook (EMEP/EEA, 2019), which states all the HCB present as a contaminant will be volatilised, HCB emissions result in 118.82 kg for 1990 and 2.40 kg in 2020. An international research work at European level (Berdowski et al., 1997) estimated 400 kg of HCB emissions from pesticide use for Italy in 1990 while in the last years these emissions should be null.

Detailed information on activity data sources, methods and EFs by pollutant for 3D category is shown in Table 6.15.

Table 6.15 Activity data sources, methods and emission factors by pollutant for agriculture soils

| NFR code | Category | Method | Activity data | Emission Factor |
|----------|---|--|---------------|---|
| 3Da1 | Inorganic N-fertilizers (includes also urea application) | T2 (NH ₃), T1 (NO _x) | NS | T2 (NH ₃), D (NO _x) |
| 3Da2a | Animal manure applied to soils | T2 (NH ₃ , NO _x , NMVOC) | NS | CS (NH ₃), D (NO _x), T2 (NMVOC) |
| 3Da2b | Sewage sludge applied to soils | T1 (NH ₃ , NO _x) | NS | D (NH ₃ , NO _x) |
| 3Da2c | Other organic fertilisers applied to soils (including compost) | T1 (NH ₃ , NO _x) | NS | D (NH ₃ , NO _x) |
| 3Da3 | Urine and dung deposited by grazing animals | T2 (NH ₃ , NO _x , NMVOC) | NS | CS (NH ₃), D (NO _x), T2 (NMVOC) |
| 3Da4 | Crop residues applied to soils | | | |
| 3Db | Indirect emissions from managed soils | | | |
| 3Dc | Farm-level agricultural operations including storage, handling and transport of agricultural products | T1 (PM10, PM2.5) | NS | D (PM10, PM2.5) |
| 3Dd | Off-farm storage, handling and transport of bulk agricultural products | | | |
| 3De | Cultivated crops | CS (NH ₃), T1 (NMVOC) | NS | D (NH ₃ , NMVOC) |
| 3Df | Use of pesticides | T1 (HCB) | NS | D (HCB) |

6.4 FIELD BURNING OF AGRICULTURAL RESIDUES (3F)

NMVOC, CO, NO_x, NH₃, SO₂, PM10, PM2.5, BC, As, Cr, Cu, Ni, Se, Zn, Pb, Cd, Hg, Dioxin and PAH emissions have been estimated, applying the tier 1 and tier 2 (for heavy metals, PAH emissions and BC) approach. A detailed description of the methodology and parameters used is shown in the NIR (ISPRA, several years [a]). The same methodology to estimate emissions from open burning of waste, as reported in paragraph 7.2 of the waste section (see *Small scale waste burning (5C2)* subparagraph), is used on the basis of the amount of fixed residues instead of removable residues. Concerning NO_x, CO, NMVOC, IPCC emission factors have been used (IPCC, 1997), while for PM10 and PM2.5 emission factors from the USEPA (EPA, 1995) and BC

³ <http://www.sian.it/portale-sian/attivaserivizio.jsp?sid=174&pid=6&servizio=Banca+Dati+Fitofarmaci&bottoni=no>

⁴ http://www.appa.provincia.tn.it/fitofarmaci/programmazione_dei_controlli_ambientali/-Criteri_vendita_prodotti_fitosanitari/pagina55.html

emission factors from the EMEP/EEA Guidebook (EMEP/EEA, 2019) have been applied. NH₃, SO₂, heavy metals, dioxin and PAH emission factors are from the EMEP/EEA Guidebook (EMEP/EEA, 2019).

As concerns NO_x and CO emission factors, values used are in the range of the tier 1 emission factors from the EMEP/EEA Guidebook (EMEP/EEA, 2019).

As concerns PM, emission factors derived from the USEPA 1995 without differences between PM₁₀ and PM_{2.5} are higher than those (both tier 1 and tier 2) reported in the EMEP/EEA Guidebook (EMEP/EEA, 2019). The update of the emission factors based on the emission factors available in the 2019 EMEP/EEA Guidebook (GB) have been evaluated. The GB EFs refer to the same period and to the same country (Jenkins et al, 1996 - USA) of the EFs used in the inventory (USEPA 1995); furthermore, there is not much difference between the PM₁₀ and PM_{2.5} EFs taken from the GB. Considering also the following factors, i.e.: the uncertainty of the GB EFs; the probable error related to the EFs for the rice (the emission factor reported in the GB is very different from that in the reference publication of Jenkins); the tier 1 emission factors from the GB are similar but not equal to the average of the values relating to four types of crops reported in the reference scientific publication mentioned in the GB (Jenkins, 1996a); the low significance of the considered emission category, we decided to maintain the current EFs, pending more appropriate updates.

6.5 TIME SERIES AND KEY CATEGORIES

The following sections present an outline of the main key categories in the agriculture sector.

The agriculture sector is the main source of NH₃ emissions in Italy; for the main pollutants, in 2020 the sector accounts for:

- 95.1% of national total NH₃ emissions
- 18.5% of national total HCB emissions
- 14.0% of national total NMVOC emissions
- 14.0% of national total PM₁₀ emissions
- 9.6% of national total NO_x emissions
- 4.0% of national total PM_{2.5} emissions
- 3.1% of national total Cd emissions

Moreover, the sector comprises 0.7% of BC emissions, 0.6% of CO, 0.6% of PAH, 0.4% of Hg, 0.21% of As, 0.15% of Se, 0.09% of SO₂, 0.08% of Cr, 0.04% of Dioxins, 0.04% of Ni, 0.03% of Zn and 0.01% of Pb, 0.01% of Cu. There are no particular differences as compared to the sectoral share in 1990 when the agriculture sector accounted for 96.3% of NH₃ emissions, 11.1% of PM₁₀, 3.1% of PM_{2.5}, except for NMVOC emissions (7.4%), NO_x emissions (2.9%) and HCB emissions where agriculture accounted for 83.4% of total national emissions.

Table 6.16 reports the key categories identified in the agriculture sector while the time series of NH₃ emissions by sources is shown in Table 6.17.

Concerning NH₃ emissions, the category *manure management* (3B) represents in 2020 54.0% of national total ammonia emissions (58.0% in 1990). In particular, NH₃ emissions from *cattle* (3B1) stand for 59.2% of 3B emissions, while emissions from *swine* (3B3) and *poultry* (3B4g) represent 15.8% and 14.5%, respectively. The category *agricultural soils* (3D) represents in 2020 41.0% of national total ammonia emissions (38.2% in 1990). The animal manure applied to soils (3Da2a) and the use of synthetic N-fertilisers (3Da1) represent 44.9% and 41.1% of 3D emissions, respectively.

Regarding PM₁₀ emissions, the category *manure management* (3B) accounts for 6.6% in 2020 (6.2% in 1990) of national total PM₁₀ emissions. *Poultry* (3B4g), *cattle* (3B1) and *swine* (3B3) represent the major contributors to the total PM₁₀ emissions from category 3B with 58.0%, 22.6% and 12.6%, respectively. The category Farm-level agricultural operations including storage, handling and transport of agricultural products (3Dc) accounts for 6.2% in 2020 (4.2% in 1990) of national total PM₁₀ emissions. For PM_{2.5} emissions, the category *manure management* (3B) contributes for 2.1% in 2020 (1.9% in 1990) of national total PM_{2.5} emissions. *Cattle* (3B1) accounts for 58.0%, while *poultry* (3B4g) stands for 28.1% to the total PM_{2.5} emissions from category 3B. The category Farm-level agricultural operations including storage, handling and transport of agricultural products (3Dc) accounts for 0.3% in 2020 (0.2% in 1990) of national total PM_{2.5} emissions.

Concerning NO_x emissions, the category *manure management* (3B) represents in 2020 0.29% of national total NO_x emissions (0.09% in 1990). For NO_x emissions, the category *agricultural soils* (3D) contributes for 9.3% in 2020 (2.8% in 1990) of national total NO_x emissions. *Inorganic N-fertilizers* (3Da1) and *Animal manure applied to soils* (3Da2a) account for 43.6% and 35.5% of total 3D emissions, respectively.

Concerning NMVOC emissions, the category *manure management (3B)* represents in 2020 11.4% of national total NMVOC emissions (6.9% in 1990). For NMVOC emissions, the category *agricultural soils (3D)* contributes for 2.6% in 2020 (1.4% in 1990) of national total NMVOC emissions. For NMVOC emissions, the category *manure management (3B)* and *agricultural soils (3D)* contributes for 81.2% and 18.3% in 2020 of agricultural NMVOC emissions. *Cattle (3B1)*, *poultry (3B4g)* and *buffalo (3B4a)* represent the major contributors to the total NMVOC emissions from category 3B with 73.5%, 15.6% and 5.0%, respectively. Most of the emissions in the 3D category (69.3%) derive from *Animal manure applied to soils (3Da2a)*.

Table 6.16 Key categories in the agriculture sector in 2020

| | SO _x | NO _x | NH ₃ | NMVOC | CO | PM10 | PM2.5 | BC | Pb | Cd | Hg | PAH | DIOX | HCB | PCB |
|--------|-----------------|-----------------|-----------------|-------|------|------|-------|------|------|------|------|------|------|-------|-----|
| % | | | | | | | | | | | | | | | |
| 3B1a | | 0.06 | 13.75 | 4.20 | | 0.65 | 0.53 | | | | | | | | |
| 3B1b | | 0.09 | 18.17 | 4.18 | | 0.83 | 0.69 | | | | | | | | |
| 3B2 | | 0.02 | 0.96 | 0.10 | | 0.22 | 0.08 | | | | | | | | |
| 3B3 | | 0.00 | 8.52 | 0.35 | | 0.83 | 0.04 | | | | | | | | |
| 3B4a | | 0.01 | 2.15 | 0.57 | | 0.12 | 0.10 | | | | | | | | |
| 3B4d | | 0.00 | 0.14 | 0.01 | | 0.03 | 0.01 | | | | | | | | |
| 3B4e | | 0.01 | 0.77 | 0.11 | | 0.05 | 0.04 | | | | | | | | |
| 3B4f | | 0.00 | 0.15 | 0.01 | | 0.01 | 0.00 | | | | | | | | |
| 3B4gi | | 0.02 | 1.91 | 0.26 | | 0.77 | 0.07 | | | | | | | | |
| 3B4gii | | 0.03 | 3.68 | 0.63 | | 1.53 | 0.19 | | | | | | | | |
| 3B4gii | | 0.01 | 1.42 | 0.34 | | 0.85 | 0.19 | | | | | | | | |
| 3B4gii | | 0.01 | 0.83 | 0.55 | | 0.65 | 0.13 | | | | | | | | |
| 3B4h | | 0.01 | 1.49 | 0.11 | | 0.00 | 0.00 | | | | | | | | |
| 3Da1 | | 4.05 | 16.88 | | | | | | | | | | | | |
| 3Da2a | | 3.29 | 18.43 | 1.78 | | | | | | | | | | | |
| 3Da2b | | 0.05 | 0.24 | | | | | | | | | | | | |
| 3Da2c | | 0.86 | 2.71 | | | | | | | | | | | | |
| 3Da3 | | 1.03 | 2.36 | 0.02 | | | | | | | | | | | |
| 3Dc | | | | | | 6.16 | 0.30 | | | | | | | | |
| 3De | | | 0.42 | 0.78 | | | | | | | | | | | |
| 3Df | | | | | | | | | | | | | | 18.51 | |
| 3F | 0.09 | 0.08 | 0.14 | 0.07 | 0.65 | 1.30 | 1.62 | 0.73 | 0.01 | 3.08 | 0.36 | 0.60 | 0.04 | | |

Note: key categories are shaded in blue

Table 6.17 Time series of ammonia emissions in agriculture (Gg)

| NFR SECTOR 3 | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 3B1a Manure management - Dairy cattle | 93.01 | 75.27 | 70.57 | 62.05 | 59.71 | 54.34 | 54.41 | 53.80 | 51.06 | 49.25 | 49.85 |
| 3B1b Manure management - Non-dairy cattle | 88.77 | 86.57 | 84.55 | 73.57 | 71.01 | 61.11 | 64.04 | 64.47 | 66.00 | 66.47 | 65.91 |
| 3B2 Manure management - Sheep | 4.34 | 5.30 | 5.51 | 3.95 | 3.92 | 3.55 | 3.62 | 3.58 | 3.56 | 3.48 | 3.49 |
| 3B3 Manure management - Swine | 36.65 | 34.97 | 35.53 | 36.74 | 34.10 | 31.63 | 30.98 | 31.28 | 30.78 | 30.83 | 30.89 |
| 3B4a Manure management - Buffalo | 3.06 | 4.53 | 5.79 | 6.40 | 10.34 | 7.88 | 7.71 | 7.99 | 7.95 | 7.81 | 7.81 |
| 3B4d Manure management - Goats | 0.58 | 0.63 | 0.63 | 0.44 | 0.45 | 0.44 | 0.47 | 0.46 | 0.46 | 0.49 | 0.49 |
| 3B4e Manure management - Horses | 2.20 | 2.40 | 2.14 | 2.13 | 2.85 | 2.94 | 2.96 | 2.81 | 2.81 | 2.81 | 2.81 |

| NFR SECTOR 3 | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| 3B4f Manure management - Mules and asses | 0.64 | 0.29 | 0.25 | 0.23 | 0.35 | 0.54 | 0.57 | 0.55 | 0.55 | 0.55 | 0.55 |
| 3B4gi Manure management - Laying hens | 13.87 | 12.68 | 9.56 | 6.30 | 7.84 | 7.04 | 6.99 | 7.09 | 6.89 | 6.95 | 6.93 |
| 3B4gii Manure management - Broilers | 12.37 | 12.96 | 12.17 | 11.77 | 12.00 | 12.85 | 12.84 | 13.07 | 12.84 | 12.91 | 13.36 |
| 3B4giii Manure management - Turkeys | 7.30 | 7.24 | 7.18 | 5.42 | 6.08 | 5.80 | 5.46 | 5.10 | 4.96 | 4.96 | 5.16 |
| 3B4giv Manure management - Other poultry | 1.76 | 4.06 | 3.15 | 4.62 | 2.27 | 2.97 | 3.54 | 3.32 | 3.30 | 3.20 | 3.00 |
| 3B4h Manure management - Other animals (*) | 7.51 | 8.41 | 8.79 | 10.00 | 8.69 | 7.72 | 7.43 | 6.89 | 5.93 | 5.77 | 5.42 |
| 3Da1 Inorganic N-fertilizers (includes also urea application) | 69.58 | 72.51 | 72.33 | 70.69 | 48.27 | 53.30 | 62.97 | 54.12 | 49.99 | 50.40 | 61.21 |
| 3Da2a Animal manure applied to soils | 95.21 | 86.91 | 79.05 | 70.19 | 68.11 | 66.08 | 66.27 | 66.68 | 66.05 | 66.18 | 66.83 |
| 3Da2b Sewage sludge applied to soils | 0.66 | 1.06 | 1.42 | 1.15 | 1.31 | 1.08 | 0.96 | 0.93 | 0.94 | 0.91 | 0.88 |
| 3Da2c Other organic fertilisers applied to soils (including compost) | 1.32 | 1.45 | 1.81 | 1.78 | 3.29 | 4.68 | 4.55 | 7.02 | 5.64 | 6.84 | 9.82 |
| 3Da3 Urine and dung deposited by grazing animals | 10.09 | 11.33 | 11.48 | 8.74 | 8.97 | 8.60 | 8.79 | 8.65 | 8.60 | 8.51 | 8.55 |
| 3De Cultivated crops | 2.38 | 1.58 | 1.59 | 1.41 | 1.38 | 1.48 | 1.50 | 1.55 | 1.59 | 1.56 | 1.54 |
| 3F Field burning of agricultural residues | 0.49 | 0.48 | 0.48 | 0.52 | 0.50 | 0.51 | 0.55 | 0.49 | 0.49 | 0.49 | 0.49 |
| Total | 451.74 | 430.63 | 413.98 | 378.10 | 351.46 | 334.55 | 346.60 | 339.84 | 330.38 | 330.38 | 344.99 |

Note: (*) 3B4h includes rabbits and fur animals

The largest and most intensive agricultural area in Italy is the Po River catchment with the following characteristics: high crop yields due to climatic factors, double cropping system adopted by livestock farms, flooded rice fields, high livestock density and animal production that keep animals in stables all the year (Bassanino et al 2011, Bechini and Castoldi 2009). 64%, 78% and 88% of cattle, poultry and swine production are located in Piedmont, Lombardy, Emilia-Romagna, and Veneto Regions (Northern Italy/Po River Basin). At regional level, the presence of large cattle, poultry and swine farms in the Po basin assume a particular relevance for air quality issues, especially, for the specific meteorological conditions of this area.

The reduction of NH₃ emissions from 3B is mainly related to the reduction in the number of animals. Between 1990 and 2020 total NH₃ emissions from 3B have reduced by 23.6%. Cattle livestock decreased by 22.7% (from 7,752,152 to 5,993,015 heads). Dairy cattle and non-dairy cattle have decreased by 38.0% and 14.8%, respectively. The so-called first pillar of the EU Common Agriculture Policy (CAP), dealing with market support, had a strong impact through the milk quota system by reducing animal numbers in the dairy sector to compensate for increasing animal productivity (EEA, 2016). On the contrary, swine and poultry have increased between 1990 and 2020 by 2.4% and 3.2%, respectively (see Table 6.1). Abatement technologies are considered in the EFs used for NH₃ estimations. Research studies funded by ISPRA, such as the MeditAiraneo project, or by the Ministry of Environment have allowed us to collect information on the inclusion of abatement technologies in Italy, especially those related to the swine and poultry recovery and treatment of manure and to land spreading (CRPA, 2006[b]; Córdor et al., 2008; CRPA, 2010[b]).

NH₃ emissions of 3D category are driven by the animal manure applied to soils and the use of inorganic N-fertilizers. Between 1990-2020 emissions have respectively decreased by 29.8% and 12.0% mainly due to the reduction of the number of animals and the use of inorganic N-fertilizers, that are decreased overall by 24.0% (while urea increased by 1.2%), in terms of nitrogen content. According to the Italian Fertilizer Association (AIF, *Associazione Italiana Fertilizzanti*) the use of fertilisers is determined by their cost and particularly by the price of agricultural products. Because of the agriculture product price decreasing, minor amount of fertilisers has been used by farmers to reduce costs (Perelli, 2007). Furthermore, the EU Nitrates Directive which aims at reducing and preventing water pollution caused by nitrates from agricultural sources has addressed the lower use of synthetic and nitrogen-based fertilisers (EEA, 2016).

Every 4 years (first every 5) the national emission inventory is disaggregated at NUTS3 level as requested by CLRTAP (Córdor *et al.*, 2008). A database with the time series for all sectors and pollutants has been published (ISPRA, 2021; ISPRA, 2009; ISPRA, several years [c]; ISPRA, several years [d]). The disaggregation of 2019 agricultural emissions has also been finalised and figures are available at the following web site: <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>. The disaggregation (NUTS3) of the NH₃ agricultural emissions is shown in Figure 6.2. In 2019, four regions contributed with more than 60% of agricultural NH₃ emissions: Lombardia, Veneto, Emilia Romagna and Piemonte.




Figure 6.2 *NH₃ emissions from Agriculture in 2019 (t)*

6.6 QA/QC AND VERIFICATION

QA/QC procedures for the agriculture sector are in line with the 2006 IPCC Guidelines and consistent with the EMEP/EEA Guidebook. Italy has drawn up a QA/QC procedure manual and elaborates annually a QA/QC plan both for the UNFCCC and UNECE/CLRTAP inventories. In the QA/QC Agriculture section GHG and NH₃ emissions improvements are specified (ISPRA, several years [b]). Furthermore, feedbacks for the agricultural emission inventory derive also from communication of data to different institutions (ISTAT, UNAITALIA, CRPA etc.) and/or at local level (regional environmental institutions). In addition, ISPRA participates in a technical working group on agriculture within the National Statistical System, composed by producers and users of agricultural statistics.

Data used to estimate emissions were verified with census data. Slight differences in the livestock number (cattle and other swine) are found between conjunctural surveys (used for emissions estimation) and Agricultural Census for the year 2010; while for the other categories the differences are more significant. In the conjunctural surveys, the number of heads of the sows, sheep, goats, mules and asses, broilers, laying hens categories is on average 15% higher than the census, whereas for other poultry the difference is 30% and for horses and rabbits is more than double.

Ammonia emissions for swine and poultry manure management from housing and storage were compared with data reported in the E-PRTR registry for the year 2014, which represent 62.1% and 19.5%, respectively, of national NH₃ emissions for the same categories (3B).

Data on national sales of synthetic nitrogen fertilizers (by type of fertilizers) as provided by *Assofertilizzanti-Federchimica*⁵ (data are available online - <https://assofertilizzanti.federchimica.it/fertilizzanti/statistiche-fertilizzanti>) for the period 2012-2019 have been compared to official statistics provided by ISTAT. Differences were mainly found for the amount of simple mineral nitrogen fertilizers, where data from *Assofertilizzanti* are higher by 17%, on average, for the years 2013-2019. This could be due to a possible double counting of some product which could be considered as a single product and as a compound with other fertilizers. Further investigations will be conducted.

A check on the urea data has been made. Yara, the only Italian producer, provided an estimate of urea consumption in the various production sectors, such as SCR engines, NO_x emissions reduction, industrial and agricultural uses. All these uses have been considered in the national emissions inventory. Further checks will be made between apparent consumption and end uses.

In 2021, a technical report (CREA, 2021) produced by CREA (Council for Agricultural Research and Economics), in collaboration with ISPRA, on the assessment of emissions related to the use of nitrogen fertilizers was published, which contains several analyses of nitrogen fertilizer consumption data from different data sources: IFASTAT/Fertilizer Europe, FAO, EUROSTAT, *Assofertilizzanti*, National Integrated Production Specifications, Farm Accountancy Data Network (FADN). These analyses suggest further investigation to better understand the differences found between these data sources and the ISTAT data used for the estimates.

In 2021, NH₃ emissions for dairy cows were also calculated with the AgrEE tool - Agricultural Emission Estimation tool (made by the JRC) and differences with national estimates for the year 2015 were analysed. National estimates are slightly higher than the AgrEE tool results (16.5 kt N-NH₃ for spreading vs 16.1 kt N-NH₃) for several reasons. In particular two of them are: for N₂O emissions from storage, according to the Guidebook, it is necessary to remove emissions from housing, while IPCC considers the N excreted at the housing; for NH₃ emissions from spreading, the Guidebook does not remove N lost through leaching and run-off at storage, unlike the IPCC methodology. The verification with AgrEE tool revealed other differences that will be taken into account for improved emission estimates.

6.7 RECALCULATIONS

In 2022, recalculations were implemented for the agricultural emission inventory.

Major changes are: corrected EF NH₃ housing for 2019 for non-dairy cattle and buffaloes categories; included in storage estimates NH₃ emission factors for equine and sheep and goat for the whole time series; modified the formula for estimating NH₃ emissions from storage for cattle, pigs and poultry by subtracting from N at housing also the amount of N at digesters before multiplying by the emission factor; for the estimation of N at spreading, the percentages of N remaining after emissions in the housing, storage and after other losses during storage (NO₂, N₂O, N₂, N leached in manure management, NH₃ from digesters) were recalculated and N bedding was added; updated PM emission factors of the other poultry category for the whole time series; corrected the formula for calculating the biogas produced, which affects the estimate of N at digesters the whole time series for dairy cattle and non-dairy cattle and, to a small extent, from 1991 to 2004 also for pigs; updated the estimate of straw use for dairy cattle, non-dairy cattle and buffaloes to take into account sheltered animals only (removing grazing animals).

Minor changes are: corrected error in total biomass available in 2018 and 2019, implying change in NH₃ field burning emissions; updated equine and sheep and goat housing NH₃ emission factors for the whole time series; updated N excreted dairy cattle for 2019 based on updated protein percentage in milk; updated 2019 rabbit data; corrected 2016 fertiliser data, based on an ISTAT update, that implies a slight change in the years 2017-2019 also in the estimation of N ammonium nitrate and calcium nitrate, whose total (reported by ISTAT) remains unchanged but changed the values of the two fertilizers, which have slightly different FE NH₃; corrected soil improver data for the years 2015-2019, based on an update by ISTAT; updated turkey estimate 2017-2019; change in total arable crops per PM estimate for 2019 due to updated area data for aubergines grown in greenhouses.

⁵ *Federchimica* is the National Association of the Chemical Industry and *Assofertilizzanti* represents the production companies of the fertilizer industry.

Changes in NH₃ emissions from housing and storage imply changes in the estimated NMVOC emissions from storage and spreading. HCB emissions from the use of pesticides have been updated for the year 2019.

6.8 PLANNED IMPROVEMENTS

Currently, uncertainty analysis, for the agricultural emission sector, is carried out only for the GHG emission inventory. We plan to estimate uncertainties also for the other pollutants, including NH₃ and PM. Monte Carlo analysis has also been performed for one key category of the GHG agricultural emission inventory; initial results are shown in the NIR (ISPRA, several years [a]).

No emissions are estimated for 3Da4 *Crop residues applied to soils*, 3Db *Indirect emissions from managed soils* and 3Dd *off-farm storage, handling and transport of bulk agricultural products*. However, Italy will assess the availability of AD and EFs for these categories.

In the coming years, the Permanent census of agriculture will provide valuable information on animal and agronomic production methods. The focus of the Permanent census is to provide a comprehensive information framework on the structure of the agricultural system and the livestock at national, regional and local level by integrating archive data and carrying out statistical support surveys. Statistical registers will be created with the aim of increasing the quantity and quality of information in order to reduce the response.

The results of the AgrEE tool verification will be analysed in order to improve the estimates made.

7 WASTE (NFR SECTOR 5)

7.1 OVERVIEW OF THE SECTOR

Italy estimates the categories of the waste sector, as reported in the following box. In this submission Cd, Hg and Pb emissions from 5E have been estimated too. Conversely, Italy does not consider NH₃ emissions from latrines because this activity does not occur or it can be considered negligible. In the last available national census on wastewater treatment plants (ISTAT, 2015) the following data are reported: 99.4% of people are served by the sewage system, 17,897 wastewater system plants serve a total of 98,360,724 people equivalent; 8,377 are the Imhoff tanks present in Italy, 1,607 are the primary wastewater treatment plants, 5,604 are the secondary wastewater treatment plants and 2,309 are the advanced wastewater treatment plants. The biogas collected from the anaerobic digestion of wastewaters is burned with heat/energy recovery and relevant emissions are reported in Category 1 while emissions from the exceeding biogas which is flared are not estimated at the moment because emission factors are under investigation, but anyway it should be negligible.

| NFR | | SNAP | |
|-------|-------------------------------|----------|--|
| 5A | Solid waste disposal on land | 09 04 01 | Managed waste disposal on land |
| | | 09 04 02 | Unmanaged waste disposal on land |
| 5B | Biological treatment of waste | 09 10 05 | Compost production |
| | | 09 10 06 | Anaerobic digestion at biogas facilities |
| 5C1a | Municipal waste incineration | 09 02 01 | Incineration of municipal wastes |
| 5C1b | Other waste incineration | 09 02 02 | Incineration of industrial wastes |
| | | 09 02 05 | Incineration of sludge from wastewater treatment |
| | | 09 02 07 | Incineration of hospital wastes |
| | | 09 02 08 | Incineration of waste oil |
| 5C1bv | Cremation | 09 09 01 | Cremation of corpses |
| 5C2 | Small scale waste burning | 09 07 00 | Open burning of agricultural wastes |
| 5D | Wastewater handling | 09 10 01 | Waste water treatment in industry |
| | | 09 10 02 | Waste water treatment in residential and commercial sector |
| 5E | Other waste | | Car and building fires |

Concerning air pollutants, emissions estimated for each sector are reported in Table 7.1.

Table 7.1 Air pollutant emissions estimated for each sector

| Main pollutants | 5A | 5B | 5C1a | 5C1bi | 5C1bii | 5C1biii | 5C1biv | 5C2 | 5C1bv | 5D | 5E |
|------------------------------|----|----|------|-------|--------|---------|--------|-----|-------|----|----|
| NO _x | | | x | x | x | x | x | x | x | | |
| CO | | | x | x | x | x | x | x | x | | |
| NMVOG | x | x | x | x | x | x | x | x | x | x | |
| SO _x | | | x | x | x | x | x | x | x | | |
| NH ₃ | x | x | | | | | | | | | |
| Particulate matter | | | | | | | | | | | |
| TSP | x | | x | x | x | x | x | x | x | | x |
| PM10 | x | | x | x | x | x | x | x | x | | x |
| PM2.5 | x | | x | x | x | x | x | x | x | | x |
| BC | | | x | x | x | x | x | x | x | | x |
| Priority heavy metals | | | | | | | | | | | |
| Pb | | | x | x | x | x | x | x | x | | x |
| Cd | | | x | x | x | x | x | x | x | | x |
| Hg | | | x | x | x | x | x | x | x | | x |
| POPs Annex II | | | | | | | | | | | |
| PCB | | | x | x | | x | x | | x | | |

| Main pollutants | 5A | 5B | 5C1a | 5C1bi | 5C1bii | 5C1biii | 5C1biv | 5C2 | 5C1bv | 5D | 5E |
|---------------------------|----|----|------|-------|--------|---------|--------|-----|-------|----|----|
| POPs Annex III | | | | | | | | | | | |
| Dioxins | | | x | x | x | x | x | x | x | | x |
| PAH | | | x | x | x | x | x | x | x | | |
| HCB | | | x | x | | x | x | | x | | |
| Other heavy metals | | | | | | | | | | | |
| As | | | x | x | x | x | x | x | x | | |
| Cr | | | x | x | x | x | x | x | x | | |
| Cu | | | x | x | x | x | x | x | x | | |
| Ni | | | x | x | x | x | x | | x | | |
| Se | | | x | x | | x | | x | x | | |
| Zn | | | x | x | x | | x | x | x | | |

In 2020, *open burning of waste* (5C2) is key category for Cd and BC, *car and building fires* (5E) is a key category for PM and dioxins emissions while clinical waste incineration (5C1biii) is key category for HCB. In 1990, *municipal waste incineration* (5C1a), *industrial waste incineration* (5C1 bi) and *car and building fires* (5E) are key categories for dioxins emissions. As regard the trend, *municipal waste incineration* (5C1a) and *car and building fires* (5E) are key categories for dioxins emissions, *biological treatment of waste* (5B2) is key category for NH₃ emissions, *car and building fires* (5E) is a key category for PM₁₀ and PM_{2.5}, whereas *open burning of agricultural waste* (5C2) is key category for Cd emissions.

The waste sector, and in particular Waste incineration (5C), is a source of different pollutants; for the main pollutants, in 2020, the sector accounts for:

- 17.4 % in national total Dioxin emissions;
- 9.4 % in national total Cd emissions;
- 6.1 % in national total BC emissions;
- 6.2% in national total HCB emissions.

Moreover, the sector comprises 2.5% of total NH₃ emissions, 2.4% of CO, 1.8% of Hg, 1.2% of NMVOC, 1.8% of PAH, 4.6% and 3.9% in national total of PM_{2.5} and PM₁₀ emissions respectively and for what concerns all remaining pollutants are below 1%.

7.2 METHODOLOGICAL ISSUES

7.2.1 Solid waste disposal on land (5A)

Solid waste disposal on land is a major source concerning greenhouse gas emissions but not concerning air pollutants. Notwithstanding, NMVOC and NH₃ emissions are estimated, as a percentage of methane emitted, calculated using the IPCC Tier 2 methodology (IPCC, 1997; IPCC, 2000), through the application of the First Order Decay Model (FOD). As a consequence of the last review process also PM emissions have been estimated. A detailed description of the model and its application to Italian landfills is reported in the National Inventory Report on the Italian greenhouse gas inventory (ISPRA, 2022 [a]).

Following the suggestion of NEC review (EEA, 2017 [a]) more info about the extraction and use of biogas is provided below.

The amount of biogas recovery in landfills has increased as a result of the implementation of the European Directive on the landfill of waste (EC, 1999); the amounts of biogas recovered and flared have been estimated taking into account the amount of energy produced, the energy efficiency of the methane recovered, the captation efficiency and the efficiency in recovering methane for energy purposes assuming that the rest of methane captured is flared.

Emissions for all the relevant pollutants from biogas recovered from landfills and used for energy purposes are reported in the energy sector in “1A4a biomass” category together with wood, the biomass fraction of incinerated waste and biogas from wastewater plants. In the following scheme consumptions and low calorific values are reported for the year 2020.

1A4a biomass detailed activity data. Year 2020

| Fuels | | Consumption (Gg) | LCV (TJ/Gg) |
|--------------------------------------|-------------------|-------------------------|--------------------|
| <i>Wood and similar</i> | <i>Wood</i> | 292.51 | 10.47 |
| | <i>Steam Wood</i> | 0.00 | 30.80 |
| <i>Incinerated waste (biomass)</i> | | 2140.03 | 11.39 |
| <i>Biogas from landfills</i> | | 218.78 | 55.07 |
| <i>Biogas from wastewater plants</i> | | 25.90 | 55.07 |

It is assumed that landfill gas composition is 50% VOC. The percentage by weight of CH₄ compared to the total VOC emitted is 98.7%. The remaining 1.3% (NMVOC) consists of paraffinic, aromatic and halogenated hydrocarbons (Gaudioso et al., 1993): this assumption refers to US EPA data (US EPA, 1990). As regard ammonia, emission factor has been assumed equal to 1 volume per cent of VOC too (Tchobanoglous et al., 1993).

According with the discussion during the ESD review about CH₄ emissions from landfills and the consequent technical correction (EEA, 2017 [b]), Italy revised the half-life values considering the distribution of dry and wet regions in Italy. New data (CREA, 2017) regarding raining and evapotranspiration have been elaborated allowing to distinguish between dry and wet region and the estimates have been splitted in two components considering the location of SWDS.

Methane, and consequently NMVOC and NH₃ air pollutants, is emitted from the degradation of waste occurring in municipal landfills, both managed and unmanaged (due to national legislation, from 2000 municipal solid wastes are disposed only into managed landfills). The main parameters that influence the estimation of emissions from landfills are, apart from the amount of waste disposed into managed landfill: the waste composition (which vary through the years in the model); the fraction of methane in the landfill gas (included in VOC, which has been assumed equal to 50%) and the amount of landfill gas collected and treated. These parameters are strictly dependent on the waste management policies throughout the waste streams which consist of: waste generation, collection and transportation, separation for resource recovery, treatment for volume reduction, stabilisation, recycling and energy recovery and disposal at landfill sites.

Basic data on waste production and landfills system are those provided by the national Waste Cadastre, basically built with data reported through the Uniform Statement Format (MUD). The Waste Cadastre is formed by a national branch, hosted by ISPRA, and by regional and provincial branches.

These figures are elaborated and published by ISPRA yearly since 1999: the yearbooks report waste production data, as well as data concerning landfilling, incineration, composting, anaerobic digestion and generally waste life-cycle data (APAT-ONR, several years; ISPRA, several years [a]).

For inventory purposes, a database of waste production, waste disposal in managed and unmanaged landfills and sludge disposal in landfills was created and it has been assumed that waste landfilling started in 1950.

For the year 2020, the non-hazardous landfills in Italy disposed 5,817 kt of MSW and 2,910 kt of industrial wastes, as well as 162 kt of sludge from urban wastewater treatment plants.

In Table 7.2, the time series of AMSW and domestic sludge disposed into non-hazardous landfills from 1990 is reported.

Table 7.2 *Trend of MSW production and MSW, AMSW and domestic sludge disposed in landfills (Gg)*

| ACTIVITY DATA (Gg) | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|---|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|--------------|--------------|
| MSW production | 22,231 | 25,780 | 28,959 | 31,664 | 32,479 | 29,524 | 29,572 | 30,158 | 30,079 | 28,945 |
| MSW disposed in landfills for non-hazardous waste | 17,432 | 22,459 | 21,917 | 17,226 | 15,015 | 7,819 | 6,927 | 6,496 | 6,283 | 5,817 |
| Assimilated MSW disposed in landfills for non-hazardous waste | 2,828 | 2,978 | 2,825 | 2,914 | 3,508 | 3,222 | 3,899 | 3,512 | 3,256 | 2,910 |
| Sludge disposed in managed landfills for non-hazardous waste | 2,454 | 1,531 | 1,326 | 544 | 346 | 387 | 342 | 261 | 232 | 162 |
| Total Waste to managed landfills for non-hazardous waste | 16,363 | 21,897 | 26,069 | 20,684 | 18,870 | 11,428 | 11,167 | 10,269 | 9,771 | 8,889 |
| Total Waste to unmanaged landfills for non-hazardous waste | 6,351 | 5,071 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total Waste to landfills for non-hazardous waste | 22,714 | 26,968 | 26,069 | 20,684 | 18,870 | 11,428 | 11,167 | 10,269 | 9,771 | 8,889 |

7.2.2 Biological treatment of waste (5B)

Under this category, NMVOC and NH₃ emissions from compost production and from anaerobic digestion are reported.

The amount of waste treated in biological treatments has shown a great increase from 1990 to 2020 (from 283,879 Mg to 6,818,914 Mg for composting and from 0 to 81,151 Mg of N-excreted from manure management).

Information on input waste to composting plants is published yearly by ISPRA since 1996, including data for 1993 and 1994 (ANPA, 1998; APAT-ONR, several years; ISPRA, several years [a]), while for 1987 and 1995 only data on compost production are available (MATTM, several years [a]; AUSITRA-Assoambiente, 1995); on the basis of this information the whole time series has been reconstructed.

The composting plants are classified in two different kinds: the plants that treat a selected waste (food, market, garden waste, sewage sludge and other organic waste, mainly from the agro-food industry); and the mechanical-biological treatment plants, that treat the unselected waste to produce compost, refuse derived fuel (RDF), and a waste with selected characteristics for landfilling or incinerating system.

It is assumed that 100% of the input waste to the composting plants from selected waste is treated as compost, while in mechanical-biological treatment plants 30% of the input waste is treated as compost on the basis of national studies and references (Favoino and Cortellini, 2001; Favoino and Girò, 2001). NMVOC emission factor (51g NMVOC kg⁻¹ treated waste) is from international scientific literature too (Finn and Spencer, 1997).

NH₃ emissions from biogas facilities (anaerobic digesters) in the agriculture sector have been updated in the previous submission on the basis of the study carried out by CRPA (CRPA, 2018) and in particular data relative to the percentages of the different substrates that feed the anaerobic digesters and data relative to the average content of volatile solids by type of substrates have been changed. As a result of these changes, the amount of manure sent to the digesters decreases considerably and also the NH₃ emissions. These emissions have been subtracted from 3B manure management category (cattle, swine and poultry) and allocated in the anaerobic digestion at biogas facilities (5B2 of the waste sector). In the current submission, Ni and Hg emissions from combustion of removable cereal residues have been estimated while recalculations occur because of the correction of the amount of biomass available for open burning since 1990.

7.2.3 Waste Incineration (5C1a – 5C1b)

Regarding waste incineration, methodology used for estimating emissions is based on and consistent with the EMEP/CORINAIR Guidebook (EMEP/EEA, 2019).

In this sector only emissions from facilities without energy recovery are reported, whereas emissions from waste incineration facilities with energy recovery are reported in the Energy Sector 1A4a because energy produced in incinerators is still prevalently used to satisfy the internal energy demand of the plants (auto

production) and in this sense it would be wrong, according to the guidelines, to report them under 1A1a Public Electricity and Heat Production instead of 1A4a. In 2020, about 99% of the total amount of waste incinerated is treated in plants with energy recovery system.

Existing incinerators in Italy are used for the disposal of municipal waste, together with some industrial waste, sanitary waste and sewage sludge for which the incineration plant has been authorized by the competent authority. Other incineration plants are used exclusively for industrial and sanitary waste, both hazardous and not, and for the combustion of waste oils, whereas there are plants that treat residual waste from waste treatments, as well as sewage sludge.

A complete database of the incineration plants is now available, updated with the information reported in the yearly report on waste production and management published by ISPRA (APAT-ONR, several years; ISPRA, several years). For each plant a lot of information is reported, among which the year of the construction and possible upgrade, the typology of combustion chamber and gas treatment section, energy recovery section (thermal or electric), and the type and amount of waste incinerated (municipal, industrial, etc.). A specific emission factor is therefore used for each pollutant combined with plant specific waste activity data.

In Table 7.3, emission factors for each pollutant and waste typology are reported. Emission factors have been estimated on the basis of a study conducted by ENEA (De Stefanis P., 1999), based on emission data from a large sample of Italian incinerators (FEDERAMBIENTE, 1998; AMA-Comune di Roma, 1996), legal thresholds (Ministerial Decree 19 November 1997, n. 503 of the Ministry of Environment; Ministerial Decree 12 July 1990) and expert judgements.

Since 2010, emission factors for urban waste incinerators have been updated on the basis of data provided by plants (ENEA-federAmbiente, 2012; De Stefanis P., 2012) concerning the annual stack flow, the amount of waste burned and the average concentrations of the pollutants at the stack. As the emission factors are considerably lower than the old ones due to the application of very efficient abatement systems it was necessary to apply a linear smoothing methodology assuming a progressive application of the abatement systems between 2005 and 2010. In a similar way, emission factors for industrial waste incinerators have been updated from 2010 onwards on the basis of the 2019 EMEP/EEA Guidebook. Similarly to municipal waste smoothing has been applied between 2005 and 2010 supposing a linear application of the abatement systems. In particular, for 5C1bi the HCB emission factors comes from table 3-1 of the 2019 EMEP/EEA Guidebook considering an abatement efficiency of 90% while PCB emission factors derives from 2007 Guidebook but considering 90% abatement (no value in the 2019 EMEP/EEA Guidebook). For 5C1biii the emission factors from 2019 EMEP/EEA Guidebook have been used for PCB while HCB emission factors derives from 2007 Guidebook because of a lack of information in the last guidebook. For 5C1biv emission factors from 2019 EMEP/EEA Guidebook have been considered without abatement because information about this specific category of plant are under investigation.

Table 7.3 Emission factors for waste incineration

| Air Pollutant | u. m | Municipal 1990-2009 | Municipal since 2010 | Industrial 1990-2009 | Industrial since 2010 | Clinical 1990-2009 | Clinical since 2010 | Sludge 1990-2009 | Sludge since 2010 | Oil 1990-2009 | Oil since 2010 |
|-----------------|------|---------------------|----------------------|----------------------|-----------------------|--------------------|---------------------|------------------|-------------------|---------------|----------------|
| NO _x | kg/t | 1.15 | 0.62 | 2.00 | 2.00 | 0.60 | 0.60 | 3.00 | 3.00 | 2.00 | 2.00 |
| CO | kg/t | 0.07 | 0.07 | 0.56 | 0.56 | 0.08 | 0.08 | 0.60 | 0.60 | 0.08 | 0.08 |
| NM VOC | kg/t | 0.46 | 0.46 | 7.40 | 7.40 | 7.40 | 7.40 | 0.25 | 0.25 | 7.40 | 7.40 |
| SO ₂ | kg/t | 0.39 | 0.02 | 1.28 | 1.28 | 0.03 | 0.03 | 1.80 | 1.80 | 1.28 | 1.28 |
| PM10 | g/t | 46.00 | 6.06 | 240.00 | 0.70 | 25.68 | 25.68 | 180.00 | 41.00 | 240.00 | 0.07 |
| PM2.5 | g/t | 46.00 | 6.06 | 240.00 | 0.40 | 25.68 | 25.68 | 180.00 | 11.00 | 240.00 | 0.04 |
| As | g/t | 0.05 | 0.02 | 0.12 | 0.00 | 0.00 | 0.00 | 0.50 | 0.24 | 0.12 | 0.00 |
| Cu | g/t | 1.00 | 0.00 | 1.20 | 0.12 | 0.56 | 0.56 | 10.00 | 2.00 | 1.20 | 0.01 |
| Se | g/t | 0.01 | 0.01 | 0.01 | 0.00 | 0.04 | 0.04 | - | 0.01 | 0.01 | 0.00 |
| Zn | g/t | 0.02 | 0.02 | 12.60 | 1.26 | - | - | 10.00 | 3.30 | 12.60 | 0.13 |
| Cd | g/t | 0.25 | 0.01 | 0.80 | 0.01 | 0.00 | 0.00 | 1.20 | 0.80 | 0.80 | 0.00 |
| Cr | g/t | 0.45 | 0.00 | 1.60 | 0.16 | 0.01 | 0.01 | 3.00 | 0.70 | 1.60 | 0.02 |
| Hg | g/t | 0.15 | 0.03 | 0.80 | 0.01 | 0.04 | 0.04 | 1.20 | 1.15 | 0.80 | 0.01 |
| Ni | g/t | 16.35 | 0.00 | 0.80 | 0.01 | 0.03 | 0.03 | 3.00 | 0.40 | 0.80 | 0.00 |
| Pb | g/t | 1.35 | 1.04 | 24.00 | 0.13 | 0.02 | 0.02 | 3.00 | 0.40 | 24.00 | 0.01 |
| PAH | g/t | 0.05 | 0.00 | 0.48 | 0.00 | 0.00 | 0.00 | 0.60 | 0.00 | 0.48 | 0.00 |
| PCB | g/t | 0.005 | 0.00005 | 0.0050 | 0.0005 | 0.020 | 0.020 | 0.005 | 0.0045 | - | - |
| HCB | g/t | 0.001 | 0.00002 | 0.0001 | 0.0002 | 0.019 | 0.019 | 0.500 | 0.0047 | - | - |

Concerning dioxin emissions, clinical and industrial emission factors are also derived from data collected from a large sample of Italian incinerators and legal thresholds, as well as expert judgement; in particular for municipal solid waste, emission factors vary within the years and the facility on the basis of plant technology (i.e. typology of combustion chamber and gas treatment section) and the year of the upgrade. This site specific evaluation has been possible thanks to a study conducted in the past for a sample of municipal waste incinerators located in Regione Lombardia in order to produce an assessment of field-based values applicable to other facilities with the same characteristics (Pastorelli et al., 2001) and, since 2010 urban waste data, thanks to the abovementioned survey (ENEA-federAmbiente, 2012). Moreover, for the incineration plants reported in the national EPER/PRTR register, verification of emissions has been carried out.

In Table 7.4 dioxin emission factors for waste incineration are reported for 1990 and 2020.

Table 7.4 Dioxin emission factors for 1990 and 2020

| Waste Typology | u.m | 1990 | 2020 |
|----------------|-----|-----------|------|
| Municipal | g/t | 115 - 1.6 | 0.1 |
| Clinical | g/t | 200 | 0.5 |
| Industrial | g/t | 80 - 135 | 0.5 |
| Sludge | g/t | 77 | 0.5 |
| Oil | g/t | 200 | 0.5 |

In Table 7.5 activity data are reported by type of waste.

Table 7.5 Amount of waste incinerated by type (Gg)

| Waste incinerated | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|---------|---------|---------|
| <i>Gg</i> | | | | | | | | | | | |
| Total waste | 1,656. 2 | 2,149. 1 | 3,061. 7 | 4,964. 2 | 6,977. 3 | 7,534. 7 | 7,587. 7 | 7,477. 2 | 7,726.0 | 7,509.5 | 7,514.3 |
| with energy recovery | 911.2 | 1,557. 8 | 2,749. 7 | 4,720. 6 | 6,795. 9 | 7,431. 5 | 7,501. 1 | 7,386. 3 | 7,633.9 | 7,420.0 | 7,427.4 |
| without energy recovery | 745.0 | 591.3 | 312.0 | 243.5 | 181.4 | 103.2 | 86.7 | 91.0 | 92.1 | 89.5 | 86.9 |
| Municipal waste (5C1a) | 1,025. 6 | 1,436. 6 | 2,324. 9 | 3,219. 9 | 4,336. 9 | 4,698. 4 | 4,453. 9 | 4,325. 1 | 4,577.3 | 4,358.9 | 4,280.1 |
| with energy recovery | 626.4 | 1,185. 5 | 2,161. 4 | 3,168. 0 | 4,284. 0 | 4,698. 4 | 4,453. 9 | 4,325. 1 | 4,577.3 | 4,358.9 | 4,280.1 |
| without energy recovery | 399.2 | 251.1 | 163.5 | 51.9 | 52.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Industrial waste (5C1b i-ii-iv) | 496.1 | 560.7 | 626.5 | 1,618. 1 | 2,505. 3 | 2,734. 7 | 3,029. 1 | 3,051. 9 | 3,041.2 | 3,072.7 | 3,158.7 |
| with energy recovery | 259.5 | 331.2 | 511.6 | 1,447. 0 | 2,399. 4 | 2,676. 1 | 2,988. 0 | 2,999. 1 | 2,991.7 | 3,022.5 | 3,109.9 |
| without energy recovery | 236.6 | 229.6 | 114.8 | 171.1 | 105.9 | 58.6 | 41.1 | 52.8 | 49.5 | 50.2 | 48.7 |
| Clinical waste (5C1biii) | 134.5 | 151.7 | 110.3 | 126.2 | 135.1 | 101.6 | 104.7 | 100.2 | 107.5 | 77.9 | 75.6 |
| with energy recovery | 25.3 | 41.1 | 76.7 | 105.7 | 112.5 | 57.0 | 59.1 | 62.0 | 64.9 | 38.6 | 37.5 |
| without energy recovery | 109.2 | 110.6 | 33.6 | 20.5 | 22.6 | 44.6 | 45.5 | 38.2 | 42.7 | 39.3 | 38.1 |

7.2.4 Cremation of corpses (5C1bv)

Emissions from incineration of human bodies in crematoria have been carried out for the entire time series. The methodology used for estimating emissions is based on and conform to the EMEP/EEA Air Pollutant Emission Inventory Guidebook (EMEP/EEA, 2019).

Activity data have been supplied by a specific branch of Federutility, which is the federation of energy and water companies (SEFIT, several years), whereas emission factors derive from a survey carried out by the

same subject in 2015. For some metal, such as Pb, Cd, As, Cr, Cu, Ni and Se EFs are those reported in the Guidebook 2019.

Up to some years ago cremation was not so popular in Italy also because the Catholic Church encouraged burial. Partly because cemeteries are becoming overcrowded, the number of cremations in Italy has risen from 5,809 in 1990 to 247,840 in 2020. Moreover, it is practice to cremate also mortal remains: activity data have been supplied too by SEFIT, from 1999, whereas mortal remains from 1990 to 1998 have been reconstructed on the basis of an expert judgment (SEFIT, several years).

In Table 7.6 time series of number of cremations, mortal remains, as well as annual deaths and crematoria in Italy are reported. The major emissions from crematoria are nitrogen oxides, carbon monoxide, sulphur dioxide, particulate matter, mercury, hydrogen fluoride (HF), hydrogen chloride (HCl), NMVOCs, other heavy metals, and some POPs. As a consequence of the review process, emissions have been revised to take in account the implementation of abatement technology over the years. In Table 7.7 emission factors for cremation 2020 are reported.

Table 7.6 Cremation time series (activity data)

| Cremation of corpses | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2017 | 2018 | 2019 | 2020 |
|----------------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| Cremations | 5,809 | 15,436 | 30,167 | 48,196 | 77,379 | 137,168 | 170,903 | 183,146 | 194,669 | 247,840 |
| Deaths | 543,700 | 555,203 | 560,241 | 567,304 | 587,488 | 653,000 | 649,061 | 633,133 | 634,432 | 746,146 |
| Mortal remains | 1,000 | 1,750 | 1,779 | 9,880 | 18,899 | 34,178 | 36,425 | 37,538 | 38,305 | 29,266 |
| % of cremation | 1.07 | 2.78 | 5.38 | 8.50 | 13.17 | 21.01 | 26.33 | 28.93 | 30.68 | 33.22 |
| Crematoria | ND | 31 | 35 | 43 | 53 | 70 | 79 | 83 | 85 | 87 |

Table 7.7 Emission factors for cremation of corpses - 2020

| Air pollutant | u.m. | Cremation |
|----------------------|---------|-----------|
| NO _x | kg/body | 0.4739 |
| CO | kg/body | 0.0047 |
| NMVOC | kg/body | 0.0091 |
| SO _x | kg/body | 0.0093 |
| PM10 | g/body | 2.1483 |
| PM2.5 | g/body | 2.1483 |
| Pb | mg/body | 30.0300 |
| Cd | mg/body | 5.0300 |
| Hg | mg/body | 0.0070 |
| As | mg/body | 13.6100 |
| Cr | mg/body | 13.5600 |
| Cu | mg/body | 12.4300 |
| Ni | mg/body | 17.3300 |
| PAH (benzo(a)pyrene) | µg/body | 176.2000 |
| Dioxins | µg/body | 0.0190 |

7.2.5 Small scale waste burning (5C2)

The open burning of agricultural waste is a key category for Cd and BC emissions. Dioxins, TSP, PM10, PM2.5, BC, CO, NMVOC, PAH, SO_x, NO_x and heavy metals emissions have been estimated. No estimations were performed for NH₃ emissions as well as other POPs.

A country-specific methodology has been used. Parameters taken into consideration are the following:

1. Amount of removable residues (t), estimated with annual crop production (ISTAT, several years [a], [b]; ISTAT, 2017 [a], [b]) and removable residues/product ratio (IPCC, 1997; CESTAAT, 1988; Borgioli E., 1981).
2. Amount of dry residues in removable residue (t dry matter), calculated with amount of removable fixed residues and fraction of dry matter (IPCC, 1997; CESTAAT, 1988; Borgioli E., 1981).
3. Amount of removable dry residues oxidized (t dry matter), assessed with amount of dry residues in the removable residues, burnt fraction of removable residues (CESTAAT, 1988) and fraction of residues oxidized during burning (IPCC, 1997).
4. Amount of carbon from removable residues burning release in air (t C), calculated with the amount of removable dry residue oxidized and the fraction of carbon from the dry matter of residues (IPCC, 1997; CESTAAT, 1988).

5. C-CH₄ from removable residues burning (t C-CH₄), calculated with the amount of carbon from removable residues burning release in air and default emissions rate for C-CH₄, equal to 0.005 (IPCC, 1997).
6. C-CO from removable residues burning (t C-CO), calculated with the amount of carbon from removable residues burning release in air and default emissions rate for C-CO, equal to 0.06 (IPCC, 1997).
7. Amount of nitrogen from removable residues burning release in air (t N), calculated with the amount of removable dry residue oxidized and the fraction of nitrogen from the dry matter of residues. The fraction of nitrogen has been calculated considering raw protein content from residues (dry matter fraction) divided by 6.25.
8. N-NO_x from removable residues burning (t N-NO_x), calculated with the amount of nitrogen from removable residues burning release in air and the default emissions rate for N- NO_x, equal to 0.121 (IPCC, 1997).

NMVOC emissions have been considered equal to CH₄ emissions. As regards the other pollutants, heavy metals, Dioxin and PAH emission factors are from the EMEP/EEA Guidebook (EMEP/EEA, 2016) and emissions have been added as requested by the NECD review process (EEA, 2018) (Table 7.8).

Table 7.8 Emission factors for burning of agriculture residues

| Air pollutant | u.m. | Removable residues | | | | References |
|------------------------|------|----------------------------------|-------|------|-----|---------------------|
| | | Wheat – Barley – Rice – Orchards | | | | |
| Benzo(a)pyrene | g/t | 67.7 | 98.8 | 19 | 1.5 | EMEP/EEA, 2016 |
| Benzo(b)fluoranthene | g/t | 189.1 | 307.4 | 31.5 | 2.8 | |
| Benzo(k)fluoranthene | g/t | 80.7 | 77 | 23.1 | 6.2 | |
| Indeno(1,2,3-cd)pyrene | g/t | 57.9 | 38.2 | 14.5 | | |
| PM10 | g/t | | | 3.3 | | EMEP/CORINAIR, 2007 |
| PM2.5 | g/t | | | 2.8 | | EMEP/CORINAIR, 2007 |
| Dioxins | g/t | | | 10 | | EMEP/CORINAIR, 2016 |
| BC | g/t | | | 1.2 | | EMEP/EEA, 2013 |

Removable residues from agriculture production are estimated for each crop type (cereal, green crop, permanent cultivation) taking into account the amount of crop produced, from national statistics (ISTAT, several years [a], [b]; ISTAT, 2017 [a], [b]), the ratio of removable residue in the crop, the dry matter content of removable residue, the ratio of removable residue burned, the fraction of residues oxidised in burning, the carbon and nitrogen content of the residues. Most of these wastes refer especially to the prunes of olives and wine, because of the typical national cultivation. Activity data (agricultural production) used for estimating burning of agriculture residues are reported in Table 7.9. Emissions due to stubble burning, which are emissions only from the agriculture residues burned on field, are reported in the agriculture sector, under 3.F.

Under the waste sector the burning of removable agriculture residues that are collected and could be managed in different ways (disposed in landfills, used to produce compost or used to produce energy) is reported. Different percentages of the removable agriculture residue burnt for different residues are assumed, varying from 10% to 90%, according to national and international literature. Moreover, these removable wastes are assumed to be all burned in open air (e.g. on field), taking in consideration the highest available CO, NMVOC, PM and dioxins emission factors as reported in the table above. The amount of biomass from pruning used for domestic heating is reported in the energy sector in the 1A4b category as biomass fuel.

Table 7.9 Time series of crop productions (Gg)

| Production | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|----------------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|---------|
| Gg | | | | | | | | | | | |
| Cereals | | | | | | | | | | | |
| Wheat | 8,108.5 | 7,946.1 | 7,427.7 | 7,717.1 | 6,849.9 | 7,394.5 | 8,037.9 | 6,966.5 | 6,932.9 | 6,576.6 | 6,553.9 |
| Rye | 20.8 | 19.8 | 10.3 | 7.9 | 13.9 | 13.2 | 13.2 | 11.1 | 10.6 | 12.5 | 11.5 |
| Barley | 1,702.5 | 1,387.1 | 1,261.6 | 1,214.1 | 944.3 | 955.1 | 988.3 | 984.3 | 1,010.3 | 1,072.4 | 1,090.6 |
| Oats | 298.4 | 301.3 | 317.9 | 429.2 | 288.9 | 261.4 | 260.8 | 229.0 | 243.4 | 238.1 | 242.7 |

| Production | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <i>Gg</i> | | | | | | | | | | | |
| Rice | 1,290.7 | 1,320.9 | 1,245.6 | 1,444.8 | 1,574.3 | 1,505.8 | 1,598.0 | 1,516.0 | 1,480.9 | 1,498.1 | 1,513.1 |
| Maize | 5,864 | 8,454 | 10,140 | 10,428 | 8,496 | 7,074 | 6,839 | 6,048 | 6,179 | 6,259 | 6,771 |
| Sorghum | 114.2 | 214.8 | 215.2 | 184.9 | 275.6 | 294.2 | 313.8 | 240.7 | 293.9 | 312.4 | 361.7 |
| Woody crops | | | | | | | | | | | |
| Grapes | 8,438.0 | 8,447.7 | 8,869.5 | 8,553.6 | 7,839.7 | 7,915.0 | 8,044.1 | 7,169.7 | 8,513.6 | 7,862.9 | 8,222.4 |
| Olives | 912.5 | 3,323.5 | 2,810.3 | 3,774.8 | 3,117.8 | 2,732.9 | 2,016.0 | 2,598.5 | 1,953.5 | 2,194.1 | 2,207.2 |
| Citrus Orchards | 2,868.8 | 2,607.7 | 3,100.2 | 3,518.1 | 3,820.6 | 3,151.5 | 2,766.4 | 2,811.3 | 2,631.3 | 2,895.9 | 2,940.0 |
| Orchards | 5,793.5 | 5,406.6 | 5,952.2 | 6,034.5 | 5,777.3 | 5,988.8 | 5,927.5 | 5,360.3 | 5,608.0 | 5,318.0 | 5,408.7 |
| Carobs | 29.2 | 44.4 | 38.1 | 31.7 | 25.3 | 31.5 | 28.9 | 28.9 | 37.0 | 35.9 | 36.9 |
| Total | 35,441 | 39,474 | 41,388 | 43,339 | 39,023 | 37,318 | 36,834 | 33,965 | 34,894 | 34,276 | 35,360 |

7.2.6 Wastewater treatments (5D)

The biogas collected from the anaerobic digestion of wastewaters is burned with heat/energy recovery and relevant emissions are reported in the energy sector. As regards NMVOC emissions from wastewater handling, consequently to the NECD review (EEA, 2017 [a]). At present, no reliable EFs are available for wastewater treatment plants at national level. By using EFs from the 2019 EEA/EMEP Guidebook both for domestic and industrial wastewater and the volumes of wastewater produced NMVOC emissions resulted in the time series reported in the table below (Table 7.10).

Table 7.10 Time series of NMVOC emissions (Gg)

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|-----------------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| <i>NMVOC (Gg)</i> | | | | | | | | | | | |
| Industrial wastewater | 0.0136 | 0.0139 | 0.0138 | 0.0130 | 0.0104 | 0.0099 | 0.0104 | 0.0104 | 0.0106 | 0.0105 | 0.0098 |
| Domestic wastewater | 0.0636 | 0.0821 | 0.0898 | 0.1005 | 0.1063 | 0.1030 | 0.1170 | 0.1189 | 0.1209 | 0.1228 | 0.1247 |
| Total | 0.0772 | 0.0961 | 0.1036 | 0.1135 | 0.1160 | 0.1129 | 0.1275 | 0.1293 | 0.1315 | 0.1333 | 0.1345 |

7.2.7 Other waste (5E)

On the basis of the Final review report of the 2017 Comprehensive technical review of national emission inventories (EEA, 2017 [a]) emissions from category 5E – Car and Building Fires have been estimated. Buildings have been subdivided into 4 subcategories: detached house, undetached house, apartment buildings and industrial buildings and the distribution of population in the different typology of building has been derived from Eurostat. Data regarding the number of car and building fires have been derived from the Annually statistics of fire service in Italy (Annually statistics of fire service in Italy, several years) while EFs are coherent with the Guidebook EMEP/EEA 2016 deriving from Aasestad, 2007 for particulate matter (TSP=PM10=PM2.5) while BC EF has been derived from IIASA report (IIASA, 2004). No data about car and building fires are available before 2000 so 90's data have been reconstructed on the basis of the national population and the resulting time series are reported in Tab. 7.11. On the basis of the last review reports (EEA, 2019; EEA, 2020; EEA 2021) PCDD/F emissions have been estimated and revised. After the technical correction during the 2019 NECD review, Italy investigated emission factors because those reported in the guidebook seems to be unreliable or not fitting to the Italian context. In particular, the emission factors reported in the EMEP/EEA Guidebook “used for particles in the inventory are given by scaling the emission factors used for combustion of fuelwood in the households” (Aasestad, 2007) but the Italian buildings are made up of the vast majority of reinforced concrete. Despite this, following the last review process, Italy decided to use these emission factors pending the availability of values deriving from new studies. Further, Cd, Hg and Pb emissions have been estimated for this source on the basis of 2019 EMEP/EEA Guidebook.

Table 7.11 PM2.5 and BC emissions from the category 5E

| | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|--------------------------------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Detached house fires (n°) | 10,580 | 10,592 | 10,614 | 10,861 | 9,213 | 9,577 | 10,542 | 11,610 | 11,472 | 13,222 | 13,171 |
| Undetached house fires (n°) | 8,186 | 8,195 | 8,212 | 8,404 | 6,928 | 11,484 | 12,641 | 13,921 | 13,756 | 12,728 | 12,679 |
| Apartment building fires (n°) | 18,843 | 18,865 | 18,904 | 19,344 | 18,891 | 23,278 | 25,624 | 28,219 | 27,884 | 28,913 | 28,802 |
| Industrial building fires (n°) | 4,931 | 4,936 | 4,947 | 5,062 | 4,560 | 4,872 | 8,033 | 6,814 | 4,257 | 3,599 | 3,192 |
| Car fires (n°) | 25,924 | 25,954 | 26,008 | 26,614 | 22,735 | 22,680 | 22,696 | 23,537 | 22,761 | 22,170 | 18,641 |
| 5E PM2.5 (Gg) | 3.04 | 3.05 | 3.08 | 3.13 | 2.76 | 3.29 | 3.69 | 4.00 | 3.89 | 0.48 | 4.07 |
| 5E BC (Gg) | 0.5621 | 0.5644 | 0.5694 | 0.5783 | 0.5098 | 0.6085 | 0.6824 | 0.7406 | 0.7191 | 0.0892 | 0.0892 |
| | 55 | 10 | 40 | 79 | 53 | 49 | 06 | 21 | 12 | 71 | 71 |
| 5E PCDD/F (g Iteq) | 31.108 | 31.233 | 31.511 | 32.006 | 28.197 | 33.557 | 37.551 | 40.735 | 39.560 | 41.689 | 41.256 |
| | 627 | 380 | 758 | 410 | 442 | 548 | 092 | 381 | 174 | 395 | 888 |

7.3 TIME SERIES AND KEY CATEGORIES

The following Table 7.12 presents an outline of the weight of the different categories for each pollutant in the waste sector for the year 2020. Key categories are those shaded.

Table 7.12 Key categories in the waste sector in 2020

| | 5A | 5B1 | 5B2 | 5C1bi | 5C1biii | 5C1biv | 5C1bv | 5C2 | 5D1 | 5D2 | 5E |
|-----------------------|--------|------|------|----------|---------|---------|--------|------|------|-------|-------|
| | % | | | | | | | | | | |
| SO_x | | | | 0.05 | 0.001 | 0.03 | 0.003 | 0.10 | | | |
| NO_x | | | | 0.01 | 0.004 | 0.01 | 0.02 | 0.34 | | | |
| NH₃ | 1.68 | 0.05 | 0.75 | | | | | | | | |
| NMVOC | 0.85 | 0.04 | | 0.03 | 0.03 | 0.0004 | 0.0003 | 0.25 | 0.01 | 0.001 | |
| CO | | | | 0.001 | 0.0002 | 0.0004 | 0.0001 | 2.40 | | | |
| PM10 | 0.001 | | | 0.00001 | 0.001 | 0.0003 | 0.0004 | 1.49 | | | 2.45 |
| PM2.5 | 0.0002 | | | 0.00001 | 0.001 | 0.0001 | 0.0004 | 1.59 | | | 3.05 |
| BC | | | | 0.000003 | 0.0002 | 0.00003 | | 5.63 | | | 0.57 |
| Pb | | | | 0.003 | 0.001 | 0.004 | 0.01 | 0.10 | | | 0.02 |
| Cd | | | | 0.01 | 0.001 | 0.28 | 0.04 | 8.73 | | | 0.31 |
| Hg | | | | 0.003 | 0.02 | 0.27 | 0.03 | 1.02 | | | 0.41 |
| PAH | | | | 0.0001 | 0.00001 | 0.00003 | 0.0001 | 1.79 | | | |
| Dioxins | | | | 0.01 | 0.01 | 0.002 | 0.002 | 2.67 | | | 14.73 |
| HCB | | | | 0.05 | 5.58 | 0.49 | 0.06 | | | | |
| PCB | | | | 0.02 | 0.74 | 0.06 | 0.02 | | | | |

Note: key categories are shaded in blue

The following pie charts show, for the main pollutants, the contribution of each sub-category to the total emissions from the waste sector, both for 1990 and 2020 (Figure 7.1, Figure 7.2, Figure 7.3 and Figure 7.4).

Finally, in Table 7.13, emissions time series for each pollutant of the waste sector are reported. In the period 1990-2020, total emissions from incineration plants increase, but whereas emissions from plants with energy recovery show a strong growth, emissions from plants without energy recovery decreased because of the legal constraints which impose the energy production. For 2020, about 99% of the total amount of waste incinerated is treated in plants with energy recovery system reported in 1A4a.




Figure 7.1 Contribution of CO, NO_x and NMVOC sub-category emissions to waste sector total emissions




Figure 7.2 Contribution of POPs Annex III sub-category emissions to waste sector total emissions




Figure 7.3 Contribution of priority heavy metals sub-category emissions to waste sector total emissions




Figure 7.4 Contribution of PM10 and PM2.5 sub-category emissions to waste sector total emissions




Figure 7.5 Time series of dioxin emissions of the waste sector by category (g I-Teq)

Table 7.13 Time series emissions in the waste sector by category and pollutant

| WASTE SECTOR | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Solid waste disposal (5A) | | | | | | | | | | | |
| NMVOG (Gg) | 6.43 | 7.97 | 9.06 | 8.96 | 8.20 | 7.39 | 7.21 | 7.19 | 7.18 | 7.08 | 7.51 |
| NH ₃ (Gg) | 5.21 | 6.46 | 7.35 | 7.26 | 6.65 | 5.99 | 5.85 | 5.83 | 5.82 | 5.74 | 6.09 |
| PM10 (Gg) | 0.004 | 0.006 | 0.005 | 0.004 | 0.004 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 | 0.002 |
| PM2.5 (Gg) | 0.0007 | 0.0008 | 0.0008 | 0.0007 | 0.0006 | 0.0004 | 0.0003 | 0.0004 | 0.0003 | 0.0003 | 0.0003 |
| Biological treatment of waste (5B) | | | | | | | | | | | |
| NMVOG (Gg) | 0.01 | 0.03 | 0.14 | 0.28 | 0.36 | 0.37 | 0.38 | 0.37 | 0.37 | 0.36 | 0.35 |
| NH ₃ (Gg) | 0.01 | 0.06 | 0.10 | 0.31 | 0.46 | 2.62 | 2.63 | 2.69 | 2.68 | 2.86 | 2.87 |
| Waste incineration (5C) | | | | | | | | | | | |
| CO (Gg) | 40.68 | 46.90 | 45.38 | 50.47 | 47.18 | 47.04 | 48.90 | 46.01 | 44.34 | 43.95 | 44.90 |
| NO _x (Gg) | 2.68 | 2.81 | 2.36 | 2.58 | 2.34 | 2.22 | 2.21 | 2.16 | 2.10 | 2.11 | 2.18 |
| NMVOG (Gg) | 4.55 | 4.75 | 3.26 | 3.78 | 3.25 | 2.90 | 3.02 | 2.79 | 2.72 | 2.74 | 2.78 |
| SO _x (Gg) | 0.55 | 0.49 | 0.31 | 0.34 | 0.24 | 0.18 | 0.15 | 0.16 | 0.16 | 0.16 | 0.16 |
| PM10 (Gg) | 2.31 | 2.55 | 2.45 | 2.69 | 2.53 | 2.53 | 2.66 | 2.47 | 2.43 | 2.41 | 2.47 |
| PM2.5 (Gg) | 1.97 | 2.17 | 2.09 | 2.29 | 2.17 | 2.17 | 2.28 | 2.11 | 2.08 | 2.07 | 2.12 |
| BC (Gg) | 0.81 | 0.89 | 0.87 | 0.95 | 0.91 | 0.91 | 0.96 | 0.89 | 0.87 | 0.87 | 0.87 |
| PAH (t) | 1.28 | 1.27 | 1.20 | 1.25 | 1.04 | 1.10 | 1.20 | 1.06 | 1.07 | 1.06 | 1.08 |
| Dioxins (g I-Teq) | 103.22 | 79.68 | 28.76 | 8.18 | 7.73 | 7.73 | 8.11 | 7.52 | 7.39 | 7.35 | 7.53 |
| HCB (kg) | 12.86 | 13.97 | 9.87 | 8.27 | 0.49 | 0.98 | 0.90 | 0.82 | 0.92 | 0.83 | 0.80 |
| PCB (kg) | 5.36 | 4.62 | 2.08 | 1.55 | 0.57 | 1.03 | 0.96 | 0.88 | 0.97 | 0.88 | 0.86 |
| As (t) | 0.09 | 0.09 | 0.07 | 0.07 | 0.05 | 0.05 | 0.05 | 0.05 | 0.05 | 0.05 | 0.05 |
| Cd (t) | 0.60 | 0.59 | 0.48 | 0.51 | 0.34 | 0.37 | 0.40 | 0.35 | 0.36 | 0.34 | 0.35 |
| Cr (t) | 0.66 | 0.58 | 0.35 | 0.39 | 0.09 | 0.10 | 0.09 | 0.09 | 0.09 | 0.09 | 0.09 |
| Cu (t) | 0.99 | 0.87 | 0.55 | 0.49 | 0.12 | 0.16 | 0.12 | 0.14 | 0.14 | 0.13 | 0.13 |
| Hg (t) | 0.33 | 0.32 | 0.23 | 0.30 | 0.21 | 0.09 | 0.08 | 0.08 | 0.09 | 0.08 | 0.08 |
| Ni (t) | 6.79 | 4.37 | 2.84 | 1.05 | 0.04 | 0.05 | 0.04 | 0.05 | 0.05 | 0.04 | 0.04 |
| Pb (t) | 5.90 | 5.53 | 2.76 | 4.03 | 0.24 | 0.18 | 0.17 | 0.17 | 0.16 | 0.16 | 0.17 |
| Se (t) | 0.04 | 0.04 | 0.03 | 0.04 | 0.04 | 0.04 | 0.04 | 0.04 | 0.04 | 0.04 | 0.04 |
| Zn (t) | 5.28 | 6.33 | 4.74 | 5.96 | 3.61 | 3.42 | 3.07 | 3.27 | 3.08 | 3.10 | 3.26 |
| Wastewater (5D) | | | | | | | | | | | |
| NMVOG (Gg) | 0.08 | 0.10 | 0.10 | 0.11 | 0.12 | 0.11 | 0.13 | 0.13 | 0.13 | 0.13 | 0.13 |
| Other waste (5E) | | | | | | | | | | | |
| PM2.5 (Gg) | 3.04 | 3.05 | 3.08 | 3.13 | 2.76 | 3.29 | 3.69 | 4.00 | 3.89 | 0.48 | 4.07 |
| BC (Gg) | 0.56 | 0.56 | 0.57 | 0.58 | 0.51 | 0.61 | 0.68 | 0.74 | 0.72 | 0.09 | 0.09 |
| PCDD/F (g I-Teq) | 31.11 | 31.23 | 31.51 | 32.01 | 28.20 | 33.56 | 37.55 | 40.74 | 39.56 | 41.69 | 41.26 |
| Cd (t) | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| Hg (t) | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 |
| Pb (t) | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 |

7.4 RECALCULATIONS

In the following table the recalculations occurred in the 2022 submission with respect the last year submission are reported at category level.

Table 7.14 Recalculations in the waste sector by category and pollutant

| WASTE SECTOR | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 |
|---|------|------|------|------|------|------|------|-------|-------|-------|
| Solid waste disposal (5A) | | | | | | | | | | |
| NMVOG | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.1% | -0.4% | -1.0% | -1.6% |
| NH ₃ | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.1% | 0.1% | -0.4% | -1.0% | -1.6% |
| PM10 (Gg) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| PM2.5 (Gg) | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| Biological treatment of waste (5B) | | | | | | | | | | |

| WASTE SECTOR | 1990 | 1995 | 2000 | 2005 | 2010 | 2015 | 2016 | 2017 | 2018 | 2019 |
|--------------------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| NMVOG | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 2.7% |
| NH ₃ | 0.0% | -0.2% | -0.1% | -0.2% | -0.2% | -0.2% | -0.2% | -0.2% | -0.2% | -0.1% |
| Waste | | | | | | | | | | |
| incineration (5C) | | | | | | | | | | |
| CO | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| NO _x | 0.1% | 0.2% | 0.5% | 0.9% | 1.7% | 0.4% | 0.4% | 0.5% | 1.7% | 0.5% |
| NMVOG | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 9.2% | 9.8% |
| SO _x | 0.1% | 0.3% | 0.9% | 1.5% | 3.8% | 0.0% | -1.7% | -1.8% | 2.1% | -6.3% |
| PM10 | 0.0% | 0.0% | 0.0% | 0.1% | 0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| PM2.5 | -1.2% | -1.0% | -0.5% | -0.7% | 0.2% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| BC | -0.1% | -0.1% | 0.0% | -0.1% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| PAH | -8.1% | -12.5% | -12.5% | -13.8% | -14.4% | -12.9% | -10.6% | -12.6% | -11.7% | -12.0% |
| Dioxins | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.2% | 0.1% |
| HCB | 0.0% | 0.0% | 0.0% | 0.1% | 3.0% | 0.5% | 0.6% | 0.7% | 126.1% | 86.6% |
| PCB | 0.1% | 0.2% | 0.6% | 1.6% | 7.4% | 1.0% | 1.1% | 1.4% | 128.5% | 90.7% |
| As | -45.9% | -58.7% | -64.3% | -65.6% | -71.5% | -68.9% | -67.3% | -68.2% | -66.7% | -68.1% |
| Cd | -18.5% | -27.2% | -30.3% | -32.0% | -38.5% | -34.5% | -30.3% | -34.5% | -32.6% | -35.1% |
| Cr | -2.9% | -5.1% | -7.8% | -8.0% | -24.3% | -22.2% | -21.4% | -22.2% | -20.1% | -25.8% |
| Cu | -21.6% | -33.7% | -43.0% | -49.8% | -78.6% | -71.4% | -74.0% | -72.8% | -69.6% | -73.5% |
| Hg | 25.3% | 36.1% | 85.7% | 98.0% | 2008% | 203.1% | 906.4% | 264.7% | 253.6% | 150.6% |
| Ni | 0.4% | 0.7% | 1.1% | 3.2% | 538.7% | 229.4% | 544.4% | 263.5% | 259.6% | 182.5% |
| Pb | -22.8% | -31.6% | -46.8% | -40.9% | -91.3% | -92.9% | -92.9% | -93.0% | -92.8% | -92.9% |
| Se | -61.5% | -71.7% | -74.0% | -75.5% | -73.2% | -71.4% | -68.3% | -70.2% | -68.5% | -68.5% |
| Zn | -86.8% | -89.9% | -91.8% | -91.2% | -93.7% | -93.6% | -93.4% | -93.6% | -93.4% | -93.5% |
| Wastewater (5D) | | | | | | | | | | |
| NMVOG | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% | 0.0% |
| Other waste (5E) | | | | | | | | | | |
| PM2.5 | 636.1% | 636.1% | 636.1% | 636.1% | 668.6% | 611.0% | 642.3% | 545.7% | 705.5% | 0.0% |
| BC | 636.1% | 636.1% | 636.1% | 636.1% | 668.6% | 611.0% | 642.3% | 545.7% | 705.5% | 0.0% |
| PCDD/F (g I-Teq) | 550.3% | 550.3% | 550.3% | 550.3% | 577.5% | 544.6% | 576.8% | 498.6% | 631.5% | 670.9% |

Recently, in the previous submission, estimates of PM emissions have been included in the inventory for solid waste disposed on landfills. For this category, some minor recalculations occurred because of the update of sludge activity data since 2007 while the main methodological update is due to the new waste characterization applied starting from 2016.

About biological treatment, recalculations occur in 2019 because of the update of activity data for mechanical biological treatments. Furthermore, the formula for calculating the biogas produced has been corrected, which affects the estimate of N at digesters the whole time series.

As regards incineration, recalculations occur for the update of activity data from 2018. As concern open burning: correction of the amount of biomass available for open burning since 1990 and addition of the estimation of Ni and Hg emissions for open burning from combustion of removable cereal residues. Recalculations in the category 5E occurred because of the last review process and the consequent technical correction.

The analysis regarding incineration plants has been conducted through verifications and comparisons with data reported in E-PRTR registry, Emissions Trading Scheme and updated data of incinerated waste amount by plants.

For what concerns cremation of corpses, in 2020 SEFIT, the specific branch of Federutility, which is the federation of energy and water companies which is responsible for crematoria, has carried out together with ISPRA a new survey on emissions from national crematoria. Results from this survey have been utilized for the new estimations taking in account the implementation of abatement technology over the years.

7.5 PLANNED IMPROVEMENTS

Emissions from 5E are still under investigation, the results will be reported in the next submissions.

8 RECALCULATIONS AND IMPROVEMENTS

8.1 RECALCULATIONS

To meet the requirements of transparency, consistency, comparability, completeness and accuracy of the inventory, the entire time series is checked and revised every year during the annual compilation of the inventory. Measures to guarantee and improve these qualifications are undertaken and recalculations should be considered as a contribution to the overall improvement of the inventory.

Recalculations are elaborated on account of changes in the methodologies used to carry out emission estimates, changes due to different allocation of emissions as compared to previous submissions, changes due to error corrections and in consideration of new available information.

The complete NFR files from 1990 to 2020 have been submitted. The percentage difference between the time series reported in the 2021 submission and the series reported this year (2022 submission) are shown in Table 8.1 by pollutant.

Improvements in the calculation of emission estimates have led to a recalculation of the entire time series of the national inventory. Considering the total emissions, the emission levels for the year 2019 show a decrease for some pollutants and an increase for others; in particular, according to the review process, a significant decrease has been observed for lead emissions, equal to 11.3% , for cadmium (-6.1%), mercury (-1.7%) and PAH (-0.03%) mainly due to the update of emission factors for iron and steel activities on the basis of plant specific data, and the update of emission factor for open burning of agriculture residues and for PM 2.5 (-1.1%) and BC (-2.4%) for the update of emission factors for iron and steel activities. On the other side a significant increase of emissions has been observed in 2019 for HCB (36.1%) and PCB (3.9%) due to the estimation of emissions from secondary aluminium production and for Dioxins (13.4%) and PM10 (2.4%), due to the update according to the review process of emissions from building and cars fires. The update of COPERT version, resulted in a recalculation of NO_x (+2.0%) and NH₃ (-1.6%). The decrease of NMVOC emissions (-0.7%) and CO (-0.04%) is driven by the update of emission factor for open burning of agriculture residues. The decrease of SO_x emissions (-0.4%) is driven by from one side the decrease of emissions due to the update of diesel fuel consumption for agriculture off-road and fishing, from the other side by the increase of emissions due to the update of the sulphur content of biodiesel for road transport. For the other pollutants recalculations for 2019 are less than 1%.

In the *energy* sector a further revision of the emission estimates regarded the road transport sector. Specifically, the upgraded version of COPERT model, COPERT 5.5.1 (EMISIA SA, 2021), has been applied to calculate emissions of all pollutants for the whole period 1990-2020, with a revision of emission factors especially for the new vehicles. It resulted in a recalculation of the time series for all the pollutants. For 1.A.2 sector a revision of emission factors on the basis of EPRTR plant level data involved SO_x, NO_x and NH₃ for glass and lime production, and on the basis of plant data SO_x, NO_x, PM, Heavy metals and HCB for the iron and steel sector. Moreover HCB and PCB emissions from secondary aluminum production has been included for the whole timeseries.

For 1.A.3.a category, aviation, there has been an update for the entire time series of PAH from 1990 to 2017 due to the correction of an error in the calculation of the emissions. For 1.A.3.d maritime activities, the total number of ships arrived has been updated from 2017.

For 1.A.4 energy recovery from waste reported in the commercial heating has been updated from 2018 because of the update of activity data.

In the *industrial processes* sector, recalculations occur for PM10, PM2.5 along the whole time series because this submission includes PM emission estimates from category 2.A.5b construction and demolition according to the recommendation of the review (EEA, 2021). Minor recalculations occurred because of the update of activity data for the following source categories: chipboard, bread, glass, and lime production.

For the *solvent* sector minor recalculations occurred because of the update of activity data in the manufacture of fat edible, in 2019.

In 2022, recalculations were implemented for the *agricultural* emission inventory. Major changes are: corrected EF NH₃ housing for 2019 for non-dairy cattle and buffaloes categories; included in storage estimates NH₃ emission factors for equine and sheep and goat for the whole time series; modified the formula for estimating NH₃ emissions from storage for cattle, pigs and poultry by subtracting from N at housing also the amount of N at digesters before multiplying by the emission factor; for the estimation of N at spreading, the percentages of N remaining after emissions in the housing, storage and after other losses during storage (NO₂,

N₂O, N₂, N leached in manure management, NH₃ from digesters) were recalculated and N bedding was added; updated PM emission factors of the other poultry category for the whole time series; corrected the formula for calculating the biogas produced, which affects the estimate of N at digesters the whole time series for dairy cattle and non-dairy cattle and, to a small extent, from 1991 to 2004 also for pigs; updated the estimate of straw use for dairy cattle, non-dairy cattle and buffaloes to take into account sheltered animals only (removing grazing animals).

As regards the *waste* sector, recalculations occur for the category 5C, because of the update of activity data for industrial waste incineration. Recalculations in the category 5E occurred because of the last review process and the consequent technical correction. As concern open burning: correction of the amount of biomass available for open burning since 1990 and addition of the estimation of Ni and Hg emissions for open burning from combustion of removable cereal residues. About biological treatment, recalculations occur in 2019 because of the update of activity data for mechanical biological treatments. Furthermore, the formula for calculating the biogas produced has been corrected, which affects the estimate of N at digesters the whole time series. For what concerns cremation of corpses, in 2020 SEFIT, the specific branch of Federutility, which is the federation of energy and water companies which is responsible for crematoria, has carried out together with ISPRA a new survey on emissions from national crematoria.

Table 8.1 Recalculation between 2021 and 2022 submissions

| | SO _x | NO _x | NH ₃ | NMVOC | CO | PM10 | PM2.5 | BC | Pb | Hg | Cd | DIOX | PAH | HCB | PCB |
|------|-----------------|-----------------|-----------------|-------|-------|------|-------|-------|--------|-------|-------|-------|-------|-------|------|
| | % | | | | | | | | | | | | | | |
| 1990 | 0.00 | -0.01 | 0.53 | -0.02 | 0.00 | 3.27 | 1.29 | 0.90 | -0.04 | 0.55 | -1.15 | 5.24 | -0.16 | 2.13 | 1.73 |
| 1991 | 0.00 | -0.01 | 0.53 | -0.01 | 0.00 | 3.16 | 1.22 | 0.88 | -0.09 | 0.67 | -2.00 | 5.05 | -0.26 | 2.19 | 1.70 |
| 1992 | 0.00 | -0.01 | 0.54 | -0.01 | 0.00 | 3.10 | 1.23 | 0.88 | -0.10 | 0.64 | -1.74 | 5.28 | -0.23 | 2.38 | 1.82 |
| 1993 | 0.00 | -0.01 | 0.55 | -0.01 | 0.00 | 2.77 | 1.18 | 0.85 | -0.11 | 0.66 | -1.91 | 5.57 | -0.24 | 2.45 | 1.71 |
| 1994 | 0.05 | -0.01 | 0.59 | -0.02 | 0.00 | 2.87 | 1.21 | 0.87 | -0.09 | 0.73 | -1.82 | 5.69 | -0.22 | 2.81 | 1.86 |
| 1995 | 0.03 | -0.01 | 0.59 | -0.02 | 0.00 | 3.23 | 1.26 | 0.86 | -0.11 | 0.72 | -1.97 | 5.45 | -0.23 | 3.16 | 1.83 |
| 1996 | 0.03 | -0.01 | 0.61 | -0.02 | 0.00 | 3.18 | 1.29 | 0.87 | -0.10 | 0.80 | -1.73 | 5.88 | -0.22 | 3.67 | 1.93 |
| 1997 | 0.00 | -0.02 | 0.59 | -0.02 | 0.00 | 2.96 | 1.27 | 0.88 | -0.15 | 0.75 | -2.05 | 5.80 | -0.23 | 3.40 | 2.15 |
| 1998 | 0.01 | -0.02 | 0.60 | -0.02 | 0.00 | 3.01 | 1.28 | 0.84 | -0.14 | 0.86 | -1.86 | 5.90 | -0.22 | 3.07 | 2.53 |
| 1999 | 0.01 | -0.02 | 0.60 | -0.02 | 0.00 | 3.26 | 1.34 | 0.86 | -0.18 | 0.93 | -2.37 | 6.21 | -0.26 | 8.72 | 2.52 |
| 2000 | 0.01 | -0.02 | 0.69 | -0.02 | 0.00 | 3.62 | 1.47 | 0.92 | -0.25 | 0.87 | -1.95 | 6.54 | -0.34 | 18.19 | 2.95 |
| 2001 | 0.01 | -0.01 | 0.54 | -0.03 | 0.00 | 3.69 | 1.55 | 0.96 | -0.35 | 0.91 | -2.15 | 8.28 | -0.38 | 13.63 | 2.83 |
| 2002 | 0.02 | -0.01 | 0.54 | -0.03 | 0.00 | 4.69 | 1.94 | 1.07 | -0.94 | 1.00 | -2.43 | 9.08 | -0.48 | 15.32 | 3.03 |
| 2003 | 0.04 | -0.01 | 0.55 | -0.03 | 0.00 | 4.05 | 1.73 | 1.06 | -0.92 | 0.98 | -2.57 | 8.45 | -0.37 | 15.03 | 2.86 |
| 2004 | 0.05 | -0.01 | 0.54 | -0.03 | 0.00 | 4.77 | 2.06 | 1.15 | -1.09 | 1.32 | -3.08 | 9.20 | -0.55 | 20.93 | 2.91 |
| 2005 | 0.00 | -0.01 | 0.47 | -0.04 | 0.00 | 4.02 | 1.77 | 1.16 | -0.93 | 1.37 | -2.67 | 8.10 | -0.36 | 25.77 | 2.83 |
| 2006 | 0.00 | -0.00 | 0.37 | -0.05 | 0.00 | 3.63 | 1.50 | 1.03 | -0.89 | 1.35 | -2.72 | 6.59 | -0.31 | 23.71 | 2.80 |
| 2007 | -0.02 | -0.01 | 0.34 | -0.06 | 0.00 | 3.17 | 1.31 | 0.92 | -0.87 | 1.61 | -2.64 | 5.97 | -0.24 | 28.05 | 2.92 |
| 2008 | 0.01 | 0.01 | 0.42 | -0.09 | 0.00 | 2.73 | 1.19 | 0.86 | -0.95 | 1.97 | -3.07 | 5.88 | -0.22 | 30.01 | 2.72 |
| 2009 | 0.01 | 0.05 | 0.46 | -0.10 | 0.00 | 2.49 | 1.22 | 0.86 | -1.29 | 2.74 | -3.65 | 7.51 | -0.24 | 29.64 | 3.15 |
| 2010 | 0.04 | 0.06 | 0.53 | -0.10 | 0.00 | 2.32 | 1.21 | 0.80 | -1.15 | 2.82 | -3.81 | 7.57 | -0.24 | 30.49 | 3.55 |
| 2011 | -0.03 | 0.02 | 0.12 | -0.13 | 0.00 | 2.95 | 1.76 | 0.96 | -1.10 | 1.08 | -3.86 | 10.01 | -0.32 | 37.94 | 3.97 |
| 2012 | -0.55 | -0.05 | -0.50 | 0.11 | 0.61 | 2.26 | 1.37 | 0.82 | -1.03 | 1.01 | -3.71 | 9.46 | -0.22 | 29.53 | 3.47 |
| 2013 | -0.63 | -0.31 | -1.64 | -0.29 | -0.07 | 1.95 | 1.30 | 0.66 | -1.16 | 1.17 | -4.19 | 9.63 | -0.25 | 38.15 | 4.19 |
| 2014 | -0.82 | -0.38 | -2.04 | -0.39 | -0.16 | 2.03 | 1.39 | 0.58 | -0.95 | 1.07 | -3.34 | 10.30 | -0.24 | 40.56 | 4.76 |
| 2015 | -0.77 | -0.41 | -1.91 | -0.40 | -0.15 | 2.00 | 1.37 | 0.64 | -1.18 | 1.32 | -4.16 | 10.08 | -0.28 | 35.94 | 4.90 |
| 2016 | 0.55 | 0.21 | -1.85 | -0.41 | -0.13 | 1.78 | 1.22 | 1.24 | -14.52 | -2.31 | -6.37 | 11.44 | -0.24 | 36.15 | 4.75 |
| 2017 | -0.33 | 1.95 | -1.96 | -0.39 | -0.12 | 1.94 | 1.27 | 1.35 | -12.17 | -1.63 | -6.14 | 11.44 | -0.07 | 41.16 | 4.79 |
| 2018 | 0.20 | 3.10 | -2.03 | -0.34 | -0.09 | 2.34 | 1.53 | 1.30 | -11.05 | -1.41 | -5.76 | 12.32 | 0.01 | 41.55 | 4.43 |
| 2019 | -0.44 | 2.01 | -1.55 | -0.74 | -0.04 | 2.35 | -1.06 | -2.42 | -11.31 | -1.66 | -6.11 | 13.36 | -0.03 | 36.09 | 3.96 |

8.2 PLANNED IMPROVEMENTS

Specific improvements are specified in the QA/QC plan (ISPRA, 2022[b]); they can be summarized as follows.

For the *energy* and *industrial processes* sectors, a major progress regards the harmonisation of information collected in the framework of different obligations, Large Combustion Plant, E-PRTR and Emissions Trading, thus highlighting the main discrepancies in data and detecting potential errors, the use of data and country specific emission factors collected in national research involving road transport and biomass consumption in residential and for POPs emissions the use of the results of a national research in the potential update of emission factors and methodologies in the iron and steel sector. For the *agriculture* and *waste* sectors, improvements will be related to the availability of new information, on emission factors, activity data as well as parameters necessary to carry out the estimates; specifically, a study on the best available technologies used in agriculture practices and the elaboration of data from the 2016 farm structure survey and availability of information on the landfill gas combustion in landfills flaring and emissions from the exceeding biogas flared at wastewater treatment plants are under investigation.

The EMEP/EEA Guidebook 2019 chapters (EMEP/EEA, 2019) has continued to be considered, and update emission factors will be applied in the next year submission of the inventory with a focus to PAH estimates in order to improve the completeness, e.g. for PAH compounds, accuracy and reduce the uncertainty.

The comparison between local inventories and national inventory and the meetings and exchange of information with local environmental agencies will continue.

Further analyses will concern the collection of statistical data and information to estimate uncertainty in specific sectors.

9 PROJECTIONS

In 2021 submission, the pollutant emission projections presented in this chapter were consistent with the emission projections submitted at EU level according to the Directive (EU) 2016/2284 of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants.

After the review process carried out on those projections, a re-submission was provided in June 2021.

In the following months a major revision of data and methodologies has been started at national level, in order to improve the overall quality of projections and to take into account the changes in activity levels due to the COVID-19 pandemic. This process is not concluded yet.

For this reason, the present chapter is updated only to take into account the re-submission made in June 2021. Next year submission will present new projections and address the improvements required by the Technical Review Team.

9.1 THE NATIONAL FRAMEWORK

At national level, the Legislative Decree n. 155 of 2010 (D.Lgs. 2010), that implements the European Directive on air quality, 2008/50/EC (EC, 2008), and the Legislative Decree n. 81 of 2018 (D.Lgs. 2018), that implements the Directive (EU) 2016/2284, provide that ISPRA develops the energy scenario and the scenario of national production activities while ENEA, based on these scenarios, calculates the emission projections using the methodology developed for these purposes at European level.

In this framework, ENEA has elaborated the new national baseline emission scenario using the GAINS-Italy model.

GAINS-Italy is part of the MINNI model, an Integrated Modelling System that links atmospheric science with the economics of emission abatement measures and policy analysis and consists of several interdependent and interconnected components: the national AMS (Atmospheric Modeling System, Mircea et al., 2014) and the national GAINS-Italy. They interact in a feedback system through ATMs (Atmospheric Transfer Matrices) and RAIL (RAINS-Atmospheric Inventory link).

The GAINS-Italy model (fig. 9.1) explores cost-effective multi-pollutant emission control strategies (Ciucci et al., 2016) that meet environmental objectives on air quality impacts (on human health and ecosystems) and greenhouse gases. The current legislation (CLE) scenario represents the ‘baseline’ and reflects all policies legally in force, both those affecting activity levels (such as energy and agriculture policies), as well as pollution control policies for the period 1990-2050.

The GAINS-Italy model elaborates emission scenarios for air pollutants and greenhouse gases on 5-year time intervals, starting from 1990 to 2050, and evaluates cost-effective multi-pollutant emission control strategies to reach environmental objectives on air quality impacts. Moreover, GAINS-Italy performs fast-response calculations of regional background concentrations of PM_{2.5} and NO₂ in consequence of hypothesized emission reductions on the Italian territory. This last feature is enhanced by the Atmospheric Transfer Matrices (ATMs), simplified (quasi-linear) relations between total regional emissions and concentrations, calibrated through a set of national Atmospheric Modelling System simulations, based on controlled pollutant emission reductions (Briganti et al., 2011).

The development of an emission scenario with the GAINS-Italy model requires the definition of anthropogenic activity levels, both energy and non-energy, and of a control strategy with a 5-year interval for the period 1990-2050 in the format required by the model. Starting from these information, GAINS-Italy produces alternative future emission and air quality scenarios and abatement costs at a 5-year interval starting from 1990 to 2050.

For the preparation of national emission scenarios, an acceptable harmonization, at a given base year, between the national emission inventory and the GAINS-Italy emissions (D’Elia and Peschi, 2013) has been carried out.

More details about the procedure to build an emission scenario could be found in D’Elia and Peschi, 2016. In the present chapter, the scenarios that will be discussed are the Baseline scenario (with measures, WM_NECP) and the Policy scenario (with additional measures, WAM_NECP) elaborated for the National Energy and Climate Plan (NECP, https://ec.europa.eu/energy/sites/ener/files/documents/it_final_necp_main_en.pdf).




Figure 9.1 – The simplified functional flow-chart of the MINNI national modelling scheme.

9.2 INPUT SCENARIOS

9.2.1 The energy scenario

The Energy scenario used as input to the GAINS-Italy model has been produced by ISPRA with the TIMES (The Integrated MARKAL-EFOM1 System / EFOM Energy Flow Optimization Model, Loulou et al., 2004; Loulou et al., 2005) model developed as part of the IEA-ETSAP (Energy Technology Systems Analysis Program). TIMES is a technology rich, bottom-up model generator, which uses linear-programming to produce a least-cost energy system, optimized according to a number of user constraints, over medium to long-term time horizons.

The model has been developed considering the detailed energy input needed by GAINS-Italy so that the two models are fully integrated and all the information needed by GAINS-Italy can be found in the TIMES output, that describes, for each sector, the amount of energy carriers, raw materials used, and goods or services produced.

In 2020 the National Energy and Climate Plan (NECP) has been submitted to the European Commission, the scenarios used as input to the GAINS model are consistent with the scenarios included in the NECP. Scenario are also consistent with those submitted to the EU Commission under regulation 525/2013 and available at http://cdr.eionet.europa.eu/it/eu/mmr/art04-13-14_lcds_pams_projections/envvxfh1a/.

9.2.2 The scenario of non-energy activities

To develop an emission scenario, the GAINS-Italy model requires the definition also of non-energy activities level. The definition of such scenario is based on economic variables, like GDP (gross domestic product) or added value derived from the energy scenario, population data or specific sector statistics.

Livestock projection has been carried out with a statistical model where the number of animals has been linked to the projections of other variables, like meat consumption and production, or milk consumption and production (see equation 9.1)

$$(n^{\circ}heads_i) = \left(\frac{n^{\circ}heads_i}{MP_i}\right) \times \left(\frac{MP_i}{MC_i}\right) \times \left(\frac{MC_i}{MC_{tot}}\right) \times \left(\frac{MC_{tot}}{Pop}\right) \times (Pop) \quad (9.1)$$

where the head number of livestock i is linked to meat production (MP) and consumption (MC) of livestock i , total meat consumption (MC_{tot}) and population (Pop).

All the details about this methodology are provided in D'Elia and Peschi, 2013.

The updated livestock projections have been elaborated for the baseline WM_NECP: population data (Pop) are the same of the NECP energy scenario, while for meat production and consumption sectorial studies and statistics have been considered (<http://www.ismeamercati.it/carni>).

In the following figures, the results for the main livestock are reported where ISPRA stands for the activity data considered in the 2020 emission submission, that was the last submission available when the livestock projections were elaborated;

WM_NECP are the new projections elaborated for the WM_NECP scenario.




Figure 9.2 – Livestock scenario comparison for dairy cows and other cattle.




Figure 9.3 – Livestock scenario comparison for pigs and sheep and goats.

9.2.3 The control strategy definition

In addition to energy, climate and agricultural policies assumed in the energy, non-energy and agricultural input scenarios, in the baseline emission projections a detailed inventory of national emission control legislation is considered (Amann et al., 2011).

In the WM_NECP scenario it is assumed that all the European and national regulations adopted before 2020 will be fully complied according to the foreseen time schedule. Examples of the legislations considered are the Directive on Industrial Emissions for large combustion plants, the Directives on Euro standards, Solvent Directive, the Code of Agricultural Good Practice.

In the WAM_NECP scenario it is assumed that all the policies and targets set out in the NECP will be adopted. An example of the WM_NECP control strategy for the road transport sector is reported in the following figures.




Figure 9.4 – Control strategy for gasoline (on the left) and diesel (on the right) cars.




Figure 9.5 – Control strategy for diesel light (on the left) and heavy (on the right) duty vehicles.

9.3 THE HARMONIZATION PROCESS

The first step for the preparation of a new national emission scenario is to align at a given base year the latest national emission inventory submission and the GAINS-Italy emissions, estimated with a top-down approach. Being a Party of the United Nations Economic Commission for Europe (UNECE) Convention on Long Range Transboundary Air Pollution (CLRTAP), Italy has to annually submit an emission inventory of air pollutants and provide a report on its data according to the Guidelines for reporting emissions and projections data (UNECE, 2015). On the other hand, to produce a reliable emission scenario, GAINS-Italy model produces its own emission estimates, for the years considered in the model, with its own classification system. Discrepancies between the inventory and the GAINS-Italy output exist and are due to different reasons, such as, for example, different coverage and aggregation of emission sources, different emission calculation methodologies. These discrepancies need so to be solved and the emission estimates to be aligned. This alignment step is called harmonization and is needed to validate the emission scenario to base emission time trends in GAINS-Italy on a reliable starting point. In the harmonization process, activity data, emission factors and technologies for each sector are compared. If discrepancies emerge (for example in fuel allocation across sectors or different assumptions on control measures in place in the year of comparison), the model parameters will be modified according to the inventory with the attempt to let GAINS-Italy reproduce emissions as closely as possible to the national emission inventory. Further details about the harmonization method are reported in D’Elia and Peschi, 2013. For all these reasons, a comparison between the last national emission inventory, and the GAINS-Italy emission estimates has been carried out considering three historical years, 2005, 2010 and 2015. Results of the harmonization process between the 2021 emission inventory submission (INVENTORY_sub2021) and GAINS-IT estimates are summarized in Table 9.1.

Table 9.1 – Comparison of total emissions in the last submission of the national inventory report (INVENTORY_sub2021) and GAINS-Italy estimates (GAINS-IT), for the years 2005, 2010 and 2015.

| Pollutant | Emissions 2005 | | | Emissions 2010 | | | Emissions 2015 | | |
|-----------------|------------------------|---------------|-------|------------------------|---------------|-------|------------------------|---------------|-------|
| | Inventory_sub2021 (kt) | GAINS-IT (kt) | Δ (%) | Inventory_sub2021 (kt) | GAINS-IT (kt) | Δ (%) | Inventory_sub2021 (kt) | GAINS-IT (kt) | Δ (%) |
| SO ₂ | 411 | 393 | -4% | 222 | 209 | -6% | 127 | 121 | -5% |
| NO _x | 1289 | 1231 | -4% | 934 | 985 | 5% | 719 | 719 | 0% |
| PM2.5 | 173 | 176 | 2% | 196 | 196 | 0% | 158 | 165 | 4% |
| NM VOC | 1340 | 1232 | -8% | 1117 | 993 | -11% | 901 | 803 | -11% |
| NH ₃ | 419 | 428 | 2% | 377 | 394 | 5% | 364 | 385 | 6% |

Discrepancies in reproducing the national emission inventory have been considered acceptable if differences remain within a few percentage points, i.e. in the interval between $\pm 5\%$. The higher differences in NMVOC emissions for all the three years depend on the estimates of emissions from the agricultural sector (NFR code 3) that in the model estimates have not yet been considered but they will be introduced in the emission projection update that is underway.

In the following plots, details on sectoral emissions by NFR code (Nomenclature For Reporting; Table 9.2 reports the sectors considered) are illustrated for the year 2015.

Table 9.2 – Definition of the NFR code used in the comparison between emission inventory and GAINS-IT estimates.

| NFR code | Description |
|----------|--|
| 1A1 | Energy industries (Combustion in power plants & Energy Production) |
| 1A2 | Manufacturing Industries and Construction (Combustion in industry including Mobile) |
| 1A3b | Road and Off-road Transport |
| 1A4 | Other sectors (Commercial, institutional, residential, agriculture and fishing stationary and mobile combustion) |
| 1A5 | Other |
| 1B | Fugitive emissions (Fugitive emissions from fuels) |
| 2 | Industrial Processes and Solvent use |
| 3 | Agriculture |
| 5 | Waste |
| 6A | Other (included in National Total for Entire Territory) |

For SO₂ emissions, the model shows a slight overestimate of the sectors 1A1 and 2, an underestimate of the sectors 1A2 and 1A4, while for NO_x there is a slight overestimate of the sector 1A3. Emissions from the agriculture sector (code 3) both for NO_x and NMVOC are missing in GAINS-IT and will be added in future updates.

PM_{2.5} shows a good agreement between the two estimates with the exception of the sector 6 – Other where the model estimates emissions from barbecue, fireworks not estimated in the inventory for their high uncertainties.

The model shows a slight overestimate in NMVOC emissions from the sector 2 – Industrial process and solvent use and in NH₃ emissions from the sector 3 – Agriculture in the range of uncertainty considered acceptable.




Figure 9.6 – SO₂ national emission harmonization between the last emission inventory (INVENTORY_sub2021) and GAINS-IT estimates detailed by NFR sectors for the year 2015.




Figure 9.7 – NO_x national emission harmonization between the last emission inventory (INVENTORY_sub2021) and GAINS-IT estimates detailed by NFR sectors for the year 2015.




Figure 9.8 – PM2.5 national emission harmonization between the last emission inventory (INVENTORY_sub2021) and GAINS-IT estimates detailed by NFR sectors for the year 2015.




Figure 9.9 – NMVOC national emission harmonization between the last emission inventory (*INVENTORY_sub2021*) and *GAINS-IT* estimates detailed by NFR sectors for the year 2015.




Figure 9.10 – NH₃ national emission harmonization between the last emission inventory (*INVENTORY_sub2021*) and *GAINS-IT* estimates detailed by NFR sectors for the year 2015.

9.4 THE EMISSION SCENARIO

The result of the activity input scenarios and of the harmonization process is an emission scenario. In the following figures, a comparison of the 2021 emission inventory submissions, the WM_NECP and WAM_NECP scenarios, used for NECP, are presented. Details by NFR sector are presented for the WAM_NECP; the WM_NECP scenario description was discussed in the (IIR, 2020).

A huge decrease in SO₂ emissions is projected (fig. 9.11) driven by the energy and the maritime sector for the year 2030 while the industrial sector (1A2) in the year 2030 represents the main emitting sector (fig. 9.12).




Figure 9.11 –Reported and projected (WM_NECP, WAM_NECP) SO₂ emissions elaborated by the GAINS-Italy model.




Figure 9.12 –SO₂ emissions by NFR sectors in the WAM_NECP scenario.

A huge decrease is estimated in NO_x emission scenarios (fig. 9.13) due to the diffusion of new diesel Euro 6 and electric vehicles. The road transport sector still represents the principle NO_x source (fig. 9.14).




Figure 9.13 – Reported and projected (WM_NECP, WAM_NECP) NO_x emissions elaborated by the GAINS-Italy model.




Figure 9.14 – NO_x emissions by NFR sectors in the WAM_NECP scenario.

The decrease of the PM2.5 WAM_NECP scenario (fig. 9.15) is driven by the civil sector (1A4) that continues to represent the main emitting sector (fig. 9.16).




Figure 9.15 –Reported and projected (WM_NECP, WAM_NECP) PM2.5 emissions elaborated by the GAINS-Italy model.




Figure 9.16 – PM2.5 emissions by NFR sectors in the WAM_NECP scenario.

As already underlined, the gap between the reported and projected NMVOC emissions (fig. 9.17) is due to the estimate of the sectors 3B and 3D, that will be solved in future submissions. The solvent sector will remain the main emitting sector (fig. 9.18).




Figure 9.17 –Reported and projected (WM_NECP, WAM_NECP) NMVOC emissions elaborated by the GAINS-Italy model.




Figure 9.18 – NMVOC emissions by NFR sectors in the WAM_NECP scenario.

NH₃ is the pollutant with less variations (fig. 9.19) whose main contribution to total NH₃ emissions is due by the agricultural sector (fig. 9.21).




Figure 9.19 –Reported and projected (WM_NECP, WAM_NECP) NH₃ emissions elaborated by the GAINS-Italy model.

9.5 THE NEC EMISSION TARGETS

The NEC Directive (EU, 2016), implemented in the Italian legislation in the D.Lgs. 81/2018, defines for each Member States the emission reduction targets in the year 2020 and 2030 respect to the base year 2005 for the anthropogenic emissions of SO₂, NO_x, PM_{2.5}, NMVOC and NH₃.

In Table 9.4 the attainment of the national emission reductions in the year 2030 in the comparison with the new National Emission Ceilings Directive (NECD) targets is reported.

Table 9.3 – National emission reductions in the year 2030 respect to the base year 2005 and comparison with the new National Emission Ceilings Directive (NECD) targets.

| | 2030 EMISSION REDUCTIONS | | |
|-------------------|--------------------------|---------|----------|
| | NECD targets | WM_NECP | WAM_NECP |
| SO ₂ | -71% | -78% | -80% |
| NO _x | -65% | -68% | -71% |
| PM _{2.5} | -40% | -31% | -43% |
| NMVOC | -46% | -49% | -51% |
| NH ₃ | -16% | -13% | -17% |

According to the present emission projections, additional measures have to be adopted for the 2030 targets. The WAM_NECP scenario is in compliance with all the targets.

10 REPORTING OF GRIDDED EMISSIONS AND LPS

Every four years, from 2017 with reference to 2015 emissions, ISPRA shall provide the disaggregation of the national inventory at provincial level as instituted by the Legislative Decree n. 81 of 30 May 2018. The emissions disaggregated at regional and provincial levels are compared to the results obtained by regional bottom-up inventories. Emissions disaggregated at local level are also used as input for air quality modelling. Methodologies and proxies are described in the relevant publication (ISPRA, 2009).

Simultaneously, ISPRA carries out the spatial disaggregation of national emissions on the 0.1x0.1° EMEP grid.

The grid definition for a country contains a dataset for each grid cell with information about (CEIP, 2019):

- the country or area (ISO2 or a three digits abbreviation for countries and other areas which you can see in the column Country code in the grid definition table above)
- the country-or area name
- the longitude position of the grid cell (centre of the cell)
- the latitude position of the grid cell (centre of the cell)
- and fraction of the grid cell (share of the cell area which belongs to the country/area, e.g. 1 for cells which are completely inside the country/area borders and e.g. 0.5 for cells where half of the area belongs to the country/area and the rest is outside the boundary)

According to the review process (EEA, 2020), the reporting of gridded data has been aligned with the last available reporting guidelines and the relevant GNFR codes in the next submission, as it is indicated in the Table 10.1. Proxy used for disaggregation have been reported too.

Table 10.1 Aggregation for Gridding and LPS of NFR sector

| NFR Aggregation for Gridding and LPS (GNFR) | Proxy |
|---|---|
| A_PublicPower | point sources consumptions from ETS/LPS |
| B_Industry | combustion in industry prevalently point sources consumptions data from ETS/LPS, where not available production data. For industrial processes prevalently production data from E-PRTR or producers associations, in a few of case production capacity or data about employees at NUTS3 level |
| C_OtherStationaryComb | resident people and fuels sold at NUTS3 level |
| D_Fugitive | see paragraph "fugitive" |
| E_Solvents | prevalently employees at NUTS3 level. Data from the national institut of statistics (ISTAT) |
| F_RoadTransport | see paragraph "transport" |
| G_Shipping | see paragraph "transport" |
| H_Aviation | see paragraph "transport" |
| I_Offroad | see paragraph "transport" |
| J_Waste | amount of managed waste at NUTS3 level on the basis of the management system for SWDS, biological treatments and incineration. Source: ISPRA. For domestic wastewater resident people at Nuts3 level while data about |

| | |
|-----------------|--|
| | employees at NUTS3 level in the case of industrial wastewater have been used. Source: ISPRA. |
| K_AgriLivestock | see paragraph "agriculture" |
| L_AgriOther | see paragraph "agriculture" |
| M_Other | |
| N_Natural | |
| O_AviCruise | see paragraph "transport" |
| P_IntShipping | see paragraph "transport" |
| z_Memo | |

The methodologies for spatial disaggregation are consistent with those reported in the EMEP/EEA 2019 Guidebook and described in relevant report (ISPRA, 2009). National emissions have been disaggregated at provincial level (NUTS3) because of the availability of proxy data, then the allocation to the 0.1x0.1° grid has been realized on the basis of the 2019 EMEP/EEA Guidebook (EMEP/EEA, 2019). In particular, point sources have been allocated directly to the grid within which they are contained by converting the x, y values to that of the coordinates used to geo-reference the grid. For area sources (NUTS3 polygons) the fraction of the area of the polygons has been used to distribute the emissions from the original polygon to the intersected grid cell. Finally, the spatially resolved (mapped) detailed NFR sectors have been aggregated in the GNFR code.

10.1 FUGITIVE

Proxy used for disaggregation of 1B1 and 1B2 are reported in the Table 10.2.

Tab. 10.2 Proxy for the disaggregation of fugitive emissions

| Category | Proxy |
|-------------------------|---|
| 1 B 1 a (SNAP 050102) | Amount of coal production |
| 1 B 1 a (SNAP 050103) | Amount of solid fuels consumption from ETS plants |
| 1 B 2 a iv | Amount of crude oil refined at plant level |
| 1 B 2 a v (SNAP 050502) | Amount of gasoline sold at provincial level. |
| 1 B 2 a v (SNAP 050503) | Amount of mineral oil and LPG in depots for industrial and services at regional level distributed according to the amount of gasoline sold at provincial level. |
| 1 B 2 b (SNAP 050601) | Amount of natural gas consumption in the compressor stations (ETS data) |
| 1 B 2 b (SNAP 050603) | Amount of natural gas provided to the distribution network at provincial level (other destination than industrial and thermoelectric users) |
| 1 B 2 c | Amount of crude oil refined at plant level |

10.2 TRANSPORT

Proxy used for disaggregation of the different transport mode are reported in the Table 10.3.

Tab. 10.3 Proxy for the disaggregation of transport emissions

| Category | Proxy |
|-----------|--|
| 1A2gvii | ISPRA study about industrial machinery data elaborated at NUTS 3 level. |
| 1A3ai(i) | Emissions deriving from international LTO, in the detail of Italian airports, elaborated by Eurocontrol to the aim of the estimation of emissions from aviation for Member Countries Inventories, then aggregated at NUTS 3 level. |
| 1A3aii(i) | Emissions deriving from domestic LTO, in the detail of Italian airports, elaborated by Eurocontrol to the aim of the estimation of emissions from aviation for Member Countries Inventories, then aggregated at NUTS 3 level. |
| 1A3bi | Ministry of Transport data about passenger cars fleet at NUTS 3 level, according to Copert classification; traffic flows and length of motorway sections as regards highway share. |
| 1A3bii | Ministry of Transport data about light duty vehicles fleet at NUTS 3 level, according to Copert classification; Value added as regards urban and rural share; traffic flows and length of motorway sections as regards highway share. |
| 1A3biii | Ministry of Transport data about vehicles fleet at NUTS 3 level, according to Copert classification; Value added as regards urban and rural share for heavy duty vehicles; traffic flows and length of motorway sections as regards highway share. |
| 1A3biv | Ministry of Transport data about mopeds and motorcycles fleet at NUTS 3 level, according to Copert classification; traffic flows and length of motorway sections for motorcycles categories as regards highway share. |
| 1A3bv | Same proxies as gasoline vehicles categories. Evaporative emissions are added to those due to combustion, for each vehicle category, according to Copert classification, for each driving cycle, therefore disaggregated at NUTS 3 level according to the criterion used for other road transport activities. Therefore emissions related to this activity are included in the other NFR classes. |
| 1A3bvi | Same proxies as corresponding vehicles categories, according to Copert classification. Non exhaust emissions are added to those due to combustion, for each vehicle category, according to Copert classification, for each driving cycle, therefore disaggregated at NUTS 3 level according to the criterion used for other road transport activities. Therefore emissions related to this activity are included in the other NFR classes. |
| 1A3c | Ministry of Transport data about the length of non-electrified railway sections, relating to Italian State Railways Group and to the Regional and/or local railways network, elaborated at NUTS 3 level. |
| 1A3dii | Data elaborated at NUTS 3 level covering: Eurostat data about vessels, by type, in harbours ; ships berths for: sailing boats, motorboats, watercrafts; simplified trajectories for cruise; inland waterways traffic is estimated by attributing a share to freight traffic (provinces of the Po basin) and a share to passenger traffic on the basis of Ministry of transport data about the number of boats for province. |
| 1A4bii | Data at NUTS 3 level about mechanical means used, deriving from surveys on the structure and production of farms, performed by the Italian Institute of Statistics. |
| 1A4cii | Data at NUTS 3 level about mechanical means used, deriving from surveys on the structure and production of farms, performed by the Italian Institute of Statistics. |
| 1A4ciii | Data, from Economic Observatory on the production structures of maritime fishing in Italy, about fishing boats number and consumptions at NUTS 3 level. |
| 1A5b | National Institute of Statistics data about resident population at NUTS 3 level have been used as proxy for military mobile activities. |

10.3 AGRICULTURE

Proxy used for disaggregation of the agriculture sector are reported in the Table 10.4.

Tab. 10.4 Proxy for the disaggregation of agriculture sector

| Category | Proxy |
|-----------------|--|
| 3Da1 | Quantity of annual nitrogen fertilizers (tons) distributed in the Italian provinces (National Institute of Statistics - ISTAT); for CO ₂ from liming: annual amount of lime and dolomite distributed (ISTAT); for CO ₂ from urea application: annual amount of urea distributed (ISTAT) |
| 3C | Cultivated area (hectares) for the production of rice (ISTAT) |
| 3Da2, 3Da3, 3De | For NH ₃ emissions from livestock, provincial emissions were estimated by multiplying the number of heads for each animal category available at provincial level (ISTAT) by the provincial emission factors by animal category. For N ₂ O and NO _x emissions from spreading and grazing, the proxies used are the provincial distributions of nitrogen excreted at the housing and grazing separately. |
| 3F | Annual production of cereal harvest (quintals) (ISTAT) |
| 3A1 | For cattle, provincial emissions were estimated by multiplying the number of heads available at provincial level (ISTAT) by the provincial emission factors from a 1994 study (Research Centre on Animal Production - CRPA). National emissions were then disaggregated on the basis of these provincial emissions |
| 3A | Number of heads available at provincial level (ISTAT) |
| 3B | <p>For NH₃ emissions, provincial emissions were estimated by multiplying the number of heads for each animal category available at provincial level (ISTAT) by the provincial emission factors by animal category from a 1994 study (Research Centre on Animal Production - CRPA). These provincial emissions for each animal category are the sum of emissions from all stages of manure management (i.e. housing, storage, spreading and grazing). National emissions for each animal category (which is the sum of emissions from all stages of manure management) were then disaggregated on the basis of these provincial emissions. For each animal category, the emission factors for each stage of manure management are considered. The percentage weight of the emission factors for each stage of manure management is calculated for each animal category. These percentages apply to the provincial total NH₃ emissions of all the stages, to calculate the provincial emissions (overall of all the animal categories) for the different stages distinctly. Finally, on the basis of these provincial emissions, the national emissions of each stages of manure management are disaggregated, separately.</p> <p>For CH₄ emissions, provincial emissions were estimated by multiplying the number of heads available at provincial level (ISTAT) by the provincial emission factors from a 1994 study (Research Centre on Animal Production - CRPA). National emissions were then disaggregated on the basis of these provincial emissions. For NMVOC, national emissions were disaggregated on the basis of CH₄ provincial emissions. For NO_x and PM emissions, national emissions were disaggregated on the basis of the number of heads available at provincial level (ISTAT)</p> |
| 3Df | Data on the provincial sale of pesticides containing HCB (ISPRA elaborations on ISTAT data) |
| 5C2 | Annual production of cereal and woody crops harvest (quintals) (ISTAT) |

10.4 LPS DATA

LPS data for the year 2019 have been submitted in 2021. A brief description of data is reported in the following Table 10.5.

Table 10.5 Characterization of LPS data

| GNFR | n° plant | Height class/n° |
|---------------|----------|-----------------|
| A_PublicPower | 207 | 2/194; 4/4; 5/9 |
| B_Industry | 330 | 1/328; 4/2 |
| D_Fugitive | 117 | 1/117 |
| G_Shipping | 77 | 1/77 |
| H_Aviation | 106 | 1/106 |
| I_Offroad | 22 | 1/22 |

Following the review process, some errors on longitude and latitude of plants have been corrected, also thanks the implementation procedures of the IED (Industrial Emissions Directive) European Directive.

Italy has decided not to provide all and only PRTR data as LPS data as in Italy some PRTR categories do not respond to the characteristics of large point sources. This is especially true for farms that cannot be considered large point sources and therefore have been eliminated with respect to the previous submission. The current submission is instead consistent with gridded data and includes sources outside of PRTR which are LPS such as ports or extractive activities.

11 REFERENCES

11.1 INTRODUCTION

- EMEP/CORINAIR, 2007. Atmospheric Emission Inventory Guidebook. Technical report No 16/2007.
- EMEP/EEA, 2009. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 9/2009.
- EMEP/EEA, 2013. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 12/2013.
- EMEP/EEA, 2016. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 21/2016.
- EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 13/2019.
- EU, 2016. National Emissions Ceilings Directive 2016/2284/EU.
- EEA, 2017 [a]. Final Review Report, 2017 Comprehensive Technical Review of National Emission Inventories pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Italy. 30 November 2017. Reference: No 07.0201/2016/741511/SER/ENV.C.3. Umweltbundesamt Wien.
- EEA, 2018. Final Review Report, 2018 Second phase of review of national air pollution emission inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284 or 'NECD'). Italy. 30 November 2018. Reference: No 070203/2017/765105/SER/ENV.C.3. Umweltbundesamt Wien.
- EEA, 2019. Final Review Report, 2019 Third phase of review of national air pollution emission inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284 or 'NECD'). Italy. 22 November 2019. Reference: Service request No 4 under Framework contract No ENV.C.3/FRA/2017/0012. International Institute for Applied Systems Analysis (IIASA) Laxenburg
- EEA, 2020. Final Review Report, 2020 Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 20 November 2020. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.
- EEA, 2021. Final Review Report, 2021 Review of National Air Pollutant Emission Inventory Data 2021 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 29 October 2021. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.
- IPCC, 1997. *Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories*. Three volumes: Reference Manual, Reporting Manual, Reporting Guidelines and Workbook. IPCC/OECD/IEA. IPCC WG1 Technical Support Unit, Hadley Centre, Meteorological Centre, Meteorological Office, Bracknell, UK.
- IPCC, 2000. Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan.
- IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan.
- ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009. URL: http://www.apat.gov.it/site/it-IT/APAT/Pubblicazioni/Rapporti/Documento/rapporti_92_2009.html.
- ISPRA, 2014. Quality Assurance/Quality Control plan for the Italian Emission Inventory. Procedures Manual. Report 112/2014. <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>.
- ISPRA, 2018. National Greenhouse Gas Inventory System in Italy. March 2018, <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>.
- ISPRA, 2022[a]. Italian Greenhouse Gas Inventory 1990-2019. National Inventory Report 2022. March 2022, <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>.
- ISPRA, 2022[b]. Quality Assurance/Quality Control plan for the Italian Emission Inventory. Year 2022. March 2022, <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>.
- Romano D., Bernetti A., De Lauretis R., 2004. Different methodologies to quantify uncertainties of air emissions. Environment International vol 30 pp 1099-1107.
- UNECE, 2008. Guidelines for Reporting Emission Data under the Convention on Long-range Transboundary Air Pollution, Emission Reporting Guidelines. ECE/EB.AIR/2008/4 23 September 2008.
- UNECE, 2010. Report for the Stage 3 in-depth review of emission inventories submitted under the UNECE LRTAP Convention and EU National Emissions Ceilings Directive for Italy. CEIP/S3.RR/2010/ITALY (25/11/2010).

UNECE, 2013. Report for the Stage 3 in-depth review of emission inventories submitted under the UNECE LRTAP Convention and EU National Emissions Ceilings Directive for Italy. CEIP/S3.RR/2013/ITALY (20/08/2013)

UNECE, 2014. Guidelines for Reporting Emissions and Projections Data under the Convention on Long Range Transboundary Air Pollution, ECE/EB.AIR/125 13 March 2014.

11.2 ANALYSIS OF KEY TRENDS BY POLLUTANT

Decree of President of the Republic 24 May 1988, n. 203. Attuazione delle direttive CEE numeri 80/779, 82/884, 84/360 e 85/203 concernenti norme in materia di qualità dell'aria, relativamente a specifici agenti inquinanti, e di inquinamento prodotto dagli impianti industriali, ai sensi dell'art. 15 della L. 16 aprile 1987, n. 183 (2) (2/a). Gazzetta Ufficiale del 16 giugno 1988, n. 140, S.O.

EC, 1975. Council Directive 75/716/EEC of 24 November 1975 on the approximation of the laws of the Member States relating to the sulphur content of certain liquid fuels.

EC, 1988. Council Directive 88/609/EEC of 24 November 1988 on the limitation of emissions of certain pollutants into the air from large combustion plants.

EC, 1991. Council Directive 91/441/EEC of 26 June 1991 amending Directive 70/220/EEC on the approximation of the laws of the Member States relating to measures to be taken against air pollution by emissions from motor vehicles.

EC, 1994. Directive 94/12/EC of the European Parliament and the Council of 23 March 1994 relating to measures to be taken against air pollution by emissions from motor vehicles and amending Directive 70/220/EEC.

EC, 1996. Council directive 96/61/EC of 24 September 1996 concerning integrated pollution prevention and control. Official Journal L 257 , 10/10/1996 P. 0026 – 0040.

EC, 1997 [a]. Directive 97/68/EC of the European Parliament and of the Council of 16 December 1997 on the approximation of the laws of the Member States relating to measures against the emission of gaseous and particulate pollutants from internal combustion engines to be installed in non-road mobile machinery.

EC, 1997 [b]. Directive 97/24/EC of the European Parliament and of the Council of 17 June 1997 on certain components and characteristics of two or three-wheel motor vehicles.

EC, 1998. Directive 98/69/EC of the European Parliament and of the Council of 13 October 1998 relating to measures to be taken against air pollution by emissions from motor vehicles and amending Council Directive 70/220/EEC.

EC, 1999. Council Directive 1999/13/EC of 11 March 1999 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations.

EC, 2001. National Emission Ceilings Directive: Directive 2001/81/EC.

EC, 2004. Council Directive 2004/26/EC of 21 April 2004 amending Directive 97/68/EC on the approximation of the laws of the Member States relating to measures against the emission of gaseous and particulate pollutants from internal combustion engines to be installed in non-road mobile machinery.

EU, 2016. National Emissions Ceilings Directive 2016/2284/EU.

ISTAT, 2014. I consumi energetici delle famiglie, anno 2013. <http://www.istat.it/it/archivio/142173>

Ministerial Decree 8 May 1989. Limitazione delle emissioni in atmosfera di taluni inquinanti originati dai grandi impianti di combustione. Gazzetta Ufficiale Italiana n. 124 del 30/05/1989.

Ministerial Decree 12 July 1990. Linee Guida per il contenimento delle emissioni inquinanti degli impianti industriali e la fissazione dei valori minimi di emissione. G.U. 30 luglio 1990, n. 176.

Ministerial Decree 16 May 1996, n. 392. Regolamento recante norme tecniche relative alla eliminazione degli olii usati. G. U. n. 173 del 25/07/1996.

Ministerial Decree 21 January 2000, n. 107. Regolamento recante norme tecniche per l'adeguamento degli impianti di deposito di benzina ai fini del controllo delle emissioni dei vapori. G. U. 02/05/2000 n. 100.

MSE, several years [a]. Bilancio Energetico Nazionale (BEN). Ministero delle Attività Produttive, Direzione Generale delle Fonti di Energia ed industrie di base. <http://dgerm.sviluppoeconomico.gov.it/dgerm/ben.asp> .

11.3 ENERGY (NRF SECTOR 1)

ACI, several years. Annuario statistico. Automobile Club d'Italia, Roma. <http://www.aci.it/index.php?id=54>.

AEEG, several years. Qualità del servizio gas. Autorità per l'energia elettrica e il gas.

http://www.autorita.energia.it/it/dati/elenco_dati.htm .

AISCAT, several years. Aiscat in cifre. Data and reports available on website at: http://www.aiscat.it/pubbl_cifre.htm?ck=1&sub=3&idl=4&nome=pubblicazioni&nome_sub=aiscat%20in%20cifre.

ALTROCONSUMO, 2018. Characterization of residential biomass fired appliances emissions with “real-world” combustion cycles. Technical Report (rev.2). Laboratorio Energia e Ambiente Piacenza. Società Consortile partecipata dal Politecnico di Milano. Piacenza December, 3rd 2018.

ANCMA, several years. Data available on website at: <http://www.ancma.it/statistiche/-/statistics>.

ANPA, 2001. Redazione di inventari nazionali delle emissioni in atmosfera nei settori del trasporto aereo e marittimo e delle emissioni biogeniche. Rapporto finale. Gennaio 2001

ARPA, 2007 – Stima dei consumi di legna da ardere ed uso domestico in Italia. Ricerca commissionata da APAT e ARPA Lombardia, Rapporto finale, marzo 2007.

ASSOTERMICA, several years. Studio Statistico, <http://www.anima.it/contenuti/10670/studi-di-mercato>.

Berdowski J.J.M., Baas J., Bloos J.P.J., Visschedijk A.J.H., Zandveld P.Y.J., 1997. The European Emission Inventory of Heavy Metals and Persistent Organic Pollutants for 1990. TNO Institute of Environmental Sciences, Energy Research and Process Innovation, UBA-FB report 104 02 672/03.

CESI, 2005. Caratterizzazione delle emissioni di caldaie residenziali, CESI, 2005.

CONFETRA, several years. Il trasporto merci su strada in Italia. Data and reports available on website at: <http://www.confetra.it/it/centrostudi/statistiche.htm>.

Decree of President of the Republic 24 May 1988, n. 203. Attuazione delle direttive CEE numeri 80/779, 82/884, 84/360 e 85/203 concernenti norme in materia di qualità dell'aria, relativamente a specifici agenti inquinanti, e di inquinamento prodotto dagli impianti industriali, ai sensi dell'art. 15 della L. 16 aprile 1987, n. 183 (2) (2/a). Gazzetta Ufficiale del 16 giugno 1988, n. 140, S.O.

EC, 2004. Council Directive 2004/26/EC of 21 April 2004 amending Directive 97/68/EC on the approximation of the laws of the Member States relating to measures against the emission of gaseous and particulate pollutants from internal combustion engines to be installed in non-road mobile machinery.

EDISON, several years. Rendiconto ambientale e della sicurezza.

EEA, 2019. Third phase of review of national air pollution inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report, 22 November 2019.

EEA, 2020. Final Review Report, 2020 Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 20 November 2020. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.

EEA, 2021. Final Review Report, 2021 Review of National Air Pollutant Emission Inventory Data 2021 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 29 October 2021. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.

EEA, several years, Monitoring CO₂ emissions from passenger cars and vans, EEA technical reports.

EMEP/CORINAIR, 2007. Atmospheric Emission Inventory Guidebook. Technical report No 16/2007.

EMEP/EEA, 2013. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 12/2013.

EMEP/EEA, 2016. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 21/2016.

EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 13/2019.

EMISIA SA, 2021. COPERT 5 v 5.5.1, Computer programme to calculate emissions from road transport, September 2021. <http://www.emisia.com/copert/> .

ENAC/MIT, several years. Annuario Statistico. Ente Nazionale per l'Aviazione Civile, Ministero delle Infrastrutture e dei Trasporti.

ENEA, several years. Rapporto Energia Ambiente. Ente per le Nuove tecnologie, l'Energia e l'Ambiente, Roma.

ENEA, 2000. Il riciclaggio delle batterie al piombo-acido esauste.

ENEA-AIB-MATT, 2002. "Valutazione delle emissioni di inquinanti organici persistenti da parte dell'industria metallurgica secondaria".

ENEA-federAmbiente, 2012. Rapporto sul recupero energetico da rifiuti urbani in Italia. 3° ed.

ENI, several years. Health Safety Environment report. ENI.

EUROCONTROL, several years. EUROCONTROL Fuel and Emissions Inventory. Personal Communication to EU Member States. Last communication November 2017.

Gerardi V., Perrella G., 2001 - I consumi energetici di biomasse nel settore residenziale in Italia nel 1999. ENEA, Ente Nazionale per l'Energia e l'Ambiente, 2001, ENEA RT/ERG/2001/07.

Giordano R., 2007. Trasporto merci: criticità attuali e potenziali sviluppi nel contesto europeo. National road transporters central committee.

INNOVHUB, 2016. Studio comparativo sulle emissioni di apparecchi a gas, GPL, gasolio e pellet. INNOVHUB, Stazioni sperimentali per l'industria.

INNOVHUB, 2018. Fuel consumption, regulated and unregulated exhaust emission tests on five Euro 6 b/c bifuel LPG passenger cars.

INNOVHUB, 2021. Final report on emissions from biomass combustion for heating.

INNOVHUB, several years. Report on the physico-chemical characterization of fossil fuels used in Italy.

IPCC, 1997. *Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories*. Three volumes: Reference Manual, Reporting Manual, Reporting Guidelines and Workbook. IPCC/OECD/IEA. IPCC WG1 Technical Support Unit, Hadley Centre, Meteorological Centre, Meteorological Office, Bracknell, UK.

IPCC, 2000. Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan.

IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan.

ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009. URL: http://www.apat.gov.it/site/IT/APAT/Pubblicazioni/Rapporti/Documento/rapporti_92_2009.html.

ISPRA, 2021. Update of estimates of atmospheric emissions from integrated plant for non ferrous metals production. Technical note n.1/2021.

ISPRA, 2022[a]. Italian Greenhouse Gas Inventory 1990-2020. National Inventory Report 2022. March 2022, <http://emissioni.sina.isprambiente.it/inventario-nazionale/>.

ISPRA, 2022[b]. Quality Assurance/Quality Control plan for the Italian Emission Inventory. Year 2022. March 2022, <http://emissioni.sina.isprambiente.it/inventario-nazionale/>.

ISPRA, several years. Fuel Quality Monitoring Annual Report.

ISTAT, 2009. Personal communication.

ISTAT, 2014. I consumi energetici delle famiglie, 2013. Nota metodologica. Istituto Nazionale di Statistica www.istat.it.

ISTAT, several years [a]. Annuario Statistico Italiano. Istituto Nazionale di Statistica

ISTAT, several years [b]. Trasporto merci su strada. Istituto Nazionale di Statistica. http://www3.istat.it/dati/dataset/20110729_00/.

JRC, 2013. JRC reference report. Best Available Techniques (BAT) Reference Document for Iron and Steel Production. Industrial Emissions Directive 2010/75/EU (Integrated Pollution Prevention and Control). Report EUR 25521 EN.

Katsis P., Mellios G., Ntziachristos L., 2012. Description of new elements in COPERT 4 v 10.0, December 2012.

Kouridis C., Gkatzoflias D., Kioutsioukis I., Ntziachristos L., Pastorello C., Dilara P., 2010. Uncertainty Estimates and Guidance for Road Transport Emission Calculations, European Commission, Joint Research Centre, Institute for Environment and Sustainability, 2010, <http://www.emisia.com/docs/COPERT%20uncertainty.pdf>.

Ministerial Decree 12 July 1990. Linee Guida per il contenimento delle emissioni inquinanti degli impianti industriali e la fissazione dei valori minimi di emissione. G.U. 30 luglio 1990, n. 176.

MIT, several years. Conto Nazionale delle Infrastrutture e dei Trasporti (CNT). Ministero delle Infrastrutture e dei Trasporti. http://www.mit.gov.it/mit/site.php?o=vc&lm=2&id_cat=148.

MSE, several years [a]. Bilancio Energetico Nazionale (BEN). Ministero delle Attività Produttive, Direzione Generale delle Fonti di Energia ed industrie di base. <http://dgerm.sviluppoeconomico.gov.it/dgerm/ben.asp>.

MSE, several years [b]. Bollettino Petrolifero Trimestrale (BPT). Ministero dello sviluppo economico. <http://dgerm.sviluppoeconomico.gov.it/dgerm/bollettino.asp>.

Riva A., 1997. Methodology for methane emission inventory from SNAM transmission system. Snam Spa Italy.

Romano D., Gaudioso D., De Lauretis R., 1999. Aircraft Emissions: a comparison of methodologies based on different data availability. Environmental Monitoring and Assessment. Vol. 56 pp. 51-74.

SCENARI/ISPRA, 2013. Indagine sull'uso e la disponibilità delle biomasse in Italia. 2013.

Sempos I., 2018. Note on fossil carbon content in biofuels. IPCC Working Group I, 10 October 2018.

SNAM, several years. Bilancio di sostenibilità.

SSC, 2012. Final report on emissions from biomass combustion for heating. Stazione Sperimentale dei Combustibili. Prot. ENEA/2009/34883/APU-UGA March 2012.

TECHNE, 2009. Stima delle emissioni in atmosfera nel settore del trasporto aereo e marittimo. Final report. TECHNE Consulting, March 2009.

TERNA, several years. Dati statistici sugli impianti e la produzione di energia elettrica in Italia, Gestore Rete Trasmissione Nazionale. www.terna.it.

Trozzi C., Vaccaro R., De Lauretis R., Romano D., 2002 [a]. Air pollutant emissions estimate from global air traffic in airport and in cruise: methodology and case study. Presented at Transport and Air Pollution 2002.

Trozzi C., Vaccaro R., De Lauretis R., 2002 [b]. Air pollutant emissions estimate from global ship traffic in port and in cruise: methodology and case study. Presented at Transport and Air Pollution 2002.

UCINA, several years. La nautica in cifre. Analisi annuale del mercato. Unione Nazionale Cantieri Industrie Nautiche ed Affini.

UP, several years. Previsioni di domanda energetica e petrolifera in Italia. Unione Petrolifera.

11.4 IPPU - INDUSTRIAL PROCESSES (NRF SECTOR 2)

APAT, 2003. Il ciclo industriale dell'acciaio da forno elettrico. Agenzia per la Protezione dell'Ambiente e per i servizi tecnici, Rapporti 38/2003.

AITEC, 2017. Relazione annuale 2017. Associazione italiana tecnico economica del cemento. www.aitecweb.com.

CORINAIR, 1994. Default emission factors handbook. EUR 12586/2, Belgium.

EEA, 2017 [a]. Final Review Report. 2017 Comprehensive Technical Review of National Emission Inventories pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Italy 30 November 2017.

EEA, 2018. Second phase of review of national air pollution emission inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report 2018, 30 November 2018.

EEA, 2019. Third phase of review of national air pollution inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report, 22 November 2019.

EEA, 2020. Final Review Report 2020. Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive) Service Contract No. 070201/2019/8159797/SER/ENV.C.3. Italy, 20 November 2020.

EMEP/CORINAIR, 2006. Atmospheric Emission Inventory Guidebook. Technical report No 11/2006.

EMEP/CORINAIR, 2007. Atmospheric Emission Inventory Guidebook. Technical report No 16/2007.

EMEP/EEA, 2013. Air Pollutant Emission Inventory Guidebook. Technical report n. 12/2013.

EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 13/2019.

ENEA-AIB-MATT, 2002. "Valutazione delle emissioni di inquinanti organici persistenti da parte dell'industria metallurgica secondaria".

EUROCHLOR, 1998. Mercury process for making chlorine.

EUROCHLOR, 2001. Reduction of Mercury Emissions from the West European Chlor-Alkali Industry, June 2001 available on the web <http://www.chem.unep.ch/mercury/2001-ngo-sub/eurochlor/sub1ngoatt8.pdf>.

FEDERACCAI, several years. La siderurgia in cifre. Federazione Imprese Siderurgiche Italiane.

ILVA, 1997. Personal communication.

IPCC, 1997. *Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories*. Three volumes: Reference Manual, Reporting Manual, Reporting Guidelines and Workbook.

IPCC, 2000. Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan.

IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan.

IPPC, 2001. Best Available Techniques Reference Document on the Production of Iron and Steel. Integrated Pollution Prevention and Control. European Commission. December 2001.

ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009. <http://www.isprambiente.gov.it/it/pubblicazioni/rapporti/la-disaggregazione-a-livello-provinciale>.

ISPRA, 2021. Update of estimates of atmospheric emissions from integrated plant for non ferrous metals production. Technical note n.1/2021.

ISTAT, several years. Annuario Statistico Italiano. Istituto Nazionale di Statistica.

Legambiente, 2007. Lo stato dell'arte sulle riconversioni degli impianti cloro soda in Italia, Ottobre 2007 available on the web: http://risorse.legambiente.it/docs/Lo_stato_dell_arte_sulle_riconversioni_degli_impianti_cloro-soda_in_Italia_ott_2007.0000001526.pdf

MATTM, 2011. Autorizzazione integrata ambientale per l'esercizio dello stabilimento siderurgico della società ILVA s.p.a. ubicato nel comune di Taranto. IPPC permit.

MSE, several years. Statistiche produzione cementi. <http://www.sviluppoeconomico.gov.it/index.php/it/per-i-media/statistiche/2009708-statistiche-produzione-cementi>.

TNO, 1992. Emission factors manual PARCOM – ATMOS. Emission factors for air pollutants 1992.

11.5 IPPU - SOLVENT AND OTHER PRODUCT USE (NRF SECTOR 2)

ACI, several years. Annuario statistico. Automobile Club d'Italia, Roma. <http://www.aci.it/index.php?id=54>.

AIA, several years [a]. Personal Communication. Associazione Italiana Aerosol.

AIA, several years [b]. Relazioni annuali sulla produzione italiana aerosol. Associazione Italiana Aerosol.

Assocasa, several years. Personal Communication.

AVISA, several years. Personal Communication.

CEPE, 1999. European guidelines for car repairing.

EC, 1999. Council Directive 1999/13/EC of 11 March 1999 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations.

EC, 2002. Screening study to identify reduction in VOC emissions due to the restrictions in the VOC content of products. Final Report of the European Commission, February 2002.

EC, 2004. Directive 2004/42/ce of the european parliament and of the council of 21 April 2004 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products and amending Directive 1999/13/EC.

EEA, 2019. Third phase of review of national air pollution inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report, 22 November 2019.

EEA, 2020. Final Review Report 2020. Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive) Service Contract No. 070201/2019/8159797/SER/ENV.C.3. Italy, 20 November 2020.

-
- EMEP/CORINAIR, 2007. Atmospheric Emission Inventory Guidebook. Technical report No 16/2007.
- EMEP/EEA, 2016. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 21/2016.
- EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. Technical report n. 13/2019.
- ENEA, 1997. Piano nazionale di tutela della qualità dell'aria. Technical report produced for the Italian Ministry of Environment in the framework of a national project on air quality issue.
- ENEA/USLRMA, 1995. Lavanderie a secco.
- FAO, several years. Food balance. <http://faostat.fao.org>.
- Federchimica, several years. Personal Communication.
- FIAT, several years. Rendiconto Ambientale. Gruppo Fiat.
- GIADA, 2006. Progetto Giada and Personal Communication. ARPA Veneto – Provincia di Vicenza.
- ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009. URL: http://www.apat.gov.it/site/IT/APAT/Pubblicazioni/Rapporti/Documento/rapporti_92_2009.html
- ISTAT, several years [a]. Annuario Statistico Italiano.
- ISTAT, several years [b]. Bollettino mensile di statistica.
- ISTAT, several years [c]. Statistica annuale della produzione industriale. <http://www.istat.it/it/archivio/73150>
- MICA, 1999. L'industria Italiana delle vernici. Ministero dell'Industria del Commercio e dell'Artigianato. Dicembre 1999.
- Offredi P., several years. Professione Verniciatore del Legno. Personal communication.
- Regione Campania, 2005. Inventario regionale delle emissioni di inquinanti dell'aria della Regione Campania, marzo 2005.
- Regione Toscana, 2001. Inventario regionale delle sorgenti di emissione in aria ambiente, febbraio 2001.
- Techne, 1998. Personal communication.
- Techne, 2004. Progetto MeditAiraneo. Rassegna dei fattori di emissione nazionali ed internazionali relativamente al settore solventi. Rapporto Finale, novembre 2004.
- Techne, 2008. Fattori di emissione per l'utilizzo di solventi. Rapporto Finale, marzo 2008.
- UNIPRO, several years. Rapporto Annuale - Consumi cosmetici in Italia.
- Vetrella G., 1994. Strategie ottimali per la riduzione delle emissioni di composti organici volatili. Thesis in Statistics.

11.6 AGRICULTURE (NRF SECTOR 3)

- APAT, 2005. Methodologies used in Italy for the estimation of air emission in the agriculture sector. Technical report 64/2005. Rome - Italy. <https://www.isprambiente.gov.it/it/pubblicazioni/rapporti/methodologies-used-in-italy-for-the-estimation-of>
- Bassanino M., Sacco D., Zavattaro L., Grignania C., 2011. Nutrient balance as a sustainability indicator of different agro-environments in Italy. *Ecol Indic* 11(2): 715-723
- Bechini L., Castoldi N., 2009. On-farm monitoring of economic and environmental performances of cropping systems: Results of a 2-year study at the field scale in northern Italy. *Ecol Indic* 9: 1096-1113
- Berdowski J.J.M., Baas J, Bloos J.P.J., Visschedijk A.J.H., Zandveld P.Y.J., 1997. The European Atmospheric Emission Inventory for Heavy Metals and Persistent Organic Pollutants. Umweltforschungsplan des Bundesministers für Umwelt, Naturschutz und Reaktorsicherheit Luftreinhaltung. Forschungsbericht 104 02 672/03. TNO, Apeldoorn, The Netherlands
- Bittman S. et al, 2014. Bittman, S., Dedina, M., Howard C.M., Oenema, O., Sutton, M.A., (eds), 2014. Options for Ammonia Mitigation: Guidance from the UNECE Task Force on Reactive Nitrogen, Centre for Ecology and Hydrology, Edinburgh, UK www.clrtap-tfrn.org
- Bonazzi G., Crovetto M., Della Casa G., Schiavon S., Sirri F., 2005. Evaluation of Nitrogen and Phosphorus in Livestock manure: Southern Europe (Italy), presentato al Workshop: Nutrients in livestock manure, Bruxelles, 14 February 2005

-
- Cóndor R.D., Di Cristofaro E., De Lauretis R., 2008. Agricoltura: inventario nazionale delle emissioni e disaggregazione provinciale. Istituto superiore per la protezione e la ricerca ambientale, ISPRA Rapporto tecnico 85/2008. Roma, Italia. <https://www.isprambiente.gov.it/it/archivio/notizie-e-novita-normative/notizie-ispra/anno-2009/rapporto-agricoltura>
- Cóndor R.D., 2011. Agricoltura: emissioni nazionali in atmosfera dal 1990 al 2009. Istituto superiore per la protezione e la ricerca ambientale (ISPRA). Rapporto ISPRA 140/2011. Roma, Italia. <https://www.isprambiente.gov.it/it/pubblicazioni/rapporti/agricoltura-emissioni-nazionali-in-atmosfera-dal>
- Cóndor R.D., Valli L., 2011. Emissioni nazionali di ammoniaca e scenari emissivi derivanti dalla fase di spandimento agronomico e all'uso dei fertilizzanti azotati in Italia. e-book edited by the Centro Ricerche Produzioni Animali (CRPA). http://www.crpa.it/media/documents/crpa_wwww/Pubblicazi/E-book/Ammoniaca2011/EmissioniAmmoniaca.pdf
- CREA, 2021. Accordo di collaborazione (ex art. 15 L. 241/90) tra ISPRA e CREA - Rapporto tecnico. Valutazione delle emissioni connesse all'uso dei fertilizzanti azotati nell'ambito della direttiva NEC. Novembre 2021
- CRPA, 1997. Piani Regionali di Risanamento e tutela della qualità dell'aria. Quadro delle azioni degli enti locali per il settore zootecnico delle aree padane. Relazione di dettaglio sulla metodologia adottata per la quantificazione delle emissioni di ammoniaca. Febbraio 1997
- CRPA, 2000. Aggiornamento dell'inventario delle emissioni in atmosfera di ammoniaca, metano e protossido di azoto dal comparto agricolo. Centro Ricerche Produzioni Animali. Gennaio 2000.
- CRPA, 2006[a]. Progetto MeditAIRaneo: settore Agricoltura. Relazione finale. Technical report on the framework of the MeditAIRaneo project for the Agriculture sector, Reggio Emilia - Italy
- CRPA, 2006[b]. Predisposizione di scenari di emissione finalizzati alla progettazione di interventi per la riduzione delle emissioni nazionali di ammoniaca ed alla valutazione di misure e di progetti per la tutela della qualità dell'aria a livello regionale. Final report. Reggio Emilia - Italy
- CRPA, 2010[a]. Personal communication - experts Laura Valli and Maria Teresa Pacchioli from the Research Centre on Animal Production (expert consultation on N excretion and national production systems). Reggio Emilia, Italy
- CRPA, 2010[b]. Valutazione dell'entità delle emissioni ammoniacali derivanti dall'applicazione al suolo dei fertilizzanti, delle loro possibilità di riduzione e individuazione degli elementi per un monitoraggio statistico delle tecniche di applicazione utilizzate. Report. Reggio Emilia - Italy
- CRPA, 2018. Studio per la valutazione degli effetti sulle emissioni delle trasformazioni in corso nel settore degli allevamenti. Report. Reggio Emilia - Italy
- EC, 1986. Council Directive 86/278/EC. Council Directive 86/278/EC of 12 June 1986 on the protection of the environment, and in particular of the soil, when sewage sludge is used in agriculture. Official Journal of the European Communities 4 July 1986
- EC, 2003. Integrated Pollution Prevention and Control (IPPC). Reference Document on Best Available Techniques for Intensive Rearing of Poultry and Pigs. European Commission, July 2003.
- EEA, 2016. Analysis of key trends and drivers in greenhouse gas emissions in the EU between 1990 and 2014. Technical paper, June 2016. <http://www.eea.europa.eu/publications/analysis-of-key-trends-ghg/>
- EEA, 2019. Third phase of review of national air pollution inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report, 22 November 2019.
- EEA, 2020. Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive). Final Review Report 2020, 20 November 2020.
- EMEP/CORINAIR, 2006. Atmospheric Emission Inventory Guidebook. Technical report No 11/2006.
- EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. Technical report n. 13/2019.
- ENEA, 2003. Emissione di ammoniaca e di composti ad effetto serra dagli allevamenti di galline ovaiole. Fattori di emissione e tecniche di riduzione. RT/2003/65/PROT. Italy.
- ENEA, 2006. Valutazione della possibilità di sostituzione dell'urea con altri fertilizzanti azotati. Final report. Rome, Italy
- EPA, 1995. AP-42 Compilation of Air Emission Factors, 5th edition January 1995. <https://www.epa.gov/air-emissions-factors-and-quantification/ap-42-compilation-air-emission-factors>
- Giardini L., 1983. Agronomia Generale, Patron, Bologna

Gislon G., Colombini S., Borreani G., Crovetto G. M., Sandrucci A., Galassi G., Tabacco E., Rapetti L., 2020. Milk production, methane emissions, nitrogen, and energy balance of cows fed diets based on different forage systems. *J. Dairy Sci.* 103:8048–8061 <https://doi.org/10.3168/jds.2019-18134>

GU, Gazzetta Ufficiale della Repubblica Italiana, 2006. Criteri e norme tecniche generali per la disciplina regionale dell'utilizzazione agronomica degli effluenti di allevamento e di acque reflue di cui all'articolo 38 del decreto legislativo 11 maggio 1999 N. 152. G.U. n. 109 del 12/05/06 - Suppl. Ordinario n.120. Ministero delle Politiche Agricole e Forestali. Italy. <http://www.gazzettaufficiale.it/>

GU, Gazzetta Ufficiale della Repubblica Italiana, 2016. Criteri e norme tecniche generali per la disciplina regionale dell'utilizzazione agronomica degli effluenti di allevamento e delle acque reflue, nonché per la produzione e l'utilizzazione agronomica del digestato. G.U. n. 90 del 18/04/16 - Suppl. Ordinario n. 9. Ministero delle Politiche Agricole Alimentari e Forestali. Italy. <http://www.gazzettaufficiale.it/>

IPCC, 1997. *Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories*. Three volumes: Reference Manual, Reporting Manual, Reporting Guidelines and Workbook

IPCC, 2000. Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan

IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan

ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009. <https://www.isprambiente.gov.it/it/pubblicazioni/rapporti/la-disaggregazione-a-livello-provinciale>

ISPRA, 2021. Database della disaggregazione a livello provinciale dell'Inventario nazionale delle emissioni:1990-1995-2000-2005-2010-2015-2019. Istituto Superiore per la Protezione e la Ricerca Ambientale, ISPRA. <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>

ISPRA, several years [a]. Italian Greenhouse Gas Inventory - National Inventory Report. <http://emissioni.sina.isprambiente.it/inventario-nazionale/>

ISPRA, several years [b]. Quality Assurance/Quality Control plan for the Italian Emission Inventory, <http://emissioni.sina.isprambiente.it/inventario-nazionale/>

ISPRA, several years [c]. Annuario dei dati ambientali. <http://annuario.isprambiente.it/>

ISPRA, several years [d]. Serie storiche delle emissioni nazionali di inquinanti atmosferici, Rete del Sistema Informativo Nazionale Ambientale - SINANET. <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>

Jenkins B.M. (1996a). 'Atmospheric Pollutant Emission Factors from Open Burning of Agricultural and Forest Biomass by Wind Tunnel Simulations'. Final report (3 Vols.). CARB Project A932-126, California Air Resources Board, Sacramento, California.

MATTM, several years [b]. Personal communication: E-mail request for sewage sludge applied to agricultural soils in Italy. Ministero dell'Ambiente e della Tutela del Territorio e del Mare, Roma –Italia

Nicholson F.A., Chambers B.J., Walker A.W., 2004. Ammonia emissions from broiler litter and laying hen manure management system, *Biosystems Engineering* (2004) 89 (2), 175-185

Perelli, M., 2007. Prezzi dei prodotti agricoli e fertilizzazione. *Fertilizzanti Maggio 2007*. Anno IX N3. 10-13pp

Regione Emilia-Romagna, Servizio sviluppo sistema agroalimentare, (2001) L. R. 28/98 – p.s.a. 2000 - n. Prog. 3 tab. D - Tecniche di riduzione delle emissioni in atmosfera originate dagli allevamenti zootecnici, a cura di CRPA, Settembre 2001

Regione Emilia Romagna, 2004 L. R. 28/98 – P.S.A. 2001 - N. PROG. 3 TAB. B3 - Bilancio dell'azoto nelle specie di interesse zootecnico, Relazione finale, a cura di C.R.P.A., September 2004, Reggio Emilia, Italy

Xiccato G., Schiavon S., Gallo L., Bailoni L., Bittante G., 2005. Nitrogen excretion in dairy cow, beef and veal cattle, pig, and rabbit farms in Northern Italy. *Ital. J.Anim.Sci.* Vol. 4 (Suppl.), 103-111.

11.7 WASTE (NRF SECTOR 5)

Aasestad, 2007. The Norwegian Emission Inventory 2007. Documentation of methodologies for estimating emissions of greenhouse gases and long-range transboundary air pollutants.

-
- AMA-Comune di Roma, 1996. Nuovo impianto per l'incenerimento dei rifiuti ospedalieri. Rapporto AMA.
- ANPA, 1998. Il sistema ANPA di contabilità dei rifiuti, prime elaborazioni dei dati. Agenzia Nazionale per la Protezione dell'Ambiente.
- APAT-ONR, several years. Rapporto Rifiuti. Agenzia per la Protezione dell'Ambiente e per i servizi Tecnici.
- AUSITRA-Assoambiente, 1995. Impianti di trattamento dei rifiuti solidi urbani e assimilabili. Indagine a cura di Merzagora W., Ferrari S.P.
- Borgioli E., 1981. Nutrizione e alimentazione degli animali domestici. Ed Agricole, p. 464.
- CESTAAT, 1988. Impieghi dei sottoprodotti agricoli ed agroindustriali, Vol. 1. Centro Studi sull'Agricoltura, l'Ambiente e il Territorio, edizione fuori commercio, p. 311.
- CREA, 2017. Fornitura dati meteo-climatici georeferenziati nell'ambito della collaborazione CREA-AA/ISPRA. CREA - Consiglio per la Ricerca in Agricoltura e l'Analisi dell'Economia Agraria Centro di ricerca Agricoltura e Ambiente (CREA-AA), delivery data mail 19/10/2017.
- CRPA, 2018. Studio per la valutazione degli effetti sulle emissioni delle trasformazioni in corso nel settore degli allevamenti. Report. Reggio Emilia – Italy
- De Stefanis P., 1999. Personal communication.
- De Stefanis P., 2012. Personal communication (mail 16 November 2012)
- EC, 1999. Council Directive 1999/31/EC. Council Directive 99/31/EC of 26 April 1999 on the landfill of waste. Official Journal of the European Communities 16 July 1999.
- EEA, 2017 [a]. Final Review Report. 2017 Comprehensive Technical Review of National Emission Inventories pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Italy 30 November 2017.
- EEA, 2017 [b]. Final Review Report. 2017 annual review of national greenhouse gas inventory data pursuant to Article 19(2) of Regulation (EU) No 525/2013. Italy 30 June 2017.
- EEA, 2018. Second phase of review of national air pollution emission inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report 2018, 30 November 2018.
- EEA, 2019. Third phase of review of national air pollution inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284 or 'NECD') Italy. Draft Review Report 2019, 18 October 2019.
- EEA, 2020. Final Review Report 2020. Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive) Service Contract No. 070201/2019/8159797/SER/ENV.C.3. Italy, 20 November 2020.
- EEA, 2021. Final Review Report, 2021 Review of National Air Pollutant Emission Inventory Data 2021 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 29 October 2021. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.
- EMEP/CORINAIR, 2007. Atmospheric Emission Inventory Guidebook. Technical report No 16/2007.
- EMEP/EEA, 2016. Air Pollutant Emission Inventory Guidebook. Technical report n. 21/2016.
- EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. Technical report n. 13/2019.
- ENEA-federAmbiente, 2012. Rapporto sul recupero energetico da rifiuti urbani in Italia. 3° ed.
- Favoino E., Cortellini L., 2001. Composting and biological treatment in southern European countries: an overview. Conference Proceedings Soil and Biowaste in Southern Europe. Rome 18-19 January, 2001.
- Favoino E., Girò F., 2001. An assessment of effective, optimised schemes for source separation of organic waste in Mediterranean districts. Conference Proceedings Soil and Biowaste in Southern Europe. Rome 18-19 January, 2001.
- FEDERAMBIENTE, 1998. Impianti di smaltimento: analisi sui termocombustori RSU – prima edizione. Indagine a cura di Motawi A.
- Finn L., Spencer R., 1997. Managing biofilters for consistent odor and VOC treatment. Biocycle, January 1997 Vol. 38 Iss.1.
- Gaudio et al., 1993. Emissioni in atmosfera dalle discariche di rifiuti in Italia. RS, Rifiuti Solidi vol. VII n. 5, Sept.-Oct. 1993.

IIASA, 2004. Interim Report IR-04-079. Primary Emissions of Submicron and Carbonaceous Particles in Europe and the Potential for their Control.

IPCC, 1997. *Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories*. Three volumes: Reference Manual, Reporting Manual, Reporting Guidelines and Workbook. IPCC/OECD/IEA. IPCC WG1 Technical Support Unit, Hadley Centre, Meteorological Centre, Meteorological Office, Bracknell, UK.

IPCC, 2000. Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan.

ISPRA, 2022[a]. Italian Greenhouse Gas Inventory 1990-2020. National Inventory Report 2022. March 2022, <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/> .

ISPRA, several years [a]. Rapporto Rifiuti. Istituto Superiore per la Protezione e la Ricerca Ambientale. <http://www.isprambiente.gov.it/it/pubblicazioni/rapporti>

ISTAT, several years [a]. Statistiche dell'agricoltura, zootecnia e mezzi di produzione - Annuari (1990-1993), Istituto Nazionale di Statistica.

ISTAT, several years [b]. Statistiche dell'agricoltura - Annuari (1994-2000), Istituto Nazionale di Statistica.

ISTAT, 2015. Censimento delle acque per uso civile. Istituto nazionale di statistica, also available at website <http://www.istat.it>.

ISTAT, 2017[a]. Dati congiunturali sulle coltivazioni 2014. Istituto Nazionale di Statistica. <http://agri.istat.it/jsp/Introduzione.jsp>.

ISTAT, 2017[b]. Personal communication with R. Moro: E-mail request for last updated information on agricultural surface and production, year 2015. Istituto Nazionale di Statistica.

MATTM, several years [a]. RSA - Rapporto sullo stato dell'ambiente 1989, 1992, 1997, 2001. Ministero dell'Ambiente e della Tutela del Territorio e del Mare.

Ministerial Decree 19 November 1997, n. 503. Regolamento recante norme per l'attuazione delle Direttive 89/369/CEE e 89/429/CEE concernenti la prevenzione dell'inquinamento atmosferico provocato dagli impianti di incenerimento dei rifiuti urbani e la disciplina delle emissioni e delle condizioni di combustione degli impianti di incenerimento di rifiuti urbani, di rifiuti speciali non pericolosi, nonché di taluni rifiuti sanitari. G.U. 29 gennaio 1998, n. 23.

Ministerial Decree 12 July 1990. Linee Guida per il contenimento delle emissioni inquinanti degli impianti industriali e la fissazione dei valori minimi di emissione. G.U. 30 luglio 1990, n. 176.

Pastorelli et al., 2001. Sviluppo di fattori di emissione da inceneritori di rifiuti urbani lombardi e loro applicazione all'inventario nazionale delle diossine. *Ingegneria Ambientale*, ANNO XXX N.1 January 2001.

SEFIT, several years. Personal Communication with Daniele Fogli: E-mail request for activity data regarding cremation of corpses in Italy.

SEFIT, 2015. Emissioni inquinanti in atmosfera per i crematori italiani. Indagine conoscitiva ed elaborazione dati. Novembre 2015.

Tchobanoglous G. et al., 1993. Tchobanoglous G., Theisen H., Vigil A. *Integrated Waste Management*, McGraw-Hill, 1993.

US EPA, 1990. Air Emissions Species Manual, vol. I: Volatile Organic Compound Species Profiles, Second Edition. EPA-450/2-90-001a (United States Environmental Protection Agency – Office of Air Quality Planning and Standards, Research Triangle Park, NC 27711), January 1990.

VV.FF., several years. Annually statistics of fire service in Italy, several years.

11.8 RECALCULATIONS AND IMPROVEMENTS

EEA, 2021. Final Review Report, 2021 Review of National Air Pollutant Emission Inventory Data 2021 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 29 October 2021. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.

EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. Technical report n. 13/2019.

EMISIA SA, 2021. COPERT 5 v 5.5.1, Computer programme to calculate emissions from road transport, September 2021. <http://www.emisia.com/copert/>.

ISPRA, 2022[b]. Quality Assurance/Quality Control plan for the Italian Emission Inventory. Year 2022. March 2022, <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/> .

11.9 PROJECTIONS

Amann, M., Bertok, J., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Klimont, Z., Nguyen, B., Posch, M., Rafaj, P., Sandler, R., Schöpp, W., Wagner, F., Winiwarter, W., 2011. Cost-effective control of air quality and greenhouse gases in Europe: modelling and policy applications. *Environmental Modeling Software*, 26, 1489–1501.

Briganti, G., Calori, G., Cappelletti, A., Ciancarella, L., D'Isidoro, M., Finardi, S., Vitali, L., 2011. Determination of multi-year atmospheric transfer Matrices for GAINS-Italy model. In: *High Performance Computing on CRESCO Infrastructure: Research Activities and Results*, vol. 2010. pp. 45–51 Report ENEA Cresco, ISBN 978-88-8286-268-8.

D'Elia, I., Peschi, E., 2013. Lo scenario emissivo nazionale nella negoziazione internazionale. ENEA Technical Report, RT/2013/10/ENEA (in Italian). <http://openarchive.enea.it/handle/10840/4505>.

D'Elia, I., Peschi, E., 2016. How National integrated air quality models can be used in defining environmental policies: the revision of the NEC directive. ENEA Technical Report, RT/2016/30/ENEA. <http://openarchive.enea.it/handle/10840/8153> .

D.Lgs., 2010. Decreto Legislativo 13 agosto 2010, n. 155. Attuazione della direttiva 2008/50/CE relativa alla qualità dell'aria ambiente e per un'aria più pulita in Europa (GU n.216 del 15-9-2010 - Suppl. Ordinario n. 217).

D.Lgs., 2018. Decreto Legislativo 30 maggio 2018, n. 81. Attuazione della direttiva (UE) 2016/2284 del Parlamento europeo e del Consiglio, del 14 dicembre 2016, concernente la riduzione delle emissioni nazionali di determinati inquinanti atmosferici, che modifica la direttiva 2003/35/CE e abroga la direttiva 2001/81/CE (GU n. 151 del 2-7-2018).

EC, 2008. Directive 2008/50/EC of the European parliament and of the council of 21 may 2008 on ambient air quality and a cleaner air for Europe. *Official Journal of the European Union L*. 152 of 11.06.2008.

EU, 2016. Directive (EU) 2016/2284 of the European Parliament and of the Council of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC. *Official Journal of the European Union L*. 344/1 of 17.12.2016.

IIR, 2021. Italian Emission Inventory 1990-2019. Informative Inventory Report 2021. ISPRA Technical Report 342/2021. ISBN 978-88-448-1047-4

Loulou, R., Goldstein, G., Noble, K., 2004. Documentation for the MARKAL Family of Models. ETSAP.

Loulou, R., Remne, U., Kanudia, A., Lehtila, A., Goldstein, G., 2005. Documentation for the TIMES Model - PART I 1–78. <https://iea-etsap.org/index.php/documentation>

Mircea, M., Ciancarella, L., Briganti, G., Calori, G., Cappelletti, A., Cionni, I., Costa, M., Cremona, G., D'Isidoro, M., Finardi, S., Pace, G., Piersanti, A., Righini, G., Silibello, C., Vitali, L., Zanini, G., 2014. Assessment of the AMS-MINNI system capabilities to predict air quality over Italy for the calendar year 2005. *Atmospheric Environment*, 84, 178–188, ISSN 1352-2310, <http://dx.doi.org/10.1016/j.atmosenv.2013.11.006>

UNECE, 2015. Guidelines for reporting emissions and projections data under the Convention on Long-range Transboundary Air Pollution. ECE/EB.AIR/128/2015.

<https://www.unece.org/fileadmin/DAM/env/documents/2015/AIR/EB/English.pdf>

11.10 REPORTING OF GRIDDED EMISSIONS AND LPS

CEIP, 2019. Updated documentation of the EMEP gridding system v.2019 for the spatial disaggregation of emission data with a resolution of 0.1°x0.1°(long-lat). Technical report CEIP 06/2019.

EEA, 2020. Final Review Report 2020. Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive) Service Contract No. 070201/2019/8159797/SER/ENV.C.3. Italy, 20 November 2020.

EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 13/2019.

ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009. URL: http://www.apat.gov.it/site/IT/APAT/Pubblicazioni/Rapporti/Documento/rapporti_92_2009.html.

APPENDIX 1 SUMMARY INFORMATION ON CONDENSABLE IN PM

In order to improve atmospheric modelling and support the design of efficient and relevant policy for reducing the levels of air pollutants, emission inventory data need to be complete, accurate and comparable. With this aim, Italy immediately accepted the EMEP proposal on the necessity of accounting for condensable in PM emissions and generally applies these emission factors to all the categories. Of course, for certain categories is not possible to define if the emission factors includes condensable or not, as reported also in the 2019 Guidebook EMEP/EEA, consequently it is hard to fill the following table at category level but it is possible to provide more information. In particular, Italy uses emission factors with condensable for PM emissions from road transport thanks to the Copert model and in domestic and residential heating thanks several studies carried on in the last years about heating appliances, burning wood or other fuels. In particular, as concerns emissions from small combustion, a paper discussed during the 2019 meeting of TFEIP reported some independent verification on these estimates and Italy resulted in a good comparability with estimates of TNO (see figureA6.1).




Figure A6.1 Comparison between TNO and countries estimates

